

UNIT-I
MATRICES

- * Characteristic equation
- * Eigen values & Eigen vectors
- * Properties
- * Cayley - Hamilton theorem
- * Diagonalization of matrices
- * Canonical form
- * Nature of quadratic form
- * Applications: stretching of an elastic membrane.

CHARACTERISTIC EQUATION:-

The equation $|A - \lambda I| = 0$ is said to be the characteristic equation of the transformation or the characteristic equation of the matrix A .

Note:-

- * characteristic eqn for 2×2 matrix is $\lambda^2 - S_1\lambda + S_2 = 0$,

where $S_1 =$ Sum of the main diagonal elements

$$S_2 = |A|$$

- * characteristic eqn for 3×3 matrix is $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$,
where $S_1 =$ Sum of the main diagonal elements

$S_2 =$ Sum of the minors of the main diagonal elements

$$S_3 = |A|$$

Problem:- 1

Find the characteristic eqn of $\begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix}$

Soln:-

$$\text{Let } A = \begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix}$$

The characteristic eqn of

$$A \text{ is } \lambda^2 - S_1\lambda + S_2 = 0$$

$S_1 =$ Sum of the main diagonal elements

$$= 1 + 2 = 3$$

$$S_2 = |A| = 2 - 0 = 2$$

$\therefore$ characteristic equation is $\lambda^2 - 3\lambda + 2 = 0$

Problem: 2

Find the characteristic eqn of

$$\begin{bmatrix} 2 & -3 & 1 \\ 3 & 1 & 3 \\ -5 & 2 & -4 \end{bmatrix}$$

Soln:-

The char. eqn of $A = \begin{bmatrix} 2 & -3 & 1 \\ 3 & 1 & 3 \\ -5 & 2 & -4 \end{bmatrix}$ is $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

S_1 = Sum of the main diagonal elements

$$= 2 + 1 - 4 = -1$$

S_2 = Sum of the minors of the main diagonal elements.

$$= \begin{vmatrix} 1 & 3 \\ 2 & -4 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ -5 & -4 \end{vmatrix} + \begin{vmatrix} 2 & -3 \\ 3 & 1 \end{vmatrix}$$

$$= (-4-6) + (-8+5) + (2+9)$$

$$= 10 - 3 + 11 = -2$$

$$S_3 = \begin{vmatrix} 2 & -3 & 1 \\ 3 & 1 & 3 \\ -5 & 2 & -4 \end{vmatrix}$$

$$= 2(-4-6) + 3(-12+15) + 1(6+5)$$

$$= -20 + 9 + 11$$

$$= 0$$

$\therefore$ Char. eqn is $\lambda^3 + \lambda^2 - 2\lambda = 0$

* Eigen values and Eigen Vectors of a real matrix :-

Procedure :-

Step 1: Find the characteristic equation $|A - \lambda I| = 0$

Step 2: Solving the characteristic eqn, we get Eigen values

Step 3: To find the eigen Vectors, solve $(A - \lambda I)X = 0$

Non Symmetric matrices with non repeated Eigen values :-

Problem: 1

Find the Eigen values & Eigen Vectors of the matrix

$$A = \begin{bmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{bmatrix} \quad (\text{AP-2019})$$

Soln:-

$$\text{Let } A = \begin{bmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{bmatrix}$$

Step 1:-

The char. eqn of A is

$$\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$$

S_1 = Sum of the main diagonal elements

$$= 3 + 5 + 3 = 11$$

S_2 = Sum of the minor of the main diagonal elements.

$$= \begin{vmatrix} 5 & -1 \\ -1 & 3 \end{vmatrix} + \begin{vmatrix} 3 & 1 \\ 1 & 3 \end{vmatrix} + \begin{vmatrix} 3 & -1 \\ -1 & 5 \end{vmatrix}$$

$$= (15-1) + (9-1) + (15-1)$$

$$= 36$$

$$S_3 = \begin{vmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{vmatrix}$$

$$= 3(15-1) + 1(-3+1) + 1(1-5)$$

$$= 42 - 2 - 4 = 36$$

$$\therefore \text{char eqn is } \lambda^3 - 11\lambda^2 + 36\lambda - 36 = 0$$

Step II:-

$$\text{Put } \lambda=2, 2^3 - 11(2^2) + 36(2) - 36 = 0$$

$$2 \left| \begin{array}{ccc|c} 1 & -11 & 36 & -36 \\ 0 & 2 & -18 & 36 \\ \hline 1 & -9 & 18 & 0 \end{array} \right|$$

$$\lambda^2 - 9\lambda + 18 = 0$$

$$(\lambda-6)(\lambda-3) = 0$$

$$\therefore \lambda = 3, 6$$

$\therefore$ The Eigen values are 2, 3, 6.

Step III:-

To find Eigen vector.

$$(A - \lambda I) x = 0$$

$$\begin{pmatrix} 3-\lambda & -1 & 1 \\ -1 & 5-\lambda & -1 \\ 1 & -1 & 3-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\left. \begin{array}{l} (3-\lambda)x_1 - x_2 + x_3 = 0 \\ -x_1 + (5-\lambda)x_2 - x_3 = 0 \\ x_1 - x_2 + (3-\lambda)x_3 = 0 \end{array} \right\} \text{--- (I)}$$

Case(i) put $\lambda=2$ in (I)

$$x_1 - x_2 + x_3 = 0 \quad \text{--- (1)}$$

$$-x_1 + 3x_2 - x_3 = 0 \quad \text{--- (2)}$$

$$x_1 - x_2 + x_3 = 0 \quad \text{--- (3)}$$

Solve (1) & (2)

$$\frac{x_1}{-1} = \frac{x_2}{1} = \frac{x_3}{-1} \quad \left| \begin{array}{cc|c} -1 & 1 & 0 \\ 3 & -1 & 0 \end{array} \right| \quad \left| \begin{array}{cc|c} 1 & 1 & 0 \\ -1 & -1 & 0 \end{array} \right| \quad \left| \begin{array}{cc|c} 1 & -1 & 0 \\ -1 & 3 & 0 \end{array} \right|$$

$$\frac{x_1}{-3} = \frac{x_2}{-1+1} = \frac{x_3}{3-1}$$

$$\frac{x_1}{-2} = \frac{x_2}{0} = \frac{x_3}{2}$$

$$\therefore x_1 = \begin{bmatrix} -2 \\ 0 \\ 2 \end{bmatrix} \Rightarrow x_1 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

Case(ii) put $\lambda=3$ in (I)

$$0x_1 - x_2 + x_3 = 0 \quad \text{--- (4)}$$

$$-x_1 + 2x_2 - x_3 = 0 \quad \text{--- (5)}$$

$$x_1 - x_2 + 0x_3 = 0 \quad \text{--- (6)}$$

Solve (4) & (5)

$$\frac{x_1}{-1} = \frac{x_2}{1} = \frac{x_3}{1} \quad \left| \begin{array}{cc|c} -1 & 1 & 0 \\ 2 & -1 & 0 \end{array} \right| \quad \left| \begin{array}{cc|c} 1 & 0 & 0 \\ -1 & -1 & 0 \end{array} \right| \quad \left| \begin{array}{cc|c} 0 & -1 & 0 \\ -1 & 2 & 0 \end{array} \right|$$

$$\frac{x_1}{1-2} = \frac{x_2}{-1-0} = \frac{x_3}{0-1}$$

$$\frac{x_1}{-1} = \frac{x_2}{-1} = \frac{x_3}{-1}$$

$$\therefore x_2 = \begin{bmatrix} -1 \\ -1 \\ -1 \end{bmatrix}$$

$$\Rightarrow x_2 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Case (iii) put $\lambda = 6$ in ①

$$-3x_1 - x_2 + x_3 = 0 \quad \text{--- ⑦}$$

$$-x_1 - x_2 - x_3 = 0 \quad \text{--- ⑧}$$

$$x_1 - x_2 - 3x_3 = 0 \quad \text{--- ⑨}$$

Solve ⑦ & ⑧

$$\frac{x_1}{-1} = \frac{x_2}{-1} = \frac{x_3}{-3} = \frac{x_3}{-1}$$

$$\begin{bmatrix} -1 & 1 \\ -1 & -1 \end{bmatrix} \quad \begin{bmatrix} 1 & -3 \\ -1 & -1 \end{bmatrix} \quad \begin{bmatrix} -3 & -1 \\ -1 & -1 \end{bmatrix}$$

$$\frac{x_1}{1+1} = \frac{x_2}{-1-3} = \frac{x_3}{3-1}$$

$$\therefore x_3 = \begin{bmatrix} 2 \\ -4 \\ 2 \end{bmatrix} \Rightarrow x_3 = \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$$

Result:-

Eigenvalues	Eigen vectors
2	$\begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$
3	$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$
6	$\begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$

Problem 2:-

Find the Eigen values and Eigen vectors of the matrix

$$A = \begin{bmatrix} 11 & -4 & -7 \\ 7 & -2 & -5 \\ 10 & -4 & -6 \end{bmatrix}$$

(A/M-2018)

Soln:-

$$\text{Given } A = \begin{bmatrix} 11 & -4 & -7 \\ 7 & -2 & -5 \\ 10 & -4 & -6 \end{bmatrix}$$

Step 1:-

Let the char. eqn be

$$\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$$

$$S_1 = 11 - 2 - 6 = 3$$

$$S_2 = \begin{vmatrix} -2 & -5 \\ -4 & -6 \end{vmatrix} + \begin{vmatrix} 11 & -7 \\ 10 & -6 \end{vmatrix} + \begin{vmatrix} 11 & -4 \\ 7 & -2 \end{vmatrix}$$

$$= (12 - 20) + (-66 + 70) + (-22 + 28)$$

$$= -8 + 4 + 6$$

$$= 2$$

$$S_3 = \begin{vmatrix} 11 & -4 & -7 \\ 7 & -2 & -5 \\ 10 & -4 & -6 \end{vmatrix}$$

$$= 11(12 - 20) + 4(-42 + 50)$$

$$- 7(-28 + 20)$$

$$= -88 + 32 + 56$$

$$= 0$$

$$\therefore \text{char eqn is } \lambda^3 - 3\lambda^2 + 2\lambda = 0$$

$$\text{Step 2:- } \lambda(\lambda^2 - 3\lambda + 2) = 0$$

$$\lambda = 0, \quad \lambda^2 - 3\lambda + 2 = 0$$

$$(\lambda - 2)(\lambda - 1) = 0$$

$$\lambda = 2, 1$$

$\therefore$ The eigen values are 0, 1, 2.

Step 3:-

$$\text{Solve } (A - \lambda I)X = 0$$

$$\begin{pmatrix} 11-\lambda & -4 & -7 \\ 7 & -2-\lambda & -5 \\ 10 & -4 & -6-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\left. \begin{aligned} (11-\lambda)x_1 - 4x_2 - 7x_3 &= 0 \\ 7x_1 - (2+\lambda)x_2 - 5x_3 &= 0 \\ 10x_1 - 4x_2 - (6+\lambda)x_3 &= 0 \end{aligned} \right\} \text{---(I)}$$

Case(i) put $\lambda=0$ in (I)

$$11x_1 - 4x_2 - 7x_3 = 0 \quad \text{---(1)}$$

$$7x_1 - 2x_2 - 5x_3 = 0 \quad \text{---(2)}$$

$$10x_1 - 4x_2 - 6x_3 = 0 \quad \text{---(3)}$$

Solve (1) & (2)

$$\frac{x_1}{\begin{vmatrix} -4 & -7 \\ -2 & -5 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} -7 & 11 \\ -5 & 7 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} 11 & -4 \\ 7 & -2 \end{vmatrix}}$$

$$\frac{x_1}{20-14} = \frac{x_2}{-49+55} = \frac{x_3}{-22+28}$$

$$\frac{x_1}{6} = \frac{x_2}{6} = \frac{x_3}{6}$$

$$\therefore x_1 = \begin{bmatrix} 6 \\ 6 \\ 6 \end{bmatrix} \Rightarrow x_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Case(ii) Put $\lambda=1$ in (I)

$$10x_1 - 4x_2 - 7x_3 = 0 \quad \text{---(4)}$$

$$7x_1 - 3x_2 - 5x_3 = 0 \quad \text{---(5)}$$

$$10x_1 - 4x_2 - 7x_3 = 0 \quad \text{---(6)}$$

Solve (4) & (5)

$$\frac{x_1}{\begin{vmatrix} -4 & -7 \\ -3 & -5 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} -7 & 10 \\ -5 & 7 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} 10 & -4 \\ 7 & -3 \end{vmatrix}}$$

$$\frac{x_1}{20-21} = \frac{x_2}{-49+50} = \frac{x_3}{-30+28}$$

$$\frac{x_1}{-1} = \frac{x_2}{1} = \frac{x_3}{-2}$$

$$\therefore x_2 = \begin{bmatrix} -1 \\ 1 \\ -2 \end{bmatrix} \Rightarrow x_2 = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}$$

Case(iii) put $\lambda=2$ in (I)

$$9x_1 - 4x_2 - 7x_3 = 0 \quad \text{---(7)}$$

$$7x_1 - 4x_2 - 5x_3 = 0 \quad \text{---(8)}$$

$$10x_1 - 4x_2 - 8x_3 = 0 \quad \text{---(9)}$$

Solve (7) & (8)

$$\frac{x_1}{\begin{vmatrix} -4 & -7 \\ -4 & -5 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} -7 & 9 \\ -5 & 7 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} 9 & -4 \\ 7 & -4 \end{vmatrix}}$$

$$\frac{x_1}{20-28} = \frac{x_2}{-49+45} = \frac{x_3}{-36+28}$$

$$\frac{x_1}{-8} = \frac{x_2}{-4} = \frac{x_3}{-8}$$

$$\therefore x_3 = \begin{bmatrix} -8 \\ -4 \\ -8 \end{bmatrix} \Rightarrow x_3 = \begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix}$$

Result:-

Eigenvalues	0	1	2
Eigen vector	$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$	$\begin{bmatrix} -1 \\ 1 \\ -2 \end{bmatrix}$	$\begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix}$

Problem:3:-

Find the Eigen values and Eigen vectors of a matrix

$$A = \begin{bmatrix} -1 & 2 & -2 \\ 1 & 2 & 1 \\ -1 & -1 & 0 \end{bmatrix} \quad (\text{NID-2018})$$

Soln:-

$$\text{Given } A = \begin{bmatrix} -1 & 2 & -2 \\ 1 & 2 & 1 \\ -1 & -1 & 0 \end{bmatrix}$$

Step 1:-

Let the char. eqn be

$$\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$$

$$S_1 = -1 + 2 + 0 = 1$$

$$S_2 = \begin{vmatrix} 2 & 1 \\ -1 & 0 \end{vmatrix} + \begin{vmatrix} -1 & -2 \\ -1 & 0 \end{vmatrix} + \begin{vmatrix} -1 & 2 \\ 1 & 2 \end{vmatrix}$$

$$= (0+1) + (0-2) + (-2-2)$$

$$= -5$$

$$S_3 = \begin{vmatrix} -1 & 2 & -2 \\ 1 & 2 & 1 \\ -1 & -1 & 0 \end{vmatrix}$$

$$= -1(0+1) - 2(0+1) - 2(-1+2)$$

$$= -5$$

Step 2:-

$$\text{put } \lambda = 1 \Rightarrow 1^3 - 1^2 - 5 + 5 = 0$$

 $\therefore \lambda = 1$ is a root

$$\begin{array}{c|cccc} 1 & 1 & -1 & -5 & 5 \\ & 0 & 1 & 0 & -5 \\ \hline & 1 & 0 & -5 & 0 \end{array}$$

$$\lambda^2 - 5 = 0$$

$$\lambda^2 = 5$$

$$\lambda = \pm \sqrt{5}$$

 $\therefore$ The eigen values are, $1, -\sqrt{5}, \sqrt{5}$ Step 3:- Solve $(A - \lambda I)X = 0$

$$\begin{bmatrix} -1-\lambda & 2 & -2 \\ 1 & 2-\lambda & 1 \\ -1 & -1 & 0-\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\left. \begin{aligned} (-1-\lambda)x_1 + 2x_2 - 2x_3 &= 0 \\ x_1 + (2-\lambda)x_2 + x_3 &= 0 \\ -x_1 - x_2 - \lambda x_3 &= 0 \end{aligned} \right\} \text{ (I)}$$

Case(i) put $\lambda = 1$ in (I)

$$-2x_1 + 2x_2 - 2x_3 = 0 \quad \text{--- (1)}$$

$$x_1 + x_2 + x_3 = 0 \quad \text{--- (2)}$$

$$-x_1 - x_2 - x_3 = 0 \quad \text{--- (3)}$$

Solve (1) & (2)

$$\frac{x_1}{\begin{vmatrix} 2 & -2 \\ 1 & 1 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} -2 & -2 \\ 1 & 1 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} -2 & 2 \\ 1 & 1 \end{vmatrix}}$$

$$\frac{x_1}{2+2} = \frac{x_2}{-2+2} = \frac{x_3}{-2-2}$$

$$\frac{x_1}{4} = \frac{x_2}{0} = \frac{x_3}{-4}$$

$$\therefore X_1 = \begin{bmatrix} 4 \\ 0 \\ -4 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

Case(ii) put $\lambda = \sqrt{5}$ in (I)

$$(-1-\sqrt{5})x_1 + 2x_2 - 2x_3 = 0 \quad \text{--- (4)}$$

$$x_1 + (2-\sqrt{5})x_2 + x_3 = 0 \quad \text{--- (5)}$$

$$-x_1 - x_2 - \sqrt{5}x_3 = 0 \quad \text{--- (6)}$$

Solve (4) & (5)

$$\frac{x_1}{\begin{vmatrix} 2 & -2 \\ 2-\sqrt{5} & 1 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} -2 & -1-\sqrt{5} \\ 1 & 1 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} -1-\sqrt{5} & 2 \\ 1 & 2-\sqrt{5} \end{vmatrix}}$$

$$\frac{x_1}{2+4-2\sqrt{5}} = \frac{x_2}{-2+1\sqrt{5}} = \frac{x_3}{-2-2\sqrt{5} + \sqrt{5}+5-2}$$

$$\frac{x_1}{6-2\sqrt{5}} = \frac{x_2}{-1+\sqrt{5}} = \frac{x_3}{1-\sqrt{5}}$$

$$\frac{x_1}{(-1+\sqrt{5})(-1+\sqrt{5})} = \frac{x_2}{-1+\sqrt{5}} = \frac{x_3}{-(-1+\sqrt{5})}$$

$$\frac{x_1}{-1+\sqrt{5}} = \frac{x_2}{1} = \frac{x_3}{-1}$$

$$\therefore X_2 = \begin{bmatrix} -1+\sqrt{5} \\ 1 \\ -1 \end{bmatrix}$$

case(iii) Put $\lambda = -\sqrt{5}$ in (I)

$$(-1+\sqrt{5})x_1 + 2x_2 - 2x_3 = 0 \rightarrow (7)$$

$$x_1 + (2+\sqrt{5})x_2 + x_3 = 0 \rightarrow (8)$$

$$-x_1 - x_2 + \sqrt{5}x_3 = 0 \rightarrow (9)$$

Solve (8) & (9)

$$\frac{x_1}{2+\sqrt{5}} = \frac{x_2}{1} = \frac{x_3}{-1}$$

$$\frac{x_1}{2\sqrt{5}+5+1} = \frac{x_2}{-1-\sqrt{5}} = \frac{x_3}{-1+2+\sqrt{5}}$$

$$\frac{x_1}{2\sqrt{5}+6} = \frac{x_2}{-(1+\sqrt{5})} = \frac{x_3}{1+\sqrt{5}}$$

$$\frac{x_1}{(1+\sqrt{5})(1+\sqrt{5})} = \frac{x_2}{-(1+\sqrt{5})} = \frac{x_3}{1+\sqrt{5}}$$

$$\therefore X_3 = \begin{bmatrix} 1+\sqrt{5} \\ -1 \\ 1 \end{bmatrix}$$

Non Symmetric Matrices with Repeated Eigen Values

Problem-1:-

Find all the eigen values & Eigen Vector of the matrix

$$\begin{pmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{pmatrix}$$

Soln:-

$$\text{Let } A = \begin{pmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{pmatrix}$$

Step-1:-

Let the Char. eqn be

$$\lambda^3 - s_1\lambda^2 + s_2\lambda - s_3 = 0$$

$$s_1 = -2 + 1 + 0 = -1$$

$$s_2 = \begin{vmatrix} 1 & -6 \\ -2 & 0 \end{vmatrix} + \begin{vmatrix} -2 & -3 \\ -1 & 0 \end{vmatrix} + \begin{vmatrix} -2 & 2 \\ 2 & 1 \end{vmatrix}$$

$$= (0-12) + (0-3) + (2-4)$$

$$= -21$$

$$s_3 = \begin{vmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{vmatrix}$$

$$\therefore \text{Char. eqn is } \lambda^3 + \lambda^2 - 21\lambda - 45 = 0$$

$$\text{Step-2} \quad \text{Put } \lambda = -3, (-3)^3 + (-3)^2 - 21(-3) - 45 = 0$$

$$\begin{array}{r|rrrr} -3 & 1 & 1 & -21 & -45 \\ & 0 & -3 & 6 & 45 \\ \hline & 1 & -2 & -15 & 0 \end{array}$$

$$\lambda^2 - 2\lambda - 15 = 0$$

$$(\lambda + 3)(\lambda - 5) = 0$$

$$\lambda = -3, 5$$

$\therefore -3, -3, 5$ are eigen values.

Step 3:-

To find Eigen vectors,

Solve $(A - \lambda I)X = 0$

$$\begin{pmatrix} -2-\lambda & 2 & -3 \\ 2 & 1-\lambda & -6 \\ -1 & -2 & -\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\left. \begin{aligned} (-2-\lambda)x_1 + 2x_2 - 3x_3 &= 0 \\ 2x_1 + (1-\lambda)x_2 - 6x_3 &= 0 \\ -x_1 - 2x_2 - \lambda x_3 &= 0 \end{aligned} \right\} \text{--- (I)}$$

Case(i) Put $\lambda = 5$ in (I)

$$\begin{aligned} -7x_1 + 2x_2 - 3x_3 &= 0 & \text{--- (1)} \\ 2x_1 - 4x_2 - 6x_3 &= 0 & \text{--- (2)} \\ -x_1 - 2x_2 - 5x_3 &= 0 & \text{--- (3)} \end{aligned}$$

Solve (1) & (2)

$$\frac{x_1}{\begin{vmatrix} 2 & -3 \\ -4 & -6 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} -3 & -7 \\ -6 & 2 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} -7 & 2 \\ 2 & -4 \end{vmatrix}}$$

$$\frac{x_1}{-12-12} = \frac{x_2}{-6-42} = \frac{x_3}{28-4}$$

$$\frac{x_1}{-24} = \frac{x_2}{-48} = \frac{x_3}{24}$$

$$\therefore X_1 = \begin{bmatrix} -24 \\ -48 \\ 24 \end{bmatrix} \Rightarrow X_1 = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

Case(ii) Put $\lambda = -3$ in (I)

$$x_1 + 2x_2 - 3x_3 = 0 \rightarrow \text{--- (4)}$$

$$2x_1 + 4x_2 - 6x_3 = 0 \rightarrow \text{--- (5)}$$

$$-x_1 - 2x_2 + 3x_3 = 0 \rightarrow \text{--- (6)}$$

All the equations are same,

Take (4)

$$\text{Put } x_1 = 0$$

$$2x_2 - 3x_3 = 0$$

$$2x_2 = 3x_3$$

$$\frac{x_2}{3} = \frac{x_3}{2}$$

$$\therefore X_2 = \begin{bmatrix} 0 \\ 3 \\ 2 \end{bmatrix}$$

Case(iii)

$$\text{Put } x_2 = 0$$

$$x_1 - 3x_3 = 0$$

$$x_1 = 3x_3$$

$$\frac{x_1}{3} = \frac{x_3}{1}$$

$$\therefore X_3 = \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}$$

Problem-2:-

Find the eigenvalues &

Eigen vectors of $\begin{bmatrix} 3 & 10 & 5 \\ -2 & -3 & -4 \\ 3 & 5 & 7 \end{bmatrix}$

Soln:-

Step 1:-

Let the char. eqn be

$$\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$$

$$S_1 = 3 - 3 + 7 = 7$$

$$S_2 = \begin{vmatrix} -3 & -4 \\ 5 & 7 \end{vmatrix} + \begin{vmatrix} 3 & 5 \\ 3 & 7 \end{vmatrix} + \begin{vmatrix} 3 & 10 \\ -2 & -3 \end{vmatrix}$$

$$= (-21+20) + (21+15) + (-9+20)$$

$$= -1 + 6 + 11$$

$$= 16$$

$$S_3 = \begin{vmatrix} 3 & 10 & 5 \\ -2 & -3 & -4 \\ 3 & 5 & 7 \end{vmatrix}$$

$$= 3(-21 + 20) - 10(-14 + 12) + 5(10 + 9)$$

$$= 12$$

$$\therefore \text{The char. eqn is } \lambda^3 - 7\lambda^2 + 16\lambda - 12 = 0$$

Step 2:-

$$\lambda = 2 \Rightarrow 8 - 28 + 32 - 12 = 0$$

$\therefore \lambda = 2$ is a one root

$$\begin{array}{r|rrrr} 2 & 1 & -7 & 16 & -12 \\ & 0 & 2 & -10 & 12 \\ \hline & 1 & -5 & 6 & 0 \end{array}$$

$$\lambda^2 - 5\lambda + 6 = 0$$

$$(\lambda - 3)(\lambda - 2) = 0$$

$$\lambda = 3, 2$$

$\therefore 2, 2, 3$ are the eigen values

Step-3:-

$$\text{Solve } (A - \lambda I)x = 0$$

$$\begin{bmatrix} 3-\lambda & 10 & 5 \\ -2 & -3-\lambda & -4 \\ 3 & 5 & 7-\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\left. \begin{aligned} (3-\lambda)x_1 + 10x_2 + 5x_3 &= 0 \\ -2x_1 - (3+\lambda)x_2 - 4x_3 &= 0 \\ 3x_1 + 5x_2 + (7-\lambda)x_3 &= 0 \end{aligned} \right\} \rightarrow \text{I}$$

Case(i) put $\lambda = 3$ in (I)

$$0x_1 + 10x_2 + 5x_3 = 0 \rightarrow \text{①}$$

$$-2x_1 - 6x_2 - 4x_3 = 0 \rightarrow \text{②}$$

$$3x_1 + 5x_2 + 4x_3 = 0 \rightarrow \text{③}$$

Solve ① & ②

$$\frac{x_1}{10 \ 5} = \frac{x_2}{5 \ 0} = \frac{x_3}{0 \ 10} \Rightarrow \begin{bmatrix} 10 & 5 \\ -6 & -4 \end{bmatrix} \Rightarrow \begin{bmatrix} 5 & 0 \\ -4 & -2 \end{bmatrix} \Rightarrow \begin{bmatrix} 0 & 10 \\ -2 & -6 \end{bmatrix}$$

$$\frac{x_1}{-40+30} = \frac{x_2}{-10-0} = \frac{x_3}{0+20}$$

$$\frac{x_1}{-10} = \frac{x_2}{-10} = \frac{x_3}{20}$$

$$\therefore x_1 = \begin{bmatrix} -10 \\ -10 \\ 20 \end{bmatrix} \Rightarrow x_1 = \begin{bmatrix} +1 \\ -1 \\ -2 \end{bmatrix}$$

case(ii) Put $\lambda = 2$ in (I)

$$x_1 + 10x_2 + 5x_3 = 0 \rightarrow \text{④}$$

$$-2x_1 - 5x_2 - 4x_3 = 0 \rightarrow \text{⑤}$$

$$3x_1 + 5x_2 + 5x_3 = 0 \rightarrow \text{⑥}$$

Solve ④ & ⑤

$$\frac{x_1}{10 \ 5} = \frac{x_2}{5 \ 1} = \frac{x_3}{1 \ 10} \Rightarrow \begin{bmatrix} 10 & 5 \\ -5 & -4 \end{bmatrix} \Rightarrow \begin{bmatrix} 5 & 1 \\ -4 & -2 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 10 \\ -2 & -5 \end{bmatrix}$$

$$\frac{x_1}{-40+25} = \frac{x_2}{-10+4} = \frac{x_3}{-5+20}$$

$$\frac{x_1}{-15} = \frac{x_2}{-6} = \frac{x_3}{15}$$

$$\frac{x_1}{5} = \frac{x_2}{2} = \frac{x_3}{-5}$$

$$\therefore x_2 = \begin{bmatrix} 5 \\ 2 \\ -5 \end{bmatrix} = x_3$$

Problem 3:-

Find the Eigen values and Eigen vectors of $\begin{bmatrix} 6 & -6 & 5 \\ 14 & -13 & 10 \\ 7 & -6 & 4 \end{bmatrix}$.

Soln:-

$$\text{Let } A = \begin{bmatrix} 6 & -6 & 5 \\ 14 & -13 & 10 \\ 7 & -6 & 4 \end{bmatrix}$$

Step 1:-

Let the char. eqn be

$$\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$$

$$S_1 = 6 - 13 + 4 = -3$$

$$S_2 = \begin{vmatrix} -13 & 10 \\ -6 & 4 \end{vmatrix} + \begin{vmatrix} 6 & 5 \\ 7 & 4 \end{vmatrix} + \begin{vmatrix} 6 & -6 \\ 14 & -13 \end{vmatrix}$$

$$= (-52 + 60) + (24 - 35) + (-78 + 84)$$

$$= 8 - 11 + 6 = 3$$

$$S_3 = \begin{vmatrix} 6 & -6 & 5 \\ 14 & -13 & 10 \\ 7 & -6 & 4 \end{vmatrix}$$

$$= 6(-52 + 60) + 6(56 - 70) + 5(-84 + 91)$$

$$= 6(8) + 6(-14) + 5(7)$$

$$= -1$$

$\therefore$ the char. eqn is $\lambda^3 + 3\lambda^2 + 3\lambda + 1 = 0$.

Step 2:-

$$\text{Put } \lambda = -1 \Rightarrow -1 + 3 - 3 + 1 = 0$$

$\therefore \lambda = -1$ is a root.

$$\begin{array}{r|rrrr} -1 & 1 & 3 & 3 & 1 \\ & 0 & -1 & -2 & -1 \\ \hline & 1 & 2 & 1 & 0 \end{array}$$

$$\therefore \lambda^2 + 2\lambda + 1 = 0$$

$$(\lambda + 1)(\lambda + 1) = 0$$

$$\lambda = -1, -1$$

$\therefore \lambda = -1, -1, -1$ are the eigen values.

Step 3:-

$$\text{Solve } (A - \lambda I)X = 0$$

$$\begin{bmatrix} 6 - \lambda & -6 & 5 \\ 14 & -13 - \lambda & 10 \\ 7 & -6 & 4 - \lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{cases} (6 - \lambda)x_1 - 6x_2 + 5x_3 = 0 \\ 14x_1 - (13 + \lambda)x_2 + 10x_3 = 0 \\ 7x_1 - 6x_2 + (4 - \lambda)x_3 = 0 \end{cases} \rightarrow \textcircled{1}$$

Case (i) Put $\lambda = -1$ in $\textcircled{1}$

$$7x_1 - 6x_2 + 5x_3 = 0 \rightarrow \textcircled{1}$$

$$14x_1 - 12x_2 + 10x_3 = 0 \rightarrow \textcircled{2}$$

$$7x_1 - 6x_2 + 5x_3 = 0 \rightarrow \textcircled{3}$$

All the three eqn. are same

Put $x_1 = 0$ in $\textcircled{1}$

$$-6x_2 + 5x_3 = 0$$

$$-6x_2 = -5x_3$$

$$\frac{x_2}{-5} = \frac{x_3}{-6}$$

$$\therefore x_1 = \begin{bmatrix} 0 \\ -5 \\ -6 \end{bmatrix} \Rightarrow x_1 = \begin{bmatrix} 0 \\ 5 \\ 6 \end{bmatrix}$$

Put $x_2 = 0$ in $\textcircled{1}$

$$7x_1 + 5x_3 = 0$$

$$7x_1 = -5x_3$$

$$\frac{x_1}{-5} = \frac{x_3}{7} \therefore x_2 = \begin{bmatrix} -5 \\ 0 \\ 7 \end{bmatrix}$$

Put $x_3 = 0$ in $\textcircled{1}$

$$7x_1 - 6x_2 = 0$$

$$7x_1 = 6x_2$$

$$\frac{x_1}{6} = \frac{x_2}{7} \therefore x_3 = \begin{bmatrix} 6 \\ 7 \\ 0 \end{bmatrix}$$

Problem: 4

Find the eigen values and Eigen vectors of $\begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{bmatrix}$

Soln:-

$$\text{Let } A = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{bmatrix}$$

Given matrix is an upper triangular matrix. Hence 2, 2, 2 are the eigen values.

$$\text{Solve } (A - \lambda I)X = 0$$

$$\begin{bmatrix} 2-\lambda & 1 & 0 \\ 0 & 2-\lambda & 1 \\ 0 & 0 & 2-\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{cases} (2-\lambda)x_1 + x_2 + 0x_3 = 0 \\ 0x_1 + (2-\lambda)x_2 + x_3 = 0 \\ 0x_1 + 0x_2 + (2-\lambda)x_3 = 0 \end{cases} \quad \text{--- (1)}$$

When $\lambda = 2$

$$0x_1 + x_2 + 0x_3 = 0 \quad \rightarrow \text{--- (1)}$$

$$0x_1 + 0x_2 + x_3 = 0 \quad \rightarrow \text{--- (2)}$$

$$0x_1 + 0x_2 + 0x_3 = 0 \quad \rightarrow \text{--- (3)}$$

Solve (1) & (2)

$$\frac{x_1}{\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 0 & 0 \\ 1 & 0 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} 0 & 1 \\ 0 & 0 \end{vmatrix}}$$

$$\frac{x_1}{1} = \frac{x_2}{0} = \frac{x_3}{0}$$

$$\therefore x_1 = x_2 = x_3 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

Symmetric Matrices with Non-Repeated Eigen values:Problem: 1

Find the eigen values and Eigen vectors of $A = \begin{bmatrix} 2 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 2 \end{bmatrix}$

Soln:-

$$\text{Let } A = \begin{bmatrix} 2 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 2 \end{bmatrix}$$

Step-1

Let the char. eqn be $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$$S_1 = 2 + 2 + 2 = 6$$

$$\begin{aligned} S_2 &= \begin{vmatrix} 2 & 0 \\ 0 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 0 \\ 0 & 2 \end{vmatrix} \\ &= (4-0) + (4-1) + (4-0) \\ &= 11 \end{aligned}$$

$$\begin{aligned} S_3 &= \begin{vmatrix} 2 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 2 \end{vmatrix} \\ &= 2(4-0) - 0 + 1(0-2) \\ &= 8-2 = 6 \end{aligned}$$

$\therefore$ The char. eqn is $\lambda^3 - 6\lambda^2 + 11\lambda - 6 = 0$

Step-2:-

$$\lambda = 1 \Rightarrow 1 - 6 + 11 - 6 = 0$$

$$\begin{array}{r|rrrr} 1 & 1 & -6 & 11 & -6 \\ & 0 & 1 & -5 & 6 \\ \hline & 1 & -5 & 6 & 0 \end{array}$$

$$\lambda^2 - 5\lambda + 6 = 0$$

$$(\lambda - 2)(\lambda - 3) = 0$$

$$\lambda = 2, 3$$

$\therefore 1, 2, 3$ are eigen values.

Step-3:- solve $(A - \lambda I) = 0$

$$\begin{bmatrix} 2-\lambda & 0 & 1 \\ 0 & 2-\lambda & 0 \\ 1 & 0 & 2-\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\left. \begin{aligned} (2-\lambda)x_1 + 0x_2 + x_3 &= 0 \\ 0x_1 + (2-\lambda)x_2 + 0x_3 &= 0 \\ x_1 + 0x_2 + (2-\lambda)x_3 &= 0 \end{aligned} \right\} \rightarrow \textcircled{1}$$

Case(i) Put $\lambda = 1$ in $\textcircled{1}$

$$x_1 + 0x_2 + x_3 = 0 \rightarrow \textcircled{1}$$

$$0x_1 + x_2 + 0x_3 = 0 \rightarrow \textcircled{2}$$

$$x_1 + 0x_2 + x_3 = 0 \rightarrow \textcircled{3}$$

Solve $\textcircled{1}$ & $\textcircled{2}$

$$\frac{x_1}{\begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 1 & 1 \\ 0 & 0 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix}}$$

$$\frac{x_1}{-1} = \frac{x_2}{0} = \frac{x_3}{1}$$

$$\therefore X_1 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

Case(ii) Put $\lambda = 2$ in $\textcircled{1}$

$$0x_1 + 0x_2 + x_3 = 0 \rightarrow \textcircled{4}$$

$$0x_1 + 0x_2 + 0x_3 = 0 \rightarrow \textcircled{5}$$

$$x_1 + 0x_2 + 0x_3 = 0 \rightarrow \textcircled{6}$$

Solve $\textcircled{4}$ & $\textcircled{6}$

$$\frac{x_1}{\begin{vmatrix} 0 & 1 \\ 0 & 0 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} 1 & 0 \\ 1 & 0 \end{vmatrix}}$$

$$\frac{x_1}{0} = \frac{x_2}{1} = \frac{x_3}{0} \therefore X_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

Case(iii) put $\lambda = 3$ in $\textcircled{1}$

$$-x_1 + 0x_2 + x_3 = 0 \rightarrow \textcircled{7}$$

$$0x_1 - x_2 + 0x_3 = 0 \rightarrow \textcircled{8}$$

$$x_1 + 0x_2 - x_3 = 0 \rightarrow \textcircled{9}$$

Solve $\textcircled{7}$ & $\textcircled{8}$

$$\frac{x_1}{\begin{vmatrix} 0 & 1 \\ -1 & 0 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 1 & -1 \\ 0 & 0 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} -1 & 0 \\ 0 & -1 \end{vmatrix}}$$

$$\frac{x_1}{1} = \frac{x_2}{0} = \frac{x_3}{1}$$

$$\therefore X_3 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

Problem: 2

Find the Eigen values and Eigen vectors of the matrix

$$\begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$$

Soln:-

$$\text{Let } A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$$

Step 1:-

Let the char eqn be

$$\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$$

$$S_1 = 8 + 7 + 3 = 18$$

$$S_2 = \begin{vmatrix} 7 & -4 \\ -4 & 3 \end{vmatrix} + \begin{vmatrix} 8 & 2 \\ 2 & 3 \end{vmatrix} + \begin{vmatrix} 8 & -6 \\ -6 & 7 \end{vmatrix}$$

$$= (21 - 16) + (24 - 4) + (56 - 36)$$

$$= 45$$

$$S_3 = \begin{vmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{vmatrix}$$

$$= 8(21-16) + 6(-18+8) + 2(24-14)$$

$$= 0$$

∴ The char eqn's $\lambda^3 - 18\lambda^2 + 45\lambda = 0$

Step-2 :-

$$\lambda(\lambda^2 - 18\lambda + 45) = 0$$

$$\lambda = 0, (\lambda - 3)(\lambda - 15) = 0$$

$$\lambda = 0, 3, 15$$

∴ The eigen values are 0, 3, 15.

Step 3 :-

$$\text{Solve } (A - \lambda I) X = 0$$

$$\begin{bmatrix} 8-\lambda & -6 & 2 \\ -6 & 7-\lambda & -4 \\ 2 & -4 & 3-\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\left. \begin{aligned} (8-\lambda)x_1 - 6x_2 + 2x_3 &= 0 \\ -6x_1 + (7-\lambda)x_2 - 4x_3 &= 0 \\ 2x_1 - 4x_2 + (3-\lambda)x_3 &= 0 \end{aligned} \right\} \rightarrow \textcircled{1}$$

Case(i) put $\lambda = 0$ in $\textcircled{1}$

$$8x_1 - 6x_2 + 2x_3 = 0 \rightarrow \textcircled{1}$$

$$-6x_1 + 7x_2 - 4x_3 = 0 \rightarrow \textcircled{2}$$

$$2x_1 - 4x_2 + 3x_3 = 0 \rightarrow \textcircled{3}$$

Solve $\textcircled{1}$ & $\textcircled{2}$

$$\frac{x_1}{\begin{vmatrix} -6 & 2 \\ 7 & -4 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 2 & 8 \\ -4 & -6 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} 8 & -6 \\ -6 & 7 \end{vmatrix}}$$

$$\frac{x_1}{24-14} = \frac{x_2}{-12+32} = \frac{x_3}{56-36}$$

$$\frac{x_1}{10} = \frac{x_2}{20} = \frac{x_3}{20} \therefore X_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$$

Case (ii) put $\lambda = 3$ in $\textcircled{1}$

$$5x_1 - 6x_2 + 2x_3 = 0 \rightarrow \textcircled{4}$$

$$-6x_1 + 4x_2 - 4x_3 = 0 \rightarrow \textcircled{5}$$

$$2x_1 - 4x_2 + 0x_3 = 0 \rightarrow \textcircled{6}$$

Solve $\textcircled{4}$ & $\textcircled{5}$

$$\frac{x_1}{\begin{vmatrix} -6 & 2 \\ 4 & -4 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 2 & 5 \\ -4 & -6 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} 5 & -6 \\ -6 & 4 \end{vmatrix}}$$

$$\frac{x_1}{24-8} = \frac{x_2}{-12+20} = \frac{x_3}{20-36}$$

$$\frac{x_1}{16} = \frac{x_2}{8} = \frac{x_3}{-16}$$

$$\therefore X_2 = \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix}$$

Case(iii) put $\lambda = 15$ in $\textcircled{1}$

$$-7x_1 - 6x_2 + 2x_3 = 0 \rightarrow \textcircled{7}$$

$$-6x_1 - 8x_2 - 4x_3 = 0 \rightarrow \textcircled{8}$$

$$2x_1 - 4x_2 - 12x_3 = 0 \rightarrow \textcircled{9}$$

Solve $\textcircled{7}$ & $\textcircled{8}$

$$\frac{x_1}{\begin{vmatrix} -6 & 2 \\ -8 & -4 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 2 & -7 \\ -4 & -6 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} -7 & -6 \\ -6 & -8 \end{vmatrix}}$$

$$\frac{x_1}{24+16} = \frac{x_2}{-12-28} = \frac{x_3}{56-36}$$

$$\frac{x_1}{40} = \frac{x_2}{-40} = \frac{x_3}{20}$$

$$\therefore X_3 = \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix}$$

Symmetric Matrices with Repeated Eigen values :-

Problem 1:-

Find the Eigen values and Eigen vectors of the matrix

$$\begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

Soln:-

$$\text{Let } A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

Step 1:-

Let the char. eqn. be

$$\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$$

$$S_1 = 0$$

$$S_2 = \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} + \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} + \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix}$$

$$= -1 - 1 - 1 = -3$$

$$S_3 = \begin{vmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{vmatrix} = 0 - 1(0-1) + 1(1-0) = 2$$

$$\therefore \text{char. eqn is } \lambda^3 - 3\lambda - 2 = 0$$

$$\text{Step 2:- } \lambda = -1 \Rightarrow -1 - 3(-1) - 2 = 0$$

$$\begin{array}{r|rrrr} -1 & 1 & 0 & -3 & -2 \\ & 0 & -1 & 1 & 2 \\ \hline & 1 & -1 & -2 & 0 \end{array}$$

$$\lambda^2 - \lambda - 2 = 0$$

$$(\lambda+1)(\lambda-2) = 0$$

$$\lambda = -1, +2$$

$\therefore$ The eigen values are 2, -1, -1

Step 3:- Solve $(A - \lambda I)X = 0$

$$\begin{bmatrix} -\lambda & 1 & 1 \\ 1 & -\lambda & 1 \\ 1 & 1 & -\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\left. \begin{array}{l} -\lambda x_1 + x_2 + x_3 = 0 \\ x_1 - \lambda x_2 + x_3 = 0 \\ x_1 + x_2 - \lambda x_3 = 0 \end{array} \right\} \rightarrow \textcircled{I}$$

Case(i) $\lambda = 2$ in $\textcircled{I}$

$$-2x_1 + x_2 + x_3 = 0 \rightarrow \textcircled{1}$$

$$x_1 - 2x_2 + x_3 = 0 \rightarrow \textcircled{2}$$

$$x_1 + x_2 - 2x_3 = 0 \rightarrow \textcircled{3}$$

$$\frac{x_1}{\begin{vmatrix} 1 & 1 \\ -2 & 1 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 1 & -2 \\ 1 & 1 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} -2 & 1 \\ 1 & -2 \end{vmatrix}}$$

$$\frac{x_1}{1+2} = \frac{x_2}{1+2} = \frac{x_3}{4-1}$$

$$\frac{x_1}{3} = \frac{x_2}{3} = \frac{x_3}{3} \therefore x_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Case(ii) $\lambda = -1$ in $\textcircled{I}$

$$x_1 + x_2 + x_3 = 0 \rightarrow \textcircled{4}$$

$$x_1 + x_2 + x_3 = 0 \rightarrow \textcircled{5}$$

$$x_1 + x_2 + x_3 = 0 \rightarrow \textcircled{6}$$

All the eqns. are same.

$$\text{Put } x_1 = 0 \Rightarrow x_2 + x_3 = 0 \Rightarrow x_2 = -x_3$$

$$\frac{x_2}{-1} = \frac{x_3}{1} \therefore x_2 = \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$$

Case(iii)

Let $x_3 = \begin{bmatrix} l \\ m \\ n \end{bmatrix}$ as x_3 is orthogonal to x_1 & x_2 .

$$\begin{bmatrix} 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} l \\ m \\ n \end{bmatrix} = 0 \Rightarrow l + m + n = 0 \rightarrow \textcircled{7}$$

$$\begin{bmatrix} 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} l \\ m \\ n \end{bmatrix} = 0 \Rightarrow -l - m + n = 0 \rightarrow \textcircled{8}$$

Solve $\textcircled{7}$ & $\textcircled{8}$

$$\frac{l}{\begin{vmatrix} 1 & 1 \\ -1 & -1 \end{vmatrix}} = \frac{m}{\begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix}} = \frac{n}{\begin{vmatrix} 1 & 1 \\ 0 & -1 \end{vmatrix}}$$

$$\frac{l}{2} = \frac{m}{-1} = \frac{n}{-1} \therefore x_3 = \begin{bmatrix} 2 \\ -1 \\ -1 \end{bmatrix}$$

Properties of Eigen values and Eigen vectors.

Prop-1:-

* The Sum of the Eigen values of a matrix is the sum of the elements of the main diagonal.

(or)

The Sum of the Eigen values of a matrix is equal to the trace of the matrix.

* Product of the Eigen values is equal to the determinant of the matrix

Prop-2:-

A square matrix A and its transpose A^T have the same Eigen values

Proof:-

Let A be a square matrix of order n .

The char. eqn of A & A^T are

$$|A - \lambda I| = 0 \quad \text{L(I)} \quad \text{and} \quad |A^T - \lambda I| = 0 \quad \text{L(II)}$$

Since, the determinant value is unaltered by the inter change of rows & columns

$$\text{W.K.T } |A| = |A^T|$$

Hence (I) and (II) are identical

$\therefore$ The Eigen values of A and A^T are the same.

Prop-3:-

The Eigen values of a triangular matrix are just the diagonal elements of the matrix.

Prop-4:-

If λ is an Eigen value of a matrix A , then $\frac{1}{\lambda}$, ($\lambda \neq 0$) is the Eigen value of A^{-1} .

Proof:-

If x be the Eigen vector corresponding to λ , then

$$Ax = \lambda x \quad \text{--- (1)}$$

x by A^T

$$A^T A x = \lambda A^T x$$

$$I x = \lambda A^T x$$

$\div$ by λ

$$\frac{1}{\lambda} x = A^T x$$

$$A^T x = \frac{1}{\lambda} x.$$

This being of the same form as (1) shows that $\frac{1}{\lambda}$ is an Eigen value of the A^T .

Prop-5:-

If λ is an Eigen value of an orthogonal matrix, then $\frac{1}{\lambda}$ is also its Eigen value.

Prop-6:-

If $\lambda_1, \lambda_2, \dots, \lambda_n$ are the Eigen values of a matrix A , then A^m has the Eigen values $\lambda_1^m, \lambda_2^m, \dots, \lambda_n^m$.

Proof:-

Let λ_i be the Eigen value of A and x_i the corresponding Eigen vector. Then

$$Ax_i = \lambda_i x_i \quad \text{--- ①}$$

$$\begin{aligned} \text{We have } A^2 x_i &= A(Ax_i) \\ &= A(\lambda_i x_i) \quad \text{by ①} \\ &= \lambda_i (Ax_i) \\ &= \lambda_i (\lambda_i x_i) \\ &= \lambda_i^2 x_i \end{aligned}$$

$$\text{Similarly } A^3 x_i = \lambda_i^3 x_i$$

In general,

$$A^m x_i = \lambda_i^m x_i \quad \text{--- ②}$$

① & ② are in same form.

Prop-7:-

The Eigen values of a real Symmetric matrix are equal real numbers.

Prop-8:-

The Eigenvectors corresponding to distinct Eigen value of a real symmetric matrix are orthogonal.

Prop-9:-

The similar matrices have same Eigen values.

Reduction of Quadratic form to canonical form by orthogonal transformation:-

Quadratic form:-

A homogeneous polynomial of second degree in any number of variables is called a quadratic form.

Note:-

The matrix corresponding to the quadratic form is

$$\begin{bmatrix} \text{coeff } x_1^2 & \frac{1}{2} \text{coeff } x_1 x_2 & \frac{1}{2} \text{coeff } x_1 x_3 \\ \frac{1}{2} \text{coeff } x_2 x_1 & \text{coeff } x_2^2 & \frac{1}{2} \text{coeff } x_2 x_3 \\ \frac{1}{2} \text{coeff } x_3 x_1 & \frac{1}{2} \text{coeff } x_3 x_2 & \text{coeff } x_3^2 \end{bmatrix}$$

Index:- (S)

Number of positive square terms in the canonical form.

Signature:- (2S-r)

Difference of number of positive and negative square terms in the canonical form.

Rank:- (r)

Number of square terms in the canonical form.

Nature of the Quadratic form.

- * Positive definite $\Rightarrow$ If all the Eigen values are positive.
- * Positive Semi definite $\Rightarrow$ If all the Eigen values are positive and at least one zero.
- * Negative definite $\Rightarrow$ If all eigen values are negative.
- * Negative semi definite $\Rightarrow$ If all the Eigen values are negative and at least one zero.
- * Indefinite $\Rightarrow$ In all other cases.

Test for Nature of a quadratic form through Principal minors:

- i) The Q.F is positive definite if $D_1, D_2, \dots, D_n$ are all positive i.e., $D_n > 0$ for all n , where $D_1, D_2, \dots, D_n$ are the Principal minors of A.
- ii) The Q.F is negative definite if $D_1, D_3, D_5, \dots$ are all negative and $D_2, D_4, \dots, D_6$ are all positive. i.e., $(-1)^n D_n > 0$ for all n .

(iii) The Q.F is positive semi definite if $D_n \geq 0$ and at least one $D_i = 0$

iv) The Q.F is negative semi definite if $(-1)^n D_n \geq 0$ and at least one $D_i = 0$.

v) The Q.F is indefinite in all other cases.

Working Procedure:-

Step-1: write the matrix of the Quadratic form.

Step-2:- Find the char. eqn.

Step-3:- Find the Eigen Values

Step-4:- Find the Eigen vectors.

Step-5:- Form Normalized matrix N.

Step 6:- Find N^T

Step 7:- Find AN

Step 8: Find $D = N^T A N$

Step 9: Canonical form.

$$[y_1 y_2 y_3] D \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

Problem - 1:-

Reduce the Q.F $Q = 6x^2 + 3y^2 + 3z^2 - 4xy - 2yz + 4zx$ into canonical form by an orthogonal transformation.

Soln:-

Given
 $Q = 6x^2 + 3y^2 + 3z^2 - 4xy - 2yz + 4zx$

Step-1 Matrix form:-

$$A = \begin{bmatrix} \text{coeff } x_1^2 & \frac{1}{2} \text{coeff } x_1 x_2 & \frac{1}{2} \text{coeff } x_1 x_3 \\ \frac{1}{2} \text{coeff } x_1 x_2 & \text{coeff } x_2^2 & \frac{1}{2} \text{coeff } x_2 x_3 \\ \frac{1}{2} \text{coeff } x_1 x_3 & \frac{1}{2} \text{coeff } x_2 x_3 & \text{coeff } x_3^2 \end{bmatrix}$$

$$= \begin{bmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{bmatrix}$$

Step 2:-

To find char. eqn.

Let the char. eqn be

$$\lambda^3 - S_1 \lambda^2 + S_2 \lambda - S_3 = 0$$

$$S_1 = 6 + 3 + 3 = 12$$

$$S_2 = \begin{vmatrix} 3 & -1 \\ -1 & 3 \end{vmatrix} + \begin{vmatrix} 6 & 2 \\ 2 & 3 \end{vmatrix} + \begin{vmatrix} 6 & -2 \\ -2 & 3 \end{vmatrix}$$

$$= (9-1) + (18-4) + (18-4) \\ = 36$$

$$S_3 = \begin{vmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{vmatrix}$$

$$= 6(9-1) + 2(-6+2) + 2(2-6) \\ = 32$$

$\therefore$ The char. eqn

$$\lambda^3 - 12\lambda^2 + 36\lambda - 32 = 0$$

Step 3:-

To find Eigen values:-

$$\text{If } \lambda = 2 \Rightarrow 8 - 48 + 72 - 32 = 0$$

$\therefore \lambda = 2$ is a root.

$$2 \left| \begin{array}{ccc|c} 1 & -12 & 36 & -32 \\ 0 & 2 & -20 & 32 \\ 1 & -10 & 16 & 0 \end{array} \right|$$

$$\therefore \lambda^2 - 10\lambda + 16 = 0$$

$$(\lambda - 8)(\lambda - 2) = 0$$

$$\lambda = 2, 8$$

$\therefore$ The Eigen values are

$$8, 2, 2$$

Step 4:- To find Eigen vectors

$$\text{Solve } (A - \lambda I)X = 0$$

$$\begin{bmatrix} 6-\lambda & -2 & 2 \\ -2 & 3-\lambda & -1 \\ 2 & -1 & 3-\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\left. \begin{aligned} (6-\lambda)x_1 - 2x_2 + 2x_3 &= 0 \\ -2x_1 + (3-\lambda)x_2 - x_3 &= 0 \\ 2x_1 - x_2 + (3-\lambda)x_3 &= 0 \end{aligned} \right\} \text{--- (I)}$$

Case (i) put $\lambda = 8$ in (I)

$$-2x_1 - 2x_2 + 2x_3 = 0 \quad \text{--- (1)}$$

$$-2x_1 - 5x_2 - x_3 = 0 \quad \text{--- (2)}$$

$$2x_1 - x_2 - 5x_3 = 0 \quad \text{--- (3)}$$

Solve (1) & (2)

$$\frac{x_1}{\begin{vmatrix} -2 & 2 \\ -5 & -1 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 2 & -2 \\ -1 & -2 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} -2 & -2 \\ -2 & -5 \end{vmatrix}}$$

$$\frac{x_1}{2+10} = \frac{x_2}{-4-2} = \frac{x_3}{10-14}$$

$$\frac{x_1}{12} = \frac{x_2}{-6} = \frac{x_3}{6} \therefore X_1 = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$$

Case (i) put $\lambda = 2$ in (1)

$$4x_1 - 2x_2 + 2x_3 = 0 \rightarrow (4)$$

$$-2x_1 + x_2 - x_3 = 0 \rightarrow (5)$$

$$2x_1 - x_2 + x_3 = 0 \rightarrow (6)$$

All the eqn are same,

Take $2x_1 - x_2 + x_3 = 0$

put $x_1 = 0$, we get

$$-x_2 + x_3 = 0$$

$$x_2 = x_3 \therefore X_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

Case (ii)

To find the third Eigen vector orthogonal to x_1 and x_2 . Since the matrix A is symmetric.

Let $X_3 = \begin{bmatrix} l \\ m \\ n \end{bmatrix}$ as X_3 is orthogonal to x_1 and x_2 .

$$X_1^T X_3 = 0 \Rightarrow [2 \ -1 \ 1] \begin{bmatrix} l \\ m \\ n \end{bmatrix} = 0$$

$$\Rightarrow 2l - m + n = 0 \quad (7)$$

$$X_2^T X_3 = 0 \Rightarrow [0 \ 1 \ 1] \begin{bmatrix} l \\ m \\ n \end{bmatrix} = 0$$

$$\Rightarrow 0l + m + n = 0 \quad (8)$$

$$\frac{l}{1} = \frac{m}{1} = \frac{n}{2} = \frac{1}{2}$$

$$\frac{l}{-2} = \frac{m}{-2} = \frac{n}{2} \therefore X_3 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$$

Step 5:- Normalised matrix N :-

$$N = \begin{bmatrix} \frac{2}{\sqrt{6}} & 0 & \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{3}} \end{bmatrix}$$

Step 6:- Find N^T

$$N^T = \begin{bmatrix} \frac{2}{\sqrt{6}} & -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{3}} \end{bmatrix}$$

Step 7:- Find AN

$$AN = \begin{bmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{bmatrix} \begin{bmatrix} \frac{2}{\sqrt{6}} & 0 & \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{3}} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{12+2+2}{\sqrt{6}} & \frac{0-2+2}{\sqrt{2}} & \frac{6-2-2}{\sqrt{3}} \\ \frac{-4-3+1}{\sqrt{6}} & \frac{0+3-1}{\sqrt{2}} & \frac{-2+3+1}{\sqrt{3}} \\ \frac{4+1+3}{\sqrt{6}} & \frac{0-1+3}{\sqrt{2}} & \frac{2-1-3}{\sqrt{3}} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{16}{\sqrt{6}} & 0 & \frac{2}{\sqrt{3}} \\ -\frac{8}{\sqrt{6}} & \frac{2}{\sqrt{2}} & \frac{2}{\sqrt{3}} \\ \frac{8}{\sqrt{6}} & \frac{2}{\sqrt{2}} & -\frac{2}{\sqrt{3}} \end{bmatrix}$$

Step 8:- Find $D = N^T AN$

$$D = \begin{bmatrix} \frac{2}{\sqrt{6}} & -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{3}} \end{bmatrix} \begin{bmatrix} \frac{16}{\sqrt{6}} & 0 & \frac{2}{\sqrt{3}} \\ -\frac{8}{\sqrt{6}} & \frac{2}{\sqrt{2}} & \frac{2}{\sqrt{3}} \\ \frac{8}{\sqrt{6}} & \frac{2}{\sqrt{2}} & -\frac{2}{\sqrt{3}} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{32+8+8}{6} & \frac{0-2+2}{\sqrt{12}} & \frac{4-2-2}{\sqrt{18}} \\ \frac{0-8+8}{\sqrt{12}} & \frac{0+2+2}{2} & \frac{0+2-2}{\sqrt{6}} \\ \frac{16-8-8}{\sqrt{18}} & \frac{0+2-2}{\sqrt{6}} & \frac{2+2+2}{3} \end{bmatrix}$$

$$= \begin{bmatrix} 8 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

Step 9: canonical form:-

$$[y_1 \ y_2 \ y_3] \begin{pmatrix} 8 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

$$= 8y_1^2 + 2y_2^2 + 2y_3^2$$

∴ The canonical form is $8y_1^2 + 2y_2^2 + y_3^2 =$

Problem: 2:-

Reduce the quadratic form $2x^2 + 5y^2 + 3z^2 + 4xy$ to a canonical form through an orthogonal transformation. Find also its nature. (Apr-2018)

Soln:- Given $Q = 2x^2 + 5y^2 + 3z^2 + 4xy$

Step 1:- Matrix form

$$A = \begin{bmatrix} \text{co. eff } x_1^2 & \frac{1}{2} \text{ co. eff } x_1 x_2 & \frac{1}{2} \text{ co. eff } x_1 x_3 \\ \frac{1}{2} \text{ co. eff } x_2 x_1 & \text{co. eff } x_2^2 & \frac{1}{2} \text{ co. eff } x_2 x_3 \\ \frac{1}{2} \text{ co. eff } x_3 x_1 & \frac{1}{2} \text{ co. eff } x_3 x_2 & \text{co. eff } x_3^2 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 2 & 0 \\ 2 & 5 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

Step 2:- To find the char. eqn:-

Let the char. eqn. be

$$\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$$

$$S_1 = 2 + 5 + 3 = 10$$

$$S_2 = \begin{vmatrix} 5 & 0 \\ 0 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 0 \\ 0 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 2 \\ 2 & 5 \end{vmatrix}$$

$$= 15 + 6 + (10 - 4)$$

$$= 27$$

$$S_3 = \begin{vmatrix} 2 & 2 & 0 \\ 2 & 5 & 0 \\ 0 & 0 & 3 \end{vmatrix}$$

$$= 2(15 - 0) - 2(6 - 0) = 18$$

$$\therefore \text{char eqn is } \lambda^3 - 10\lambda^2 + 27\lambda - 18 = 0$$

Step 3:- To find the Eigen values

$$\text{Put } \lambda = 1 \Rightarrow 1 - 10 + 27 - 18 = 0$$

∴ $\lambda = 1$ is a root.

$$\begin{array}{c|ccc} 1 & 1 & -10 & 27 & -18 \\ & 0 & 1 & -9 & 18 \\ \hline & 1 & -9 & 18 & 0 \end{array}$$

$$\lambda^2 - 9\lambda + 18 = 0$$

$$(\lambda - 3)(\lambda - 6) = 0$$

$$\lambda = 3, 6$$

∴ The Eigen values are 1, 3, 6.

Step-4:- To find Eigenvectors:-

Solve $(A - \lambda I)x = 0$

$$\begin{bmatrix} 2-\lambda & 2 & 0 \\ 2 & 5-\lambda & 0 \\ 0 & 0 & 3-\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\left. \begin{aligned} (2-\lambda)x_1 + 2x_2 + 0x_3 &= 0 \\ 2x_1 + (5-\lambda)x_2 + 0x_3 &= 0 \\ 0x_1 + 0x_2 + (3-\lambda)x_3 &= 0 \end{aligned} \right\} \textcircled{I}$$

Case(i) Put $\lambda=1$ in $\textcircled{I}$

$$x_1 + 2x_2 + 0x_3 = 0 \rightarrow \textcircled{1}$$

$$2x_1 + 4x_2 + 0x_3 = 0 \rightarrow \textcircled{2}$$

$$0x_1 + 0x_2 + 2x_3 = 0 \rightarrow \textcircled{3}$$

Solve $\textcircled{1}$ & $\textcircled{2}$

$$\frac{x_1}{\begin{vmatrix} 2 & 0 \\ 0 & 2 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 0 & 1 \\ 2 & 0 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} 1 & 2 \\ 0 & 0 \end{vmatrix}}$$

$$\frac{x_1}{4} = \frac{x_2}{-2} = \frac{x_3}{0}$$

$$\therefore x_1 = \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix}$$

Case(ii) Put $\lambda=3$ in $\textcircled{I}$

$$-x_1 + 2x_2 + 0x_3 = 0 \rightarrow \textcircled{4}$$

$$2x_1 + 2x_2 + 0x_3 = 0 \rightarrow \textcircled{5}$$

$$0x_1 + 0x_2 + 0x_3 = 0 \rightarrow \textcircled{6}$$

Solve $\textcircled{4}$ & $\textcircled{5}$

$$\frac{x_1}{\begin{vmatrix} 2 & 0 \\ 2 & 0 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 0 & -1 \\ 0 & 2 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} -1 & 2 \\ 2 & 2 \end{vmatrix}}$$

$$\frac{x_1}{0} = \frac{x_2}{0} = \frac{x_3}{-2-4}$$

$$\therefore x_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Case(iii) Put $\lambda=6$ in $\textcircled{I}$

$$-4x_1 + 2x_2 + 0x_3 = 0 \rightarrow \textcircled{7}$$

$$2x_1 - x_2 + 0x_3 = 0 \rightarrow \textcircled{8}$$

$$0x_1 + 0x_2 - 3x_3 = 0 \rightarrow \textcircled{9}$$

Solve $\textcircled{8}$ & $\textcircled{9}$

$$\frac{x_1}{\begin{vmatrix} -1 & 0 \\ 0 & -3 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 0 & 2 \\ -3 & 0 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} 2 & -1 \\ 0 & 0 \end{vmatrix}}$$

$$\frac{x_1}{3} = \frac{x_2}{6} = \frac{x_3}{0}$$

$$x_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$$

Step 5:-

Normalised matrix N:-

$$N = \begin{bmatrix} \frac{2}{\sqrt{5}} & 0 & \frac{1}{\sqrt{5}} \\ -\frac{1}{\sqrt{5}} & 0 & \frac{2}{\sqrt{5}} \\ 0 & 1 & 0 \end{bmatrix}$$

Step 6:-

$$NT = \begin{bmatrix} \frac{2}{5} & -\frac{1}{\sqrt{5}} & 0 \\ 0 & 0 & 1 \\ \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} & 0 \end{bmatrix}$$

Step 7:- Find AN

$$AN = \begin{bmatrix} 2 & 2 & 0 \\ 2 & 5 & 0 \\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} \frac{2}{\sqrt{5}} & 0 & \frac{1}{\sqrt{5}} \\ -\frac{1}{\sqrt{5}} & 0 & \frac{2}{\sqrt{5}} \\ 0 & 1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{4-2}{\sqrt{5}} & 0 & \frac{2+4}{\sqrt{5}} \\ \frac{4-5}{\sqrt{5}} & 0 & \frac{2+10}{\sqrt{5}} \\ 0 & 3 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{2}{\sqrt{5}} & 0 & \frac{6}{\sqrt{5}} \\ -\frac{1}{\sqrt{5}} & 0 & \frac{12}{\sqrt{5}} \\ 0 & 3 & 0 \end{bmatrix}$$

Step 8:- Find D = N^TAN

$$D = \begin{bmatrix} \frac{2}{\sqrt{5}} & -\frac{1}{\sqrt{5}} & 0 \\ 0 & 0 & 1 \\ \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} & 0 \end{bmatrix} \begin{bmatrix} \frac{2}{\sqrt{5}} & 0 & \frac{6}{\sqrt{5}} \\ -\frac{1}{\sqrt{5}} & 0 & \frac{12}{\sqrt{5}} \\ 0 & 3 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{4+1}{5} & 0 & \frac{12-12}{5} \\ 0 & 3 & 0 \\ \frac{2-2}{5} & 0 & \frac{6+24}{5} \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 6 \end{bmatrix}$$

Step 9:- canonical form:-

$$(y_1 y_2 y_3) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 6 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \\ = y_1^2 + 3y_2^2 + 6y_3^2.$$

Step 10:-

Nature of the Q.F is positive definite.

Problem 3:-

Reduce the quadratic form $2xy - 2yz + 2xz$ into canonical form by an orthogonal transformation & find its nature, index, rank, & signature.

Soln:- Given Q = $2xy - 2yz + 2xz$ Step 1:- Matrix form:-

$$A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & -1 \\ 1 & -1 & 0 \end{bmatrix}$$

Step 2:- To find the char. eqn:-

Let the char. eqn be

$$\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$$

$$S_1 = 0$$

$$S_2 = \begin{vmatrix} 0 & -1 \\ -1 & 0 \end{vmatrix} + \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} + \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix}$$

$$= (0-1) + (0-1) + (0-1) = -3$$

$$S_3 = \begin{vmatrix} 0 & 1 & 1 \\ 1 & 0 & -1 \\ 1 & -1 & 0 \end{vmatrix}$$

$$= 0(0-1) - 1(0+1) + 1(1-0)$$

$$= -2$$

∴ The char eqn is $\lambda^3 - 3\lambda + 2 = 0$.

Step 3:- To find Eigen values:-

$$\lambda = 1 \Rightarrow 1 - 3 + 2 = 0$$

$\therefore \lambda = 1$ is a root

$$\begin{array}{c|cccc} 1 & 1 & 0 & -3 & 2 \\ & 0 & 1 & 1 & -2 \\ \hline & 1 & 1 & -2 & 0 \end{array}$$

$$\lambda^2 + \lambda - 2 = 0$$

$$(\lambda + 2)(\lambda - 1) = 0$$

$$\lambda = -2, 1$$

$\therefore$ The Eigen values are
-2, 1, 1.

Step 4:- To find Eigen Vectors

Solve $(A - \lambda I)X = 0$

$$\begin{bmatrix} 0 - \lambda & 1 & 1 \\ 1 & 0 - \lambda & -1 \\ 1 & -1 & 0 - \lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\left. \begin{array}{l} -\lambda x_1 + x_2 + x_3 = 0 \\ x_1 - \lambda x_2 - x_3 = 0 \\ x_1 - x_2 - \lambda x_3 = 0 \end{array} \right\} \text{--- (I)}$$

Case(i) $\lambda = -2$ in (I)

$$2x_1 + x_2 + x_3 = 0 \rightarrow \textcircled{1}$$

$$x_1 + 2x_2 - x_3 = 0 \rightarrow \textcircled{2}$$

$$x_1 - x_2 + 2x_3 = 0 \rightarrow \textcircled{3}$$

Solve $\textcircled{1}$ & $\textcircled{2}$

$$\frac{x_1}{\begin{vmatrix} 1 & 1 \\ 2 & -1 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 1 & 2 \\ -1 & 1 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix}}$$

$$\frac{x_1}{-3} = \frac{x_2}{3} = \frac{x_3}{3}$$

$$\therefore x_1 = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}$$

Case(ii) put $\lambda = 1$ in (I)

$$-x_1 + x_2 + x_3 = 0 \rightarrow \textcircled{4}$$

$$x_1 - x_2 - x_3 = 0 \rightarrow \textcircled{5}$$

$$x_1 - x_2 - x_3 = 0 \rightarrow \textcircled{6}$$

All eqn are same.

$$\text{Take } x_1 - x_2 - x_3 = 0$$

$$\text{Put } x_2 = 0 \Rightarrow x_1 = x_3$$

$$\therefore x_2 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

Case(iii)

Let $x_3 = \begin{bmatrix} l \\ m \\ n \end{bmatrix}$ as x_3 is orthogonal to x_1 & x_2 :

$$-l + m + n = 0 \rightarrow \textcircled{7}$$

$$l + m + n = 0 \rightarrow \textcircled{8}$$

Solve $\textcircled{7}$ & $\textcircled{8}$

$$\frac{l}{\begin{vmatrix} 1 & 1 \\ 0 & 1 \end{vmatrix}} = \frac{m}{\begin{vmatrix} 1 & -1 \\ 1 & 1 \end{vmatrix}} = \frac{n}{\begin{vmatrix} -1 & 1 \\ 1 & 0 \end{vmatrix}}$$

$$\frac{l}{1} = \frac{m}{1+1} = \frac{n}{-1}$$

$$\therefore x_3 = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

Step 6:- Normalised matrix N:-

$$N = \begin{bmatrix} -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & 0 & \frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} \end{bmatrix}$$

Step - 6 Find N^T

$$N^T = \begin{bmatrix} -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} & -\frac{1}{\sqrt{6}} \end{bmatrix}$$

Step 7:- Find AN

$$AN = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & -1 \\ 1 & -1 & 0 \end{bmatrix} \begin{bmatrix} -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & 0 & \frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{0+1+1}{\sqrt{3}} & \frac{0+0+1}{\sqrt{2}} & \frac{0+2-1}{\sqrt{6}} \\ \frac{-1+0-1}{\sqrt{3}} & \frac{1+0-1}{\sqrt{2}} & \frac{1+0+1}{\sqrt{6}} \\ \frac{-1-1+0}{\sqrt{3}} & \frac{1+0+0}{\sqrt{2}} & \frac{1-2+0}{\sqrt{6}} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{2}{\sqrt{3}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{3}} & 0 & \frac{2}{\sqrt{6}} \\ -\frac{2}{\sqrt{3}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} \end{bmatrix}$$

Step 8:- Find $D = N^T A N$

$$D = \begin{bmatrix} -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} & -\frac{1}{\sqrt{6}} \end{bmatrix} \begin{bmatrix} \frac{2}{\sqrt{3}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{3}} & 0 & \frac{2}{\sqrt{6}} \\ -\frac{2}{\sqrt{3}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{-2-2-2}{3} & \frac{-1+0+1}{\sqrt{6}} & \frac{-1+2-1}{\sqrt{18}} \\ \frac{2+0-2}{\sqrt{6}} & \frac{1+0+1}{2} & \frac{1+0-1}{\sqrt{12}} \\ \frac{2-4+2}{\sqrt{18}} & \frac{1+0-1}{\sqrt{12}} & \frac{1+4+1}{6} \end{bmatrix}$$

$$= \begin{bmatrix} -\frac{2}{3} & 0 & 0 \\ 0 & \frac{2}{2} & 0 \\ 0 & 0 & \frac{6}{6} \end{bmatrix}$$

$$= \begin{bmatrix} -2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Step 9:- canonical form

$$(y_1 y_2 y_3) \begin{pmatrix} -2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

$$= -2y_1^2 + y_2^2 + y_3^2.$$

Step 10:- Nature of the Quadratic form is IndefiniteIndex:-

No. of positive square terms in the canonical form = 2.

Rank:-

No. of square terms in the canonical form = 3

signature:-

Difference between No. of positive & negative square terms

$$= 2 - 1$$

$$= 1.$$

Problem 4:-

Reduce the quadratic form $x_1^2 + 2x_1x_2 + x_3^2 - 2x_1x_2 + 2x_2x_3$ to the Canonical form through an orthogonal transformation and hence show that it is Positive Semi definite. Also give a non-zero set of values (x_1, x_2, x_3) which makes this quadratic form zero.

Soln:-Step 1: Matrix form

$$A = \begin{bmatrix} 1 & -1 & 0 \\ -1 & 2 & 1 \\ 0 & 1 & 1 \end{bmatrix}$$

Step 2:-

Let the char eqn be $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$$S_1 = 1 + 2 + 1 = 4$$

$$S_2 = \begin{vmatrix} 2 & 1 \\ 1 & 1 \end{vmatrix} + \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} + \begin{vmatrix} 1 & -1 \\ -1 & 2 \end{vmatrix}$$

$$= (2-1) + (1-0) + (2-1) = 3$$

$$S_3 = \begin{vmatrix} 1 & -1 & 0 \\ -1 & 2 & 1 \\ 0 & 1 & 1 \end{vmatrix}$$

$$= 1(2-1) + 1(-1-0) + 0$$

$$= 0$$

$\therefore$ The char. eqn. is $\lambda^3 - 4\lambda^2 + 3\lambda = 0$

$$\text{Step 3:- } \lambda^3 - 4\lambda^2 + 3\lambda = 0$$

$$\lambda(\lambda^2 - 4\lambda + 3) = 0$$

$$\lambda(\lambda-3)(\lambda-1) = 0$$

$$\lambda = 0, 1, 3$$

$\therefore$ Eigen values are 0, 1, 3.

Step 4:-

To find the Eigen vectors

$$\text{Solve } (A - \lambda I)x = 0$$

$$\begin{bmatrix} 1-\lambda & -1 & 0 \\ -1 & 2-\lambda & 1 \\ 0 & 1 & 1-\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\left. \begin{aligned} (1-\lambda)x_1 - x_2 + 0x_3 &= 0 \\ -x_1 + (2-\lambda)x_2 + x_3 &= 0 \\ 0x_1 + x_2 + (1-\lambda)x_3 &= 0 \end{aligned} \right\} \rightarrow \textcircled{I}$$

Case (i) Put $\lambda = 0$ in $\textcircled{I}$

$$x_1 - x_2 + 0x_3 = 0 \rightarrow \textcircled{1}$$

$$-x_1 + 2x_2 + x_3 = 0 \rightarrow \textcircled{2}$$

$$0x_1 + x_2 + (1-\lambda)x_3 = 0 \rightarrow \textcircled{3}$$

Solve $\textcircled{1}$ & $\textcircled{2}$

$$\frac{x_1}{\begin{vmatrix} -1 & 0 \\ 2 & 1 \end{vmatrix}} = \frac{x_2}{\begin{vmatrix} 0 & 1 \\ 1 & -1 \end{vmatrix}} = \frac{x_3}{\begin{vmatrix} 1 & -1 \\ -1 & 2 \end{vmatrix}}$$

$$\frac{x_1}{-1} = \frac{x_2}{-1} = \frac{x_3}{2-1}$$

$$\therefore x_1 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$$

Case(ii) Put $\lambda = 1$ in (I)

$$0x_1 - x_2 + 0x_3 = 0 \rightarrow (4)$$

$$-x_1 + x_2 + x_3 = 0 \rightarrow (5)$$

$$0x_1 + x_2 + 0x_3 = 0 \rightarrow (6)$$

Solve (4) & (5)

$$\frac{x_1}{-1} = \frac{x_2}{0} = \frac{x_3}{-1}$$

$$\begin{vmatrix} -1 & 0 \\ 1 & 1 \end{vmatrix} \quad \begin{vmatrix} 0 & 0 \\ 1 & -1 \end{vmatrix} \quad \begin{vmatrix} 0 & -1 \\ -1 & 1 \end{vmatrix}$$

$$\frac{x_1}{-1} = \frac{x_2}{0} = \frac{x_3}{-1}$$

$$\therefore x_2 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

Case(iii) Put $\lambda = 3$ in (I)

$$-2x_1 - x_2 + 0x_3 = 0 \rightarrow (7)$$

$$-x_1 - x_2 + x_3 = 0 \rightarrow (8)$$

$$0x_1 + x_2 - 2x_3 = 0 \rightarrow (9)$$

$$\frac{x_1}{-1} = \frac{x_2}{2} = \frac{x_3}{2-1}$$

$$\begin{vmatrix} -1 & 0 \\ -1 & 1 \end{vmatrix} \quad \begin{vmatrix} 0 & 2 \\ 1 & -1 \end{vmatrix} \quad \begin{vmatrix} -2 & -1 \\ -1 & -1 \end{vmatrix}$$

$$\frac{x_1}{-1} = \frac{x_2}{2} = \frac{x_3}{2-1}$$

$$\frac{x_1}{-1} = \frac{x_2}{2} = \frac{x_3}{1}$$

$$\therefore x_3 = \begin{bmatrix} -1 \\ 2 \\ 1 \end{bmatrix}$$

Step-5:- Normalised matrix N:

$$N = \begin{bmatrix} \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & 0 & \frac{2}{\sqrt{6}} \\ -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \end{bmatrix}$$

Step-6:- Find N^T

$$N^T = \begin{bmatrix} \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{6}} \\ -\frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{6}} \end{bmatrix}$$

Step-7:- Find AN

$$AN = \begin{bmatrix} 1 & -1 & 0 \\ -1 & 2 & 1 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{2}} & 0 & \frac{2}{\sqrt{6}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{1-1+0}{3} & \frac{1+0+0}{\sqrt{2}} & \frac{-1-2+0}{\sqrt{6}} \\ \frac{-1+2-1}{\sqrt{3}} & \frac{-1+0+1}{\sqrt{2}} & \frac{1+4+1}{\sqrt{6}} \\ \frac{0+1-1}{\sqrt{3}} & \frac{0+0+1}{\sqrt{2}} & \frac{0+2+1}{\sqrt{6}} \end{bmatrix}$$

$$= \begin{bmatrix} 0 & \frac{1}{\sqrt{2}} & -\frac{3}{\sqrt{6}} \\ 0 & 0 & \frac{6}{\sqrt{6}} \\ 0 & \frac{1}{\sqrt{2}} & \frac{3}{\sqrt{6}} \end{bmatrix}$$

Step 8:- Find $D = N^T AN$

$$D = \begin{bmatrix} \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{6}} \\ -\frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{6}} \end{bmatrix} \begin{bmatrix} 0 & \frac{1}{\sqrt{2}} & -\frac{3}{\sqrt{6}} \\ 0 & 0 & \frac{6}{\sqrt{6}} \\ 0 & \frac{1}{\sqrt{2}} & \frac{3}{\sqrt{6}} \end{bmatrix}$$

$$= \begin{bmatrix} 0 & \frac{1+0-1}{\sqrt{6}} & \frac{-3+6-3}{\sqrt{18}} \\ 0 & \frac{1+0+1}{2} & \frac{-3+0+3}{\sqrt{12}} \\ 0 & \frac{-1+0+1}{\sqrt{12}} & \frac{3+2+3}{6} \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

Step-9:- canonical form:-

$$(y_1, y_2, y_3) D \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = 0y_1^2 + y_2^2 + 3y_3^2$$

Step-10:-

Nature of the Q.F:-
canonical form contains both Positive and atleast one Zero.

∴ It is positive definite.
To find non-zero set of values (x_1, x_2, x_3) which makes this quadratic form zero.
when $y_2=0$, $y_3=0$ and y_1 is arbitrary, the orthogonal transformation $x = Ny$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{3} & 1/\sqrt{2} & -1/\sqrt{6} \\ 1/\sqrt{3} & 0 & 2/\sqrt{6} \\ -1/\sqrt{3} & 1/\sqrt{2} & 1/\sqrt{6} \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

$$x_1 = 1/\sqrt{3} y_1 + 1/\sqrt{2} y_2 - 1/\sqrt{6} y_3$$

$$x_2 = 1/\sqrt{3} y_1 + 2/\sqrt{6} y_3$$

$$x_3 = -1/\sqrt{3} y_1 + 1/\sqrt{2} y_2 + 1/\sqrt{6} y_3$$

Taking $y_1 = \sqrt{3}$, $y_2 = 0$, and $y_3 = 0$
we get $x_1 = 1$, $x_2 = 1$, $x_3 = -1$

These values x_1, x_2, x_3 makes the Q.F zero.

Problem:5:-

The Eigen values of a 3×3 real symmetric matrices

A corresponding to the Eigen values 2, 3, 6 are $[1, 0, -1]^T$, $[1, 1, 1]^T$ and $[-1, 2, -1]^T$ respectively, find the matrix A.

Soln:-

Given: Eigen values are 2, 3, 6

Eigen vectors are $\begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$, $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$, $\begin{bmatrix} -1 \\ 2 \\ -1 \end{bmatrix}$

The Normalised matrix

$$N = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{3} & -1/\sqrt{6} \\ 0 & 1/\sqrt{3} & 2/\sqrt{6} \\ -1/\sqrt{2} & 1/\sqrt{3} & -1/\sqrt{6} \end{bmatrix}$$

$$N^T = \begin{bmatrix} 1/\sqrt{2} & 0 & -1/\sqrt{2} \\ 1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} \\ -1/\sqrt{6} & 2/\sqrt{6} & -1/\sqrt{6} \end{bmatrix}$$

$$\text{W.K.T } D = N^T A N$$

$$N D = N N^T A N$$

$$N D N^T = (N N^T) A (N N^T)$$

$$N D N^T = A$$

$$\Rightarrow A = N D N^T$$

$$= \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{3} & -1/\sqrt{6} \\ 0 & 1/\sqrt{3} & 2/\sqrt{6} \\ -1/\sqrt{2} & 1/\sqrt{3} & -1/\sqrt{6} \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 6 \end{bmatrix}$$

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & 0 & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} & -\frac{1}{\sqrt{6}} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{2}{\sqrt{2}} & \frac{3}{\sqrt{3}} & -\frac{6}{\sqrt{6}} \\ 0 & \frac{3}{\sqrt{3}} & \frac{12}{\sqrt{6}} \\ -\frac{2}{\sqrt{2}} & \frac{3}{\sqrt{3}} & -\frac{6}{\sqrt{6}} \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & 0 & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} & -\frac{1}{\sqrt{6}} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{2}{2} + \frac{3}{3} + \frac{6}{6} & 0 + \frac{3}{3} - \frac{12}{6} & -\frac{2}{2} + \frac{3}{3} + \frac{6}{6} \\ 0 + \frac{3}{3} - \frac{12}{6} & 0 + \frac{3}{3} + \frac{24}{6} & 0 + \frac{3}{3} - \frac{12}{6} \\ -\frac{2}{2} + \frac{3}{3} + \frac{6}{6} & 0 + \frac{3}{3} - \frac{12}{6} & \frac{2}{2} + \frac{3}{3} + \frac{6}{6} \end{bmatrix}$$

$$= \begin{bmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{bmatrix}$$

Cayley Hamilton theorem:-

Every square matrix satisfies its own characteristic equation.

Uses:-

- * To calculate
- (i) the positive integral powers of A.
- (ii) The inverse of a non-singular square matrix A.

Problem 1:-

Using Cayley-Hamilton thm find A^{-1} . if $A = \begin{bmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix}$

Soln:-

$$\text{Let } A = \begin{bmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix}$$

Let the char eqn be $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$$S_1 = 2 + 1 + 2 = 5$$

$$S_2 = \begin{vmatrix} 1 & 0 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix}$$

$$= (2-0) + (4-1) + (2-0)$$

$$= 7$$

$$S_3 = \begin{vmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{vmatrix}$$

$$= 2(2-0) - 1(0-0) + 1(0-1)$$

$$= 4 - 1 = 3$$

$$\therefore \text{char. eqn is } \lambda^3 - 5\lambda^2 + 7\lambda - 3 = 0$$

"By Cayley Hamilton thm,

Every square matrix satisfies its own characteristic equation."

Put $\lambda = A$

$$\therefore A^3 - 5A^2 + 7A - 3I = 0 \quad \text{--- (1)}$$

multiply by A^{-1}

$$A^2 - 5A + 7I - 3A^{-1} = 0$$

$$\Rightarrow 3A^{-1} = A^2 - 5A + 7I$$

$$A^{-1} = \frac{1}{3} [A^2 - 5A + 7I] \quad \text{--- (2)}$$

Now, $A^2 = A \cdot A$

$$= \begin{bmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 4+0+1 & 2+1+1 & 2+0+2 \\ 0+0+0 & 0+1+0 & 0+0+0 \\ 2+0+2 & 1+1+2 & 1+0+4 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & 4 & 4 \\ 0 & 1 & 0 \\ 4 & 4 & 5 \end{bmatrix}$$

$\therefore A^2 - 5A + 7I$

$$= \begin{bmatrix} 5 & 4 & 4 \\ 0 & 1 & 0 \\ 4 & 4 & 5 \end{bmatrix} - \begin{bmatrix} 10 & 5 & 5 \\ 0 & 5 & 0 \\ 5 & 5 & 10 \end{bmatrix} + \begin{bmatrix} 7 & 0 & 0 \\ 0 & 7 & 0 \\ 0 & 0 & 7 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & -1 & -1 \\ 0 & 3 & 0 \\ -1 & -1 & 2 \end{bmatrix}$$

$\therefore \textcircled{2} \Rightarrow A^{-1} = \frac{1}{3} \begin{bmatrix} 2 & -1 & -1 \\ 0 & 3 & 0 \\ -1 & -1 & 2 \end{bmatrix}$

Problem 2:-

Using Cayley Hamilton thm,

find A^{-1} if $A = \begin{bmatrix} 1 & 1 & 3 \\ 1 & 3 & -3 \\ -2 & -4 & -4 \end{bmatrix}$

Soln:-

Let $A = \begin{bmatrix} 1 & 1 & 3 \\ 1 & 3 & -3 \\ -2 & -4 & -4 \end{bmatrix}$

Let the char. eqn be $\lambda^3 - s_1\lambda^2 + s_2\lambda - s_3 = 0$

$$s_1 = 1 + 3 - 4 = 0$$

$$s_2 = \begin{vmatrix} 3 & -3 \\ -4 & -4 \end{vmatrix} + \begin{vmatrix} 1 & 3 \\ -2 & -4 \end{vmatrix} + \begin{vmatrix} 1 & 1 \\ 1 & 3 \end{vmatrix}$$

$$= (-12 - 12) + (-4 - 6) + (3 - 1)$$

$$= -20$$

$$s_3 = \begin{vmatrix} 1 & 1 & 3 \\ 1 & 3 & -3 \\ -2 & -4 & -4 \end{vmatrix}$$

$$= 1(-12 - 12) - 1(4 - 6) + 3(-4 + 6)$$

$$= -24 + 10 + 6$$

$$= -8$$

$\therefore$ The char. eqn is $\lambda^3 - 0\lambda^2 - 20\lambda + 8 = 0$

By Cayley Hamilton thm,

"Every square matrix satisfies its own characteristic eqn."

$$\therefore A^3 - 20A + 8I = 0$$

x by A^{-1}

$$\therefore A^2 - 20I + 8A^{-1} = 0$$

$$-8A^{-1} = A^2 - 20I$$

$$A^{-1} = -\frac{1}{8} [A^2 - 20I]$$

Now

$$A^2 = \begin{bmatrix} 1 & 1 & 3 \\ 1 & 3 & -3 \\ -2 & -4 & -4 \end{bmatrix} \begin{bmatrix} 1 & 1 & 3 \\ 1 & 3 & -3 \\ -2 & -4 & -4 \end{bmatrix}$$

$$= \begin{bmatrix} 1+1-6 & 1+3-12 & 3-3-12 \\ 1+3+6 & 1+9+12 & 3-9+12 \\ -2-4+8 & -2-12+16 & -6+12+16 \end{bmatrix}$$

$$= \begin{bmatrix} -4 & -8 & -12 \\ 10 & 22 & 6 \\ 2 & 2 & 22 \end{bmatrix}$$

$$\therefore A^2 - 20I = \begin{bmatrix} -4 & -8 & -12 \\ 10 & 22 & 6 \\ 2 & 2 & 22 \end{bmatrix} + \begin{bmatrix} -20 & 0 & 0 \\ 0 & -20 & 0 \\ 0 & 0 & -20 \end{bmatrix}$$

$$= \begin{bmatrix} -24 & -8 & -12 \\ 10 & 2 & 0 \\ 2 & 2 & 2 \end{bmatrix}$$

$$\therefore \textcircled{1} \Rightarrow A^T = \frac{-1}{8} \begin{bmatrix} -24 & -8 & -12 \\ 10 & 2 & 0 \\ 2 & 2 & 2 \end{bmatrix}$$

Problem 3:-

Verify Cayley Hamilton thm, find A^4 and A^T when

$$A = \begin{bmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

Soln:-

$$\text{Given } A = \begin{bmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

Let the char. eqn be

$$\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$$

$$S_1 = 2+2+2 = 6$$

$$S_2 = \begin{vmatrix} 2 & -1 \\ -1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 2 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & -1 \\ -1 & 2 \end{vmatrix}$$

$$= (4-1) + (4-2) + (4-1)$$

$$= 8$$

$$S_3 = \begin{vmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{vmatrix}$$

$$= 2(4-1) + 1(-2+1) + 2(1-2)$$

$$= 6-1-2$$

$$= 3$$

$$\therefore \text{char eqn is } \lambda^3 - 6\lambda^2 + 8\lambda - 3 = 0$$

$\therefore$ By Cayley Hamilton theorem,

"Every square matrix satisfies its own characteristic equation."

$$\therefore A^3 - 6A^2 + 8A - 3I = 0 \rightarrow \textcircled{1}$$

Verification:-

$$A^2 = \begin{bmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix} \begin{bmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 4+1+2 & -2-2-2 & 4+1+4 \\ -2-2-1 & 1+4+1 & -2-2-2 \\ 2+1+2 & -1-2-2 & 2+1+4 \end{bmatrix}$$

$$= \begin{bmatrix} 7 & -6 & 9 \\ -5 & 6 & -6 \\ 5 & -5 & 7 \end{bmatrix}$$

$$A^3 = A^2 \cdot A = \begin{bmatrix} 7 & -6 & 9 \\ -5 & 6 & -6 \\ 5 & -5 & 7 \end{bmatrix} \begin{bmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 14+6+9 & -7-12-9 & 14+6+18 \\ -10-6-6 & 5+12+6 & -10-6-12 \\ 10+5+7 & -5+10-7 & 10+5+14 \end{bmatrix}$$

$$= \begin{bmatrix} 29 & -28 & 38 \\ -22 & 23 & -28 \\ 22 & -22 & 29 \end{bmatrix}$$

$$\therefore A^3 - 6A^2 + 8A - 3I$$

$$= \begin{bmatrix} 29 & -28 & 38 \\ -22 & 23 & -28 \\ 22 & -22 & 29 \end{bmatrix} - 6 \begin{bmatrix} 7 & -6 & 9 \\ -5 & 6 & -6 \\ 5 & -5 & 7 \end{bmatrix}$$

$$+ 8 \begin{bmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix} - 3 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 29 & -28 & 38 \\ -22 & 23 & -28 \\ 22 & -22 & 29 \end{bmatrix} + \begin{bmatrix} -42 & 36 & -54 \\ 30 & -36 & 36 \\ -30 & 30 & -42 \end{bmatrix}$$

$$+ \begin{bmatrix} 16 & -8 & 16 \\ -8 & 16 & -8 \\ 8 & -8 & 16 \end{bmatrix} + \begin{bmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Hence verified.

To find A^4 :-

$$\textcircled{1} XA \Rightarrow A^4 - 6A^3 + 8A^2 - 3A = 0$$

$$A^4 = 6A^3 - 8A^2 + 3A$$

$$= 6 \begin{bmatrix} 29 & -28 & 38 \\ -22 & 23 & -28 \\ 22 & -22 & 29 \end{bmatrix} - 8 \begin{bmatrix} 7 & -6 & 9 \\ -5 & 6 & -6 \\ 5 & -5 & 7 \end{bmatrix}$$

$$+ 3 \begin{bmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 174 & -168 & 228 \\ -132 & 138 & -168 \\ 132 & -132 & 174 \end{bmatrix}$$

$$+ \begin{bmatrix} -56 & 48 & -72 \\ 40 & -48 & 48 \\ -40 & 40 & -56 \end{bmatrix} + \begin{bmatrix} 6 & -3 & 6 \\ -3 & 6 & -3 \\ 3 & -3 & 6 \end{bmatrix}$$

$$= \begin{bmatrix} 124 & -123 & 162 \\ -95 & 96 & -123 \\ 95 & -95 & 124 \end{bmatrix}$$

To find A^{-1}

$$\textcircled{1} XA^T \Rightarrow A^2 - 6A + 8I - 3A^T = 0$$

$$\Rightarrow 3A^T = A^2 - 6A + 8I$$

$$\Rightarrow A^T = \frac{1}{3} [A^2 - 6A + 8I] \rightarrow \textcircled{2}$$

Now $A^2 - 6A + 8I$

$$= \begin{bmatrix} 7 & -6 & 9 \\ -5 & 6 & -6 \\ 5 & -5 & 7 \end{bmatrix} + \begin{bmatrix} -12 & 6 & -12 \\ 6 & -12 & 6 \\ -6 & 6 & 12 \end{bmatrix} + \begin{bmatrix} 8 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & 8 \end{bmatrix}$$

$$= \begin{bmatrix} 3 & 0 & -3 \\ 1 & 2 & 0 \\ -1 & 1 & 3 \end{bmatrix} \therefore \textcircled{2} \Rightarrow A^T = \frac{1}{3} \begin{bmatrix} 3 & 0 & -3 \\ 1 & 2 & 0 \\ -1 & 1 & 3 \end{bmatrix}$$

Problem -4:-

use Cayley Hamilton thm,
to find the value of the matrix
given by

$$(i) f(A) = A^8 - 5A^7 + 7A^6 - 3A^5 + A^4 - 5A^3 + 8A^2 - 2A + I$$

$$(ii) A^8 - 5A^7 + 7A^6 - 3A^5 + 8A^4 - 5A^3 + 8A^2 - 2A + I$$

$$\text{if } A = \begin{bmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix}$$

Soln:-

Given

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix}$$

Let the char. eqn be

$$\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$$

$$S_1 = 2 + 1 + 2 = 5$$

$$S_2 = \begin{vmatrix} 1 & 0 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix}$$

$$= (2-0) + (4-1) + (2-0) \\ = 7$$

$$S_3 = \begin{vmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{vmatrix}$$

$$= 2(2-0) - (0-1) + 1(0-1)$$

$$= 3$$

$$\therefore \text{char eqn is } \lambda^3 - 5\lambda^2 + 7\lambda - 3 = 0$$

By Cayley Hamilton thm,

"Every square matrix satisfies its own characteristic equation."

$$\therefore A^3 - 5A^2 + 7A - 3I = 0 \rightarrow (1)$$

$$\text{Let } f(A) = A^8 - 5A^7 + 7A^6 - 3A^5 + A^4 - 5A^3 + 8A^2 - 2A + I$$

$$\begin{array}{r} A^5 + A \\ A^3 - 5A^2 + 7A - 3I \overline{) A^8 - 5A^7 + 7A^6 - 3A^5 + A^4 - 5A^3 + 8A^2 - 2A + I} \\ \underline{A^8 - 5A^7 + 7A^6 - 3A^5} \\ A^4 - 5A^3 + 8A^2 - 2A + I \\ \underline{A^4 - 5A^3 + 7A^2 - 3A} \\ A^2 + A + I \end{array}$$

$$\therefore f(A) = (A^3 - 5A^2 + 7A - 3I)(A^5 + A) + (A^2 + A + I)$$

$$= 0(A^5 + A) + A^2 + A + I \\ = A^2 + A + I$$

$$\text{Now } A^2 = \begin{bmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 4+0+1 & 2+1+1 & 2+0+2 \\ 0+0+0 & 0+1+0 & 0+0+0 \\ 2+0+2 & 1+1+2 & 1+0+4 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & 4 & 4 \\ 0 & 1 & 0 \\ 4 & 4 & 5 \end{bmatrix}$$

$$\therefore f(A) = \begin{bmatrix} 5 & 4 & 4 \\ 0 & 1 & 0 \\ 4 & 4 & 5 \end{bmatrix} + \begin{bmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 8 & 5 & 5 \\ 0 & 3 & 0 \\ 5 & 5 & 8 \end{bmatrix}$$

$$\text{Let } g(A) = A^8 - 5A^7 + 7A^6 - 3A^5 + 8A^4 - 5A^3 + 8A^2 - 2A + I$$

$$\begin{array}{r} A^5 + 8A + 35I \\ A^3 - 5A^2 + 7A - 3I \quad \times \quad A^5 + 8A + 35I \\ \hline A^8 - 5A^7 + 7A^6 - 3A^5 + 8A^4 - 5A^3 + 8A^2 - 2A + I \\ - (A^8 - 5A^7 + 7A^6 - 3A^5) \\ \hline 8A^4 - 5A^3 + 8A^2 - 2A + I \\ (-) 8A^4 - 40A^3 + 50A^2 - 24A \\ \hline 35A^3 - 48A^2 + 22A + I \\ (-) 35A^3 - 175A^2 + 245A - 105I \\ \hline 127A^2 - 223A + 106I \end{array}$$

$$\therefore g(A) = (A^3 - 5A^2 + 7A - 3I)(A^5 + 8A + 35I) + 127A^2 - 223A + 106I$$

$$= 127A^2 - 223A + 106I$$

$$= 127 \begin{bmatrix} 5 & 4 & 4 \\ 0 & 1 & 0 \\ 4 & 4 & 5 \end{bmatrix} - 223 \begin{bmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix} + 106 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 635 & 508 & 508 \\ 0 & 127 & 0 \\ 508 & 508 & 635 \end{bmatrix}$$

$$+ \begin{bmatrix} -446 & -223 & -233 \\ 0 & -223 & 0 \\ -223 & -223 & -446 \end{bmatrix}$$

$$+ \begin{bmatrix} 106 & 0 & 0 \\ 0 & 106 & 0 \\ 0 & 0 & 106 \end{bmatrix}$$

$$= \begin{bmatrix} 295 & 285 & 285 \\ 0 & 10 & 0 \\ 285 & 285 & 295 \end{bmatrix}.$$

Problem-5:-

Verify Cayley-Hamilton theorem and find its inverse of

$$A = \begin{bmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{bmatrix}$$

NID-2019

Soln:-

$$\text{Given } A = \begin{bmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{bmatrix}$$

Let the char. eqn be $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$$S_1 = 3 + 5 + 3 = 11$$

$$S_2 = \begin{vmatrix} 3 & -1 \\ -1 & 5 \end{vmatrix} + \begin{vmatrix} 3 & 1 \\ 1 & 3 \end{vmatrix} + \begin{vmatrix} 3 & -1 \\ -1 & 5 \end{vmatrix}$$

$$= (15 - 1) + (9 - 1) + (15 - 1)$$

$$= 14 + 8 + 14 = 36$$

$$S_3 = \begin{vmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{vmatrix}$$

$$= 3(15 - 1) + 1(-3 + 1) + 1(1 - 5)$$

$$= 3(14) + 1(-2) + 1(-4) = 36$$

∴ The char. eqn is

$$\lambda^3 - 11\lambda^2 + 36\lambda - 36 = 0$$

By Cayley-Hamilton theorem,

"Every square matrix satisfies its own characteristic equation"

$$\therefore A^3 - 11A^2 + 36A - 36I = 0$$

Verification.

$$A^2 = \begin{bmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{bmatrix} \begin{bmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 9+1+1 & -3-5-1 & 3+1+3 \\ -3-5-1 & +1+25+1 & -1+5-3 \\ 3+1+3 & -1-5-3 & 1+1+9 \end{bmatrix}$$

$$= \begin{bmatrix} 11 & -9 & 7 \\ -9 & 27 & -9 \\ 7 & -9 & 11 \end{bmatrix}$$

$$A^3 = A^2 \cdot A = \begin{bmatrix} 11 & -9 & 7 \\ -9 & 27 & -9 \\ 7 & -9 & 11 \end{bmatrix} \begin{bmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 33+9+7 & -11-45-7 & 11+9+21 \\ -27-27-9 & 9+135+9 & -9-27-27 \\ 21+9+11 & -7-45-11 & 7+9+33 \end{bmatrix}$$

$$= \begin{bmatrix} 49 & -63 & 41 \\ -63 & 153 & -63 \\ 41 & -63 & 49 \end{bmatrix}$$

$$\therefore A^3 - 11A^2 + 36A - 36I$$

$$= \begin{bmatrix} 49 & -63 & 41 \\ -63 & 153 & -63 \\ 41 & -63 & 49 \end{bmatrix} - \begin{bmatrix} 121 & -99 & 77 \\ -99 & 297 & -99 \\ 77 & -99 & 121 \end{bmatrix}$$

$$+ \begin{bmatrix} 108 & -36 & 36 \\ -36 & 180 & -36 \\ 36 & -36 & 108 \end{bmatrix} - \begin{bmatrix} 36 & 0 & 0 \\ 0 & 36 & 0 \\ 0 & 0 & 36 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

To find A^{-1}

$$\text{X by } A^{-1} \Rightarrow A^2 - 11A + 36I - 36A^{-1} = 0$$

$$36A^{-1} = A^2 - 11A + 36I$$

$$= \begin{bmatrix} 11 & -9 & 7 \\ -9 & 27 & -9 \\ 7 & -9 & 11 \end{bmatrix} - \begin{bmatrix} 33 & -11 & 11 \\ -11 & 55 & -11 \\ 11 & -11 & 33 \end{bmatrix}$$

$$+ \begin{bmatrix} 36 & 0 & 0 \\ 0 & 36 & 0 \\ 0 & 0 & 36 \end{bmatrix}$$

$$= \begin{bmatrix} 14 & 2 & -4 \\ 2 & 8 & 2 \\ -4 & 2 & 14 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{36} \begin{bmatrix} 14 & 2 & -4 \\ 2 & 8 & 2 \\ -4 & 2 & 14 \end{bmatrix} //$$

Application of Eigen values and Eigen vector- Stretching of an elastic membrane

* For any system of differential equations of the form

$$\frac{dx}{dt} = A X$$

Where A is an $n \times n$ matrix with distinct eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ and t is the independent variable.

$\therefore$ The solution is $X = Y$

where Y is the ormalized matrix of A, and

$$Y = [c_1 e^{\lambda_1 t}, c_2 e^{\lambda_2 t}, \dots, c_n e^{\lambda_n t}]^T$$

Problem:-

Solve the System of differential equation $\frac{dx}{dt} = 4x + 2y$

$$\frac{dy}{dt} = -x + y$$

with initial conditions

$$x(0) = 1, y(0) = 0.$$

Soln:-

$$\text{Let } A = \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$$

The char. eqn is $\lambda^2 - S_1 \lambda + S_2 = 0$

$$S_1 = [4 + 1] = 5$$

$$S_2 = \begin{vmatrix} 4 & 2 \\ -1 & 1 \end{vmatrix} = 4 + 2 = 6$$

$\therefore$ The char. eqn is $\lambda^2 - 5\lambda + 6 = 0$

$$\lambda = 3, \lambda = 2.$$

$\therefore$ The eigen values are 2, 3.

To find Eigen vector

$$(A - \lambda I) X = 0$$

$$\begin{bmatrix} 4 - \lambda & 2 \\ -1 & 1 - \lambda \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 0$$

$$(4 - \lambda)x + 2y = 0 \quad \text{--- (1)}$$

$$-x + (1 - \lambda)y = 0 \quad \text{--- (2)}$$

case(i) put $\lambda = 2$

$$2x + 2y = 0 \quad \text{--- (3)}$$

$$-x - y = 0 \quad \text{--- (4)}$$

The eqn (3) & (4) are same

$$2x = -2y$$

$$\frac{x}{-1} = \frac{y}{1} \quad \therefore x_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

case(ii) put $\lambda = 3$

$$x + 2y = 0 \quad \text{--- (5)}$$

$$-x - 2y = 0 \quad \text{--- (6)}$$

$\therefore$ The eqn (5) & (6) are same

$$x + 2y = 0$$

$$x = -2y \quad \therefore x_2 = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

$$\frac{x}{-2} = \frac{y}{1}$$

$$N = \begin{bmatrix} -\frac{1}{\sqrt{2}} & -\frac{2}{\sqrt{5}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{5}} \end{bmatrix}, \quad N^{-1} = \begin{bmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{bmatrix}$$

$$D = N^{-1} A N = \begin{bmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{bmatrix} \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{\sqrt{2}} & -\frac{2}{\sqrt{5}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{5}} \end{bmatrix}$$

$$= \begin{bmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{bmatrix} \begin{bmatrix} -\frac{2}{\sqrt{2}} & -\frac{6}{\sqrt{5}} \\ \frac{2}{\sqrt{2}} & \frac{3}{\sqrt{5}} \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$$

$$\therefore r(t) = c e^{3t} \quad s(t) = k e^{2t}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = N Y$$

$$= \begin{bmatrix} -\frac{1}{\sqrt{2}} & -\frac{2}{\sqrt{5}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{5}} \end{bmatrix} \begin{bmatrix} r \\ s \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -\frac{1}{\sqrt{2}} r - \frac{2}{\sqrt{5}} s \\ \frac{1}{\sqrt{2}} r + \frac{1}{\sqrt{5}} s \end{bmatrix}$$

$$\therefore x = -\frac{1}{\sqrt{2}} r - \frac{2}{\sqrt{5}} s = -\frac{1}{\sqrt{2}} c e^{3t} - \frac{2}{\sqrt{5}} k e^{2t}$$

$$y = \frac{1}{\sqrt{2}} r + \frac{1}{\sqrt{5}} s = \frac{1}{\sqrt{2}} c e^{3t} + \frac{1}{\sqrt{5}} k e^{2t}$$

$$x(0) = 1, \quad y(0) = 0$$

$$1 = -\frac{1}{\sqrt{2}} c - \frac{2}{\sqrt{5}} k \quad \text{--- (7)}$$

$$0 = \frac{1}{\sqrt{2}} c + \frac{1}{\sqrt{5}} k \quad \text{--- (8)}$$

Solve (7) & (8)

$$\begin{array}{rcl} -\frac{1}{\sqrt{2}} c - \frac{2}{\sqrt{5}} k & = & 1 \\ \frac{1}{\sqrt{2}} c + \frac{1}{\sqrt{5}} k & = & 0 \\ \hline -\frac{1}{\sqrt{5}} k & = & 1 \end{array}$$

$$\therefore \boxed{k = -\sqrt{5}}$$

$$\text{Sub } k = -\sqrt{5} \text{ in (7)}$$

$$1 = -\frac{1}{\sqrt{2}} c + 2$$

$$-1 = -\frac{1}{\sqrt{2}} c$$

$$\boxed{c = \sqrt{2}}$$

$$\therefore x(t) = -\frac{1}{\sqrt{2}} \sqrt{2} e^{3t} - \frac{2}{\sqrt{5}} (-\sqrt{5}) e^{2t}$$

$$\boxed{x(t) = -e^{3t} + 2e^{2t}}$$

$$y(t) = \frac{1}{\sqrt{2}} \sqrt{2} e^{3t} + \frac{1}{\sqrt{5}} (-\sqrt{5}) e^{2t}$$

$$\boxed{y(t) = e^{3t} - e^{2t}}$$

Problem - 2

An elastic membrane in the x_1, x_2 -plane with boundary circle $x_1^2 + x_2^2 = 1$ is stretched

so that a point $P: (x, x_2)$ goes over into the point $Q: (y_1, y_2)$

$$\text{given by } y_1 = 5x_1 + 3x_2$$

$$y_2 = 3x_1 + 5x_2.$$

Find the principal directions, that is, direction of position vector x of P for which the direction of the position vector y of Q is the same (or) exactly opposite. What shape does the boundary circle take under this deformation?

Soln:-

$$\text{Given } y_1 = 5x_1 + 3x_2$$

$$y_2 = 3x_1 + 5x_2$$

$$\text{Let } A = \begin{bmatrix} 5 & 3 \\ 3 & 5 \end{bmatrix}$$

The char. eqn is

$$\lambda^2 - S_1\lambda + S_2 = 0$$

$$S_1 = 5 + 5 = 10$$

$$S_2 = \begin{vmatrix} 5 & 3 \\ 3 & 5 \end{vmatrix} = 25 - 9 = 16$$

 $\therefore$ The char. eqn is

$$\lambda^2 - 10\lambda + 16 = 0$$

$$(\lambda - 8)(\lambda - 2) = 0$$

$$\lambda = 2, 8$$

 $\therefore$ The eigen values are

$$2, 8$$

To find Eigen vector

$$(A - \lambda I)X = 0$$

$$\begin{bmatrix} 5 - \lambda & 3 \\ 3 & 5 - \lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$(5 - \lambda)x_1 + 3x_2 = 0 \quad \text{--- (1)}$$

$$3x_1 + (5 - \lambda)x_2 = 0 \quad \text{--- (2)}$$

Case(i) Put $\lambda = 2$

$$3x_1 + 3x_2 = 0 \quad \text{--- (3)}$$

$$3x_1 + 3x_2 = 0 \quad \text{--- (4)}$$

The eqn (3) & (4) are same

$$3x_1 = -3x_2$$

$$x_1 = -x_2$$

$$\frac{x_1}{-1} = \frac{x_2}{1}$$

$$\therefore x_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Case(ii) put $\lambda = 8$ in (1) & (2).

$$-3x_1 + 3x_2 = 0 \quad \text{--- (5)}$$

$$3x_1 - 3x_2 = 0 \quad \text{--- (6)}$$

The eqn (5) & (6) are same

$$-3x_1 + 3x_2 = 0$$

$$3x_2 = 3x_1$$

$$\frac{x_2}{1} = \frac{x_1}{1}$$

$$\therefore x_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

The ~~the~~ Eigen vectors


$$\lambda = 2 \Rightarrow x_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$\lambda = 8 \Rightarrow x_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

These vectors make 45° and 135° angles with the positive x_1 -direction.

They give the Principal directions

The eigen values show that in the Principal direction the membrane is stretched by factors 8 and 2, respectively; see in Figure



Accordingly, if we choose the principal directions as direction of a new cartesian u, v -coordinate system, say with positive $z_1 = r \cos \theta$ semi axes in the first quadrant and the positive $z_2 = r \sin \theta$ semi axis in the second quadrant of the x_1, x_2 -system, then a boundary point of the unstretched circular membrane has coordinates $\cos \theta, \sin \theta$.

Hence $z_1 = \cos \theta$ — (7)

$z_2 = \sin \theta$. — (8)

(7)² + (8)²
 $\frac{z_1^2}{\cos^2 \theta} + \frac{z_2^2}{\sin^2 \theta} = \cos^2 \theta + \sin^2 \theta$

$\frac{z_1^2}{\cos^2 \theta} + \frac{z_2^2}{\sin^2 \theta} = 1$

$\therefore$ The deformed boundary is an ellipse.

Problem 3:-

Find the solution of the differential equations

$$u = 2x - 2y + 4z$$

$$v = 3y - 2z$$

$$w = -y + 2z$$

with the initial conditions

$$x(0) = 2, y(0) = 0, \text{ and } z(0) = 2.$$

Soln:-

Let $A = \begin{bmatrix} 2 & -2 & 4 \\ 0 & 3 & -2 \\ 0 & -1 & 2 \end{bmatrix}$

$$X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \quad X' = \begin{bmatrix} u \\ v \\ w \end{bmatrix}$$

The char. eqn is $\lambda^3 - S_1 \lambda^2 + S_2 \lambda - S_3 = 0$

$$S_1 = 2 + 3 + 2 = 7$$

$$S_2 = \begin{vmatrix} 2 & -2 \\ 0 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 4 \\ 0 & 2 \end{vmatrix} + \begin{vmatrix} 2 & -2 \\ 0 & 3 \end{vmatrix}$$

$$= 6 - 2 + 4 - 0 + 6 - 0 = 14$$

$$S_3 = \begin{vmatrix} 2 & -2 & 4 \\ 0 & 3 & -2 \\ 0 & -1 & 2 \end{vmatrix}$$

$$= 2(6 - 2) + 2(0 - 0) + 4(0 - 0) = 8$$

$$\therefore \lambda^3 - 7\lambda^2 + 14\lambda - 8 = 0$$

$\lambda = 1$ is a one of the root

$$\begin{array}{ccc|c} 1 & -7 & 14 & -8 \\ 0 & 1 & -6 & 8 \\ \hline 1 & -6 & 8 & 0 \end{array}$$

$$\lambda^2 - 6\lambda + 8 = 0$$

$$\lambda^2 - 4\lambda - 2\lambda + 8 = 0$$

$$(\lambda - 4)(\lambda - 2) = 0$$

$$\lambda = 2, 4.$$

∴ The eigen values are
1, 2, 4

To find Eigen vector

$$\begin{bmatrix} 2-\lambda & -2 & 4 \\ 0 & 3-\lambda & -2 \\ 0 & -1 & 2-\lambda \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\left. \begin{array}{l} (2-\lambda)x - 2y + 4z = 0 \\ 0x + (3-\lambda)y - 2z = 0 \\ 0x - y + (2-\lambda)z = 0 \end{array} \right\} \text{--- (I)}$$

case(i) put $\lambda = 1$ in (I)

$$\left. \begin{array}{l} x - 2y + 4z = 0 \text{ --- (1)} \\ 0x + 2y - 2z = 0 \text{ --- (2)} \\ 0x - y + z = 0 \text{ --- (3)} \end{array} \right\}$$

Solve (1) & (2)

$$\frac{x}{-2} = \frac{y}{4} = \frac{z}{-2}$$

$$\frac{x}{4-8} = \frac{y}{0+4} = \frac{z}{4-0}$$

$$\frac{x}{-4} = \frac{y}{2} = \frac{z}{2}$$

$$\therefore x_1 = \begin{bmatrix} -4 \\ 2 \\ 2 \end{bmatrix} \Rightarrow x_1 = \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix}$$

case(ii) put $\lambda = 2$ in (I)

$$0x - 2y + 4z = 0 \text{ --- (4)}$$

$$0x + y - 2z = 0 \text{ --- (5)}$$

$$0x - y + 0z = 0 \text{ --- (6)}$$

solve (4) & (6)

$$\frac{x}{-2} = \frac{y}{4} = \frac{z}{0}$$

$$\frac{x}{0-4} = \frac{y}{0-0} = \frac{z}{0-0}$$

$$\frac{x}{-4} = \frac{y}{0} = \frac{z}{0}$$

$$x_2 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

case(iii) $\lambda = 4$ in (I)

$$-2x - 2y + 4z = 0 \text{ --- (7)}$$

$$0x - y - 2z = 0 \text{ --- (8)}$$

$$0 - y - 2z = 0 \text{ --- (9)}$$

solve (7) & (8)

$$\frac{x}{-2} = \frac{y}{4} = \frac{z}{-2}$$

$$\frac{x}{8} = \frac{y}{-4} = \frac{z}{2}$$

$$\therefore X_3 = \begin{bmatrix} 4 \\ -2 \\ 1 \end{bmatrix}$$

Since the eigen values of the coefficient matrix A are distinct the corresponding eigenvectors are linearly independent.

This shows that A is diagonalizable and hence

$N = [x_1, x_2, x_3]$ is the diagonalizing matrix.

So we have $N^{-1}AN = D$

$$\text{i.e. } N^{-1}AN = D = \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 4 \end{bmatrix}$$

$$Y = \begin{bmatrix} u(t) \\ v(t) \\ w(t) \end{bmatrix}$$

The relation $X = PY$. then

Since P is Normalized matrix of constants, we have

$$X' = PY'$$

So $X' = AX$ becomes

$$PY' = AX$$

$$= A(NY)$$

$$Y' = (N^{-1}AN)Y$$

$$\text{i.e., } \begin{bmatrix} u'(t) \\ v'(t) \\ w'(t) \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 4 \end{bmatrix} \begin{bmatrix} u(t) \\ v(t) \\ w(t) \end{bmatrix}$$

$$= \begin{bmatrix} u(t) \\ 2v(t) \\ 4w(t) \end{bmatrix}$$

then

$$u(t) = c_1 e^t, \quad v(t) = c_2 e^{2t}$$

$$\text{and } w(t) = c_3 e^{4t}.$$

where c_1, c_2 and c_3 are constants.

$$X = NY$$

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = \begin{bmatrix} -2 & 1 & 4 \\ 1 & 0 & -2 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} u(t) \\ v(t) \\ w(t) \end{bmatrix}$$

$$= \begin{bmatrix} -2 & 1 & 4 \\ 1 & 0 & -2 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} c_1 e^t \\ c_2 e^{2t} \\ c_3 e^{4t} \end{bmatrix}$$

$$x(t) = -2c_1 e^t + c_2 e^{2t} + 4c_3 e^{4t}$$

$$y(t) = c_1 e^t + 0c_2 e^{2t} - 2c_3 e^{4t}$$

$$z(t) = c_1 e^t + 0c_2 e^{2t} + c_3 e^{4t}$$

Given $x(0) = 2$
 $y(0) = 0$
 $z(0) = 2$

$$x(0) = -2c_1 e^0 + c_2 e^0 + 4c_3 e^0$$

$$y(0) = c_1 e^0 + 0c_2 e^0 - 2c_3 e^0$$

$$z(0) = c_1 e^0 + 0c_2 e^0 + c_3 e^0$$

$$2 = -2c_1 + c_2 + 4c_3 \quad \text{--- (10)}$$

$$0 = c_1 - 2c_3 \quad \text{--- (11)}$$

$$2 = c_1 + c_3 \quad \text{--- (12)}$$

Solve (11) & (12)

$$\begin{array}{rcl} c_1 - 2c_3 & = & 0 \\ c_1 + c_3 & = & 2 \\ \hline -3c_3 & = & -2 \end{array}$$

$$\boxed{c_3 = \frac{2}{3}}$$

Sub c_3 in (11)

$$\textcircled{11} \Rightarrow c_1 = 2c_3$$

$$c_1 = 2\left(\frac{2}{3}\right)$$

$$\boxed{c_1 = \frac{4}{3}}$$

$$-2c_1 + c_2 + 4c_3 = 2$$

$$-2\left(\frac{4}{3}\right) + c_2 + 4\left(\frac{2}{3}\right) = 2$$

$$-\frac{8}{3} + c_2 + \frac{8}{3} = 2$$

$$\boxed{c_2 = 2}$$

$$x(t) = -\frac{8}{3}e^t + 2e^{2t} + \frac{8}{3}e^{4t}$$

$$y(t) = \frac{4}{3}e^t - \frac{4}{3}e^{4t}$$

$$z(t) = \frac{2}{3}e^t + \frac{2}{3}e^{4t}$$

∴ By Cayley Hamilton thm,

$$A^2 - 4A - 5I = 0 \rightarrow (1)$$

$$A^2 - 4A - 5I \begin{matrix} A^2 \\ \hline A^4 - 4A^3 - 5A^2 + A + 2I \\ A^4 - 4A^3 - 5A^2 \\ \hline A + 2I \end{matrix}$$

$$\begin{aligned} \therefore A^4 - 4A^3 - 5A^2 + A + 2I &= (A^2 - 4A - 5I)(A^2 + (A + 2I)) \\ &= A + 2I \\ &= \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix} + 2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 3 & 2 \\ 4 & 5 \end{bmatrix} \end{aligned}$$

Problem - 19 :-

Use Cayley-Hamilton theorem to find $A^4 - 8A^3 + 12A^2$ when

$$A = \begin{bmatrix} 5 & 3 \\ 1 & 3 \end{bmatrix}$$

Soln:-

The char. eqn is $\lambda^2 - 8\lambda + 12 = 0$

$$S_1 = 5 + 3 = 8$$

$$S_2 = \begin{vmatrix} 5 & 3 \\ 1 & 3 \end{vmatrix} = 15 - 3 = 12$$

$$\therefore \lambda^2 - 8\lambda + 12 = 0$$

By Cayley-Hamilton thm,

$$A^2 - 8A + 12I = 0$$

$$\times \text{ by } A^2 \quad A^4 - 8A^3 + 12A^2 = 0.$$

Problem - 20 :-

Write the matrix of the quadratic form

$$2x^2 + 8z^2 + 4xy + 10xz - 2yz$$

Soln:-

$$A = \begin{bmatrix} \text{coeff } x^2 & \frac{1}{2} \text{coeff } xy & \frac{1}{2} \text{coeff } xz \\ \frac{1}{2} \text{coeff } yx & \text{coeff } y^2 & \frac{1}{2} \text{coeff } yz \\ \frac{1}{2} \text{coeff } zx & \frac{1}{2} \text{coeff } zy & \text{coeff } z^2 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 2 & 5 \\ 2 & 0 & -1 \\ 5 & -1 & 8 \end{bmatrix}$$

Problem 21:-

Discuss the nature of the following quadratic form
 $3x_1^2 + 3x_2^2 - 5x_3^2 - 2x_1x_2 - 6x_2x_3 - 6x_3x_1$

Soln:-

$$A = \begin{bmatrix} 3 & -1 & -3 \\ -1 & 3 & -3 \\ -3 & -3 & -5 \end{bmatrix}$$

$$D_1 = |3| = 3, \text{ (+ive)}$$

$$D_2 = \begin{vmatrix} 3 & -1 \\ -1 & 3 \end{vmatrix} = 9 - 1 = 8 \text{ (+ive)}$$

$$D_3 = \begin{vmatrix} 3 & -1 & -3 \\ -1 & 3 & -3 \\ -3 & -3 & -5 \end{vmatrix}$$

$$= 3(15 - 9) + 1(5 - 9) - 3(3 - 9)$$

$$= 3(-24) + 1(-4) - 3(-6)$$

$$= -112, \text{ (-ive)}$$

∴ The Q.F is indefinite

Soln:-

$$\text{Let } A = \begin{bmatrix} 2 & 0 & 1 \\ 0 & 2 & 0 \\ a & 0 & 2 \end{bmatrix}$$

Let $\lambda_1, \lambda_2, \lambda_3$ be the eigen values of A.

$$\text{Given } \lambda_1 = 2, \lambda_2 = 3$$

$\lambda_1 + \lambda_2 + \lambda_3 =$ sum of the main diagonal elements.

$$2 + 3 + \lambda_3 = 2 + 2 + 2$$

$$5 + \lambda_3 = 6$$

$$\boxed{\lambda_3 = 1}$$

$$\text{W.K.T } \lambda_1, \lambda_2, \lambda_3 = |A|$$

$$2 \times 3 \times 1 = \begin{vmatrix} 2 & 0 & 1 \\ 0 & 2 & 0 \\ a & 0 & 2 \end{vmatrix}$$

$$\Rightarrow 6 = 2(4 - 0) - 0(0) + 1(-2a)$$

$$6 = 8 - 2a$$

$$2a = 2 \Rightarrow \boxed{a = 1}$$

Problem-16:-

Find the eigen values of a matrix

$$\begin{bmatrix} 7 & -2 & 0 \\ -2 & 6 & -2 \\ 0 & -2 & 5 \end{bmatrix}$$

corresponding to the eigen vector $[-4, -2, 4]^T$.

Soln:-

$$\text{W.K.T } (A - \lambda I)X = 0 \quad [A/M-2017]$$

$$\begin{bmatrix} 7-\lambda & -2 & 0 \\ -2 & 6-\lambda & -2 \\ 0 & -2 & 5-\lambda \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$(7-\lambda)(2) - 2 + 0 = 0$$

$$14 - 2\lambda - 2 = 0$$

$$12 - 2\lambda = 0$$

$$2\lambda = 12$$

$$\boxed{\lambda = 6}$$

Hence the corresponding eigen value is 6.

Problem-17:-

State Cayley Hamilton theorem

Soln:-

Every square matrix satisfies its own characteristic equation.

Problem-18:-

Use Cayley Hamilton theorem to find $A^4 - 4A^3 - 5A^2 + A + 2I$

$$\text{When } A = \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix}$$

Soln:-

$$\text{Given } A = \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix}$$

The char. eqn of A is

$$\lambda^2 - S_1\lambda + S_2 = 0$$

$$S_1 = 1 + 3 = 4$$

$$S_2 = \begin{vmatrix} 1 & 2 \\ 4 & 3 \end{vmatrix} = 3 - 8 = -5$$

$\therefore$ The char. eqn of A is

$$\lambda^2 - 4\lambda - 5 = 0$$

$$\lambda_1 + \lambda_2 + \lambda_3 = \lambda_1 + \lambda_2$$

$$\therefore \boxed{\lambda_3 = 0}$$

W.K.T

Product of the Eigen values
= $|A|$

$$\lambda_1 \lambda_2 \lambda_3 = \lambda_1 \lambda_2 (0) \\ = 0$$

Problem : 9

Find the sum and product of all the Eigen values of the matrix

$$\begin{bmatrix} 1 & 2 & -2 \\ 1 & 0 & 3 \\ -2 & -1 & -3 \end{bmatrix}$$

Soln:-

Sum of the Eigen values

= sum of the main diagonal elements.

$$= 1 + 0 - 3 = -2$$

Product of the Eigen values

$$= \begin{vmatrix} 1 & 2 & -2 \\ 1 & 0 & 3 \\ -2 & -1 & -3 \end{vmatrix}$$

$$= 1(0+3) - 2(-3+6) - 2(-1+0)$$

$$= 3 - 6 + 2$$

$$= -1.$$

Problem 10:-

If 3 and 6 are two Eigen values of $A = \begin{bmatrix} 1 & 1 & 3 \\ 1 & 5 & 1 \\ 3 & 1 & 1 \end{bmatrix}$

write down all the eigen values of A^T .

Soln:-

Let $\lambda_1, \lambda_2, \lambda_3$ be the Eigen values of A .

Given $\lambda_1 = 3, \lambda_2 = 6, \lambda_3 = ?$

W.K.T

Sum of the Eigen values
= sum of the main diagonal elements.

$$\lambda_1 + \lambda_2 + \lambda_3 = 1 + 5 + 1$$

$$3 + 6 + \lambda_3 = 7$$

$$\therefore \boxed{\lambda_3 = -2}$$

W.K.T

Eigen values of A^T are

$$\frac{1}{\lambda_1}, \frac{1}{\lambda_2}, \frac{1}{\lambda_3}$$

$\therefore \frac{1}{3}, \frac{1}{6}, -\frac{1}{2}$ are the eigen values of A^T .

Problem 11:-

Find the Eigen values of A^T

$$\text{where } A = \begin{bmatrix} 3 & 1 & 4 \\ 0 & 2 & 6 \\ 0 & 0 & 5 \end{bmatrix}$$

Soln:-

A is a upper triangular matrix. Hence the Eigen values of A are 3, 2, 5

$\therefore \frac{1}{3}, \frac{1}{2}, \frac{1}{5}$ are the Eigen values of A^T

Problem 12:-

Find the Eigen values of A^3 given $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 2 & -7 \\ 0 & 0 & 3 \end{bmatrix}$

Soln:-

$$\text{Given } A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 2 & -7 \\ 0 & 0 & 3 \end{bmatrix}$$

clearly given A is an upper triangular matrix.

Hence the Eigen values are 1, 2, 3

W.K.T "If λ is eigen value of A , then λ^n is an Eigen value of A^n "

$\therefore 1^2, 2^2, 3^2$ are the eigen values of A^2

$\Rightarrow 1, 8, 27$ are the eigen values of A^3 .

Problem-13:-

If 2, -1, -3 are the eigen values of the matrix A then find the Eigen values of the matrix $A^2 - 2I$.

Soln:-

Given the Eigen values of A are 2, -1, -3.

The Eigen values of $A^2 - 2I$ are $2^2 - 2(1)$, $(-1)^2 - 2(1)$, $(-3)^2 - 2(1)$

$\Rightarrow 2, -1, 7$ are the eigen values of $A^2 - 2I$.

Problem: 14:-

Find the constants a and b such that matrix $\begin{bmatrix} a & 4 \\ 1 & b \end{bmatrix}$ has 3 and -2 as its eigen values.

Soln:-

Apr-2017

Sum of the eigen values of a matrix = sum of the main diagonal elements.

$$3 + (-2) = a + b$$

$$\Rightarrow a + b = 1 \rightarrow \textcircled{1}$$

Product of the eigen values = $|A|$

$$(3) \times (-2) = \begin{vmatrix} a & 4 \\ 1 & b \end{vmatrix}$$

$$-6 = ab - 4$$

$$\Rightarrow ab = -2 \rightarrow \textcircled{2}$$

Solving $\textcircled{1}$ & $\textcircled{2}$

$$x^2 - (a+b)x + ab = 0$$

$$\Rightarrow x^2 - x - 2 = 0$$

$$(x-2)(x+1) = 0$$

$$\text{i.e., } x-2=0, \quad x+1=0$$

$$x=2, \quad x=-1$$

$$\therefore a=2, \quad b=-1 \quad (\text{or}) \quad a=-1, \quad b=2.$$

Problem-15:-

If 2 and 3 are the eigen values of $\begin{bmatrix} 2 & 0 & 1 \\ 0 & 2 & 0 \\ a & 0 & 2 \end{bmatrix}$,

find the value of 'a'.

PART-AP-1

If 3 and 15 are two eigen values of the matrix $A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$ then find its third Eigen values and hence find $|A|$.

Sol

Given $\lambda_1 = 3, \lambda_2 = 15$

W.K.T sum of the Eigen values = sum of the main diagonal elements

$$\therefore \lambda_1 + \lambda_2 + \lambda_3 = 8 + 7 + 3$$

$$3 + 15 + \lambda_3 = 18$$

$$\boxed{\lambda_3 = 0}$$

Also W.K.T

Product of the Eigen value = $|A|$

$$3 \times 15 \times 0 = |A|$$

$$\therefore \boxed{|A| = 0}$$

Problem 2:-

Show that the Eigen values of a null matrix are zero.

[APR-2018]

Soln:-

If A is a null square matrix of order 3 then its char. eqn is $(-1)^3 \lambda^3 = 0$

$$\lambda^3 = 0$$

$\therefore$ The Eigen values are 0, 0, 0.
In general,

The char eqn of a null square matrix of order n is

$$(-1)^n \lambda^n = 0$$

$$\therefore \lambda^n = 0$$

$\therefore$ The Eigen values of A are zero

Problem-3:-

If λ is the Eigen value of the matrix A , then Prove that λ^2 is the eigen value of A^2 (APR-2019)

Soln:-

Given: λ is the eigen value of the matrix A .

$$AX = \lambda X \quad \text{--- (1)}$$

$$A(AX) = A(\lambda X)$$

$$A^2 X = \lambda(AX)$$

$$= \lambda(\lambda X) \quad [\text{using (1)}]$$

$$= \lambda^2 X$$

$\therefore \lambda^2$ is an eigen value of A^2 .

Problem 4:-

If the Eigen values of the matrix A of order 3×3 are matrix 2, 3 and 1, then find the det of A .

Soln:-

Given Eigen values of A are 2, 3, 1

$|A|$ = Product of the eigen values.
 $= 2 \times 3 \times 1$

$$\boxed{|A| = 6}$$

Problem 5:-

Prove that any square matrix A and its transpose have the same Eigen values.

Soln:-

NID-2019

Let A be a square matrix of order n .

The char eqn of A and A^T are

$$|A - \lambda I| = 0 \rightarrow \textcircled{1}$$

$$\&|A^T - \lambda I| = 0 \rightarrow \textcircled{2}$$

$$\text{W.K.T } |A| = |A^T|$$

Hence $\textcircled{1}$ & $\textcircled{2}$ are identical.

$\therefore$ The Eigen values of A & A^T are same.

Problem-6:-

If $A = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$ then find

$2A^2 - 8A - 10I$, where I is the unit matrix.

Soln:-

$$\text{Given } A = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$$

The char eqn of A is $|A - \lambda I| = 0$

$$\text{i.e., } \lambda^2 - S_1\lambda + S_2 = 0$$

$$S_1 = 4$$

$$S_2 = \begin{vmatrix} 1 & 4 \\ 2 & 3 \end{vmatrix} = 3 - 8 = -5$$

$$\therefore \lambda^2 - 4\lambda + 5 = 0$$

By C-H Theorem,

$$\text{We have } A^2 - 4A - 5I = 0 \rightarrow \textcircled{1}$$

$$\therefore 2A^2 - 8A - 10I = 2[A^2 - 4A - 5I] = 0$$

Problem 7.

The Product of the two Eigen values of the matrix

$$A = \begin{bmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{bmatrix} \text{ is } 16.$$

Find the third Eigen value.

Soln:-

Let $\lambda_1, \lambda_2, \lambda_3$ are the Eigen values of A .

$$\text{Given } \lambda_1 \lambda_2 = 16$$

$$\text{W.K.T } \lambda_1 \lambda_2 \lambda_3 = |A|$$

$$\lambda_1 \lambda_2 \lambda_3 = \begin{vmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{vmatrix}$$

$$16 \lambda_3 = 6(9-1) + 2(-6+2) + 2(2-6)$$

$$16 \lambda_3 = 32$$

$$\boxed{\lambda_3 = 2}$$

Problem 8:-

If the Sum of two Eigen values and trace of a 3×3 matrix A are equal find the value of $|A|$

Soln:-

Let $\lambda_1, \lambda_2, \lambda_3$ be the Eigen values of the given 3×3 matrix A .

W.K.T

Sum of the Eigen values = Trace of A . $\textcircled{1}$

Trace of A = Sum of the two Eigen value

Problem 21:-

Find the index, signature of the Q.F. $x_1^2 + 2x_2^2 - 3x_3^2$.

Soln:-

Here Q.F = C.F

$$f(x_1, x_2, x_3) = x_1^2 + 2x_2^2 - 3x_3^2$$

Index = No. of +ive terms in the canonical form
= 2

Signature = No. of positive terms in the canonical form - No. of negative terms in the canonical form
= 2 - 1 = 1

Problem 22:-

Write down the quadratic form corresponding to the matrix $\begin{bmatrix} 0 & 5 & -1 \\ 5 & 1 & 6 \\ -1 & 6 & 2 \end{bmatrix}$

Soln

$$\text{Let } \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} 0 & 5 & -1 \\ 5 & 1 & 6 \\ -1 & 6 & 2 \end{bmatrix}$$

$\therefore$ Q.F is

$$\begin{aligned} & a_{11}x_1^2 + a_{22}x_2^2 + a_{33}x_3^2 \\ & + 2a_{12}x_1x_2 + 2a_{23}x_2x_3 \\ & + 2a_{13}x_1x_3 \\ & = x_2^2 + 2x_3^2 + 10x_1x_2 + 12x_2x_3 - 2x_1x_3. \end{aligned}$$

Problem 23:-

Write down the Q.F corresponding to the matrix $\begin{bmatrix} 2 & 0 & -2 \\ 0 & 2 & 1 \\ -2 & 1 & -2 \end{bmatrix}$

Soln:-

$$\text{Let } \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} 2 & 0 & -2 \\ 0 & 2 & 1 \\ -2 & 1 & -2 \end{bmatrix}$$

$$\begin{aligned} \text{Q.F is } & a_{11}x_1^2 + a_{22}x_2^2 + a_{33}x_3^2 \\ & + 2a_{12}x_1x_2 + 2a_{13}x_1x_3 + 2a_{23}x_2x_3 \\ & = 2x_1^2 + 2x_2^2 - 2x_3^2 - 4x_1x_3 + 2x_2x_3 \end{aligned}$$

Problem 24:-

Write down nature of a quadratic form whose matrix is $\begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -2 \end{bmatrix}$.

Soln:-

Eigen values of the given matrix are -1, -1, -2

All the eigen values are negative numbers.

$\therefore$ The nature of the Q.F is negative definite.

Problem 25:-

Can $A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ be diagonalized? why

Soln:-

The given matrix A is real symmetric and non-singular matrix.

Hence

A can be diagonalised.

UNIT - II - DIFFERENTIAL CALCULUS

Representation of functions - Limit of a function - continuity - Derivatives - Differentiation rules (sum, product, quotient, chain rules) - Implicit differentiation - Logarithmic differentiation - Applications: Maxima and minima of functions of one variable.

REPRESENTATION OF FUNCTIONS:

A function is a rule that assigns to each element 'x' in a set 'A' to exactly one element called 'f(x)' in a set 'B'.

* Domain: Let $f: A \rightarrow B$ then set 'A' is called the domain of the function.

* Co-domain: Set 'B' is called co-domain of the function.

* Range: The set of all images of all the elements of 'A' under the function 'f' is called the range of 'f' and it is denoted by $f(A)$.

PROBLEMS

1. Find the domain of the function $f(x) = \sqrt{x+2}$

Soln:

Since the square root of a negative number is not defined, the domain of 'f' must be positive.

$$\therefore x+2 \geq 0$$

$$\Rightarrow x \geq -2$$

$$\therefore \text{Domain is } [-2, \infty)$$

$$\begin{array}{c} x \geq -2 \\ \hline -2 \qquad \qquad \qquad \infty \end{array}$$

2. Find the domain of the function $f(x) = \sqrt{3-x} - \sqrt{x+2}$

Soln:

NOV/DEC 2018

$$\text{Let, } 3-x \geq 0 \text{ and } x+2 \geq 0$$

$$\Rightarrow x \leq 3 \text{ and } x \geq -2$$

$$\therefore \text{Domain is } [-2, 3]$$

$$\begin{array}{c} \hline -2 \qquad \qquad \qquad 3 \\ x \geq -2 \rightarrow \qquad \leftarrow x \leq 3 \end{array}$$

[Since the square root of a negative number is not defined, the domain of 'f' must be positive]

3. Find the domain of the function $f(x) = \frac{x+4}{x^2-9}$

Soln:

Given: $f(x) = \frac{x+4}{x^2-9}$

The function is not defined at $x=3$ and $x=-3$

Domain: $\{x / x \neq 3, x \neq -3\}$

Domain: $(-\infty, -3) \cup (-3, 3) \cup (3, \infty)$

4. Find the domain, range also sketch the graph for the following functions, $f(x) = \begin{cases} x+2, & x < 0 \\ 1-x, & x \geq 0 \end{cases}$


Soln:

Given: $f(x) = \begin{cases} x+2, & x < 0 \\ 1-x, & x \geq 0 \end{cases}$

Domain: x	-1	-2	-3	...
Range: $y = x+2$	1	0	-1	...

Domain: x	0	1	2	...
Range: $y = 1-x$	1	0	-1	...

Domain: $(-\infty, \infty)$
Range: $(-\infty, 1)$




5. Sketch the graph of the function $f(x) = \begin{cases} 1+x, & x < -1 \\ x^2, & -1 \leq x \leq 1 \\ 2-x, & x \geq 1 \end{cases}$ and use it to determine the values of 'a' for which $\lim_{x \rightarrow a} f(x)$ exists?

Soln:

Given: $f(x) = \begin{cases} 1+x, & x < -1 \\ x^2, & -1 \leq x \leq 1 \\ 2-x, & x \geq 1 \end{cases}$

From the graph, it is observed that $\lim_{x \rightarrow a} f(x)$ exists for all 'a' except

when $a = -1$, since the right and left limits are different at $a = -1$.



LIMIT OF A FUNCTION:

Let $f(x)$ be a function of a real variable ' x '. Let ' a ' and ' l ' be a fixed numbers. If ' $f(x)$ ' approaches ' l ' as ' x ' approaches ' a ', then we say ' l ' is the limit of $f(x)$ as ' x ' tends to ' a ' & we write $\lim_{x \rightarrow a} f(x) = l$.

LEFT HAND LIMIT:

If $f(x)$ approaches the value ' l ' as ' x ' approaches ' a ' from the left, then $\lim_{x \rightarrow a^-} f(x) = l$.

RIGHT HAND LIMIT:

If $f(x)$ approaches the value ' l ' as ' x ' approaches ' a ' from the right, then $\lim_{x \rightarrow a^+} f(x) = l$.

Result:

$$\lim_{x \rightarrow a} f(x) = l \text{ if and only if } \lim_{x \rightarrow a^-} f(x) = l = \lim_{x \rightarrow a^+} f(x).$$

Problems

1. Evaluate $\lim_{x \rightarrow -6} \frac{2x+12}{|x+6|}$

Soln:

We know that $\lim_{x \rightarrow a} f(x) = l$ iff $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = l$

Given: $\lim_{x \rightarrow -6} \frac{2x+12}{|x+6|}$


$$\text{Now, } f(x) = \begin{cases} \frac{2(x+6)}{x+6}, & x+6 \geq 0 \\ \frac{2(x+6)}{-(x+6)}, & x+6 < 0 \end{cases}$$

$$= \begin{cases} 2, & x \geq -6 \\ -2, & x < -6 \end{cases}$$

$$\therefore \lim_{x \rightarrow -6^-} \frac{2x+12}{|x+6|} = -2 \quad \text{--- (1)}$$

$$\& \lim_{x \rightarrow -6^+} \frac{2x+12}{|x+6|} = 2 \quad \text{--- (2)}$$

① $\neq$ ② $\therefore$ Limit does not exist.



2. Check whether $\lim_{x \rightarrow -3} \frac{3x+9}{|x+3|}$ exists.

April / May 2019

Soln:


Given: $\lim_{x \rightarrow -3} \frac{3x+9}{|x+3|}$, $f(x) = \begin{cases} \frac{3(x+3)}{x+3}, & x+3 > 0 \\ \frac{3(x+3)}{-(x+3)}, & x+3 < 0 \end{cases}$

$$= \begin{cases} 3, & x > -3 \\ -3, & x < -3 \end{cases}$$

Now, $\lim_{x \rightarrow -3^+} \frac{3x+9}{|x+3|} = 3$ — (1)

Also, $\lim_{x \rightarrow -3^-} \frac{3x+9}{|x+3|} = -3$ — (2)

(1) $\neq$ (2) $\therefore$ limit does not exist.



3. Guess the value of the limit (if it exists) for the function $\lim_{x \rightarrow 0} \frac{e^{5x} - 1}{x}$ by evaluating the function at the given numbers $x = \pm 0.5, \pm 0.1, \pm 0.01, \pm 0.001, \pm 0.0001$ [correct to six decimal places]

NOV/DEC 2018

Soln:

Let $f(x) = \frac{e^{5x} - 1}{x}$

x	-0.5	-0.1	-0.01	-0.001	-0.0001
f(x)	1.835830	3.934693	4.877058	4.987521	4.998750

x	0.5	0.1	0.01	0.001	0.0001
f(x)	22.364988	6.487213	5.127110	5.012521	5.001250

As 'x' approaches to '0', the function $f(x) = \frac{e^{5x} - 1}{x}$ approaches to '5'.

$$\therefore \lim_{x \rightarrow 0} \frac{e^{5x} - 1}{x} = 5.$$

4. Investigate $\lim_{n \rightarrow 0} \frac{1}{n^2}$

Soln:

Let $f(n) = \frac{1}{n^2}$

n	-1	-0.5	-0.1	-0.05	-0.01	-0.001	0.001	0.01	0.05	0.1	0.5	1
$f(n)$	1	4	100	400	10000	1000000	1000000	10000	400	100	4	1

As 'n' approaches to '0', the function $f(n) = \frac{1}{n^2}$ becomes very large and does not approach to a number.

$$\therefore \lim_{n \rightarrow 0} \frac{1}{n^2} = \infty.$$

5. Evaluate $\lim_{n \rightarrow 0} \frac{\sin n}{n}$

Soln:

Given: $\lim_{n \rightarrow 0} \frac{\sin n}{n}$, Here $f(n) = \frac{\sin n}{n}$

n	-1	-0.5	-0.1	-0.05	-0.01	-0.001	0.001	0.01	0.05
$f(n)$	0.8415	0.9589	0.9983	0.9996	0.9999	0.99999	0.99999	0.9999	0.9996

0.1	0.5	1
0.9983	0.9589	0.8415

As 'n' approaches to '0', the function $f(n) = \frac{\sin n}{n}$ approaches to 1.

$$\therefore \lim_{n \rightarrow 0} \frac{\sin n}{n} = 1.$$

6. Prove that $\lim_{n \rightarrow 0} |n| = 0$

Proof:

Let $f(n) = |n|$

$$= \begin{cases} n & \text{if } n \geq 0 \\ -n & \text{if } n < 0 \end{cases}$$

We know that $\lim_{n \rightarrow a} f(n) = L$ iff $\lim_{n \rightarrow a^-} f(n) = L = \lim_{n \rightarrow a^+} f(n)$

Now, $\lim_{n \rightarrow 0^-} |n| = \lim_{n \rightarrow 0^-} (-n) = 0$

Also, $\lim_{n \rightarrow 0^+} |n| = \lim_{n \rightarrow 0^+} n = 0 \therefore \lim_{n \rightarrow 0} |n| = 0$

7. Determine $\lim_{x \rightarrow \frac{\pi}{2}} \frac{1 + \cos 2x}{(\pi - 2x)^2}$

AU JAN 2016 R-15 MAT151

Soln:

Let, $\lim_{x \rightarrow \frac{\pi}{2}} \frac{1 + \cos 2x}{(\pi - 2x)^2} = \frac{1 + \cos \frac{\pi}{2}}{(\pi - 2 \cdot \frac{\pi}{2})^2} = \frac{1-1}{0} = \frac{0}{0}$

Apply L'Hospital rule,

① $\Rightarrow \lim_{x \rightarrow \frac{\pi}{2}} \frac{-2 \sin 2x}{2(\pi - 2x)(-2)} = \lim_{x \rightarrow \frac{\pi}{2}} \frac{\sin 2x}{2(\pi - 2x)}$
 $= \frac{\sin \frac{\pi}{2}}{2(\pi - 2 \cdot \frac{\pi}{2})} = \frac{0}{0}$

Apply L'Hospital rule,

② $\Rightarrow \lim_{x \rightarrow \frac{\pi}{2}} \frac{2 \cos 2x}{2(-2)} = \frac{\cos \frac{\pi}{2}}{-2} = \frac{-1}{-2} = \frac{1}{2}$

8. Evaluate $\lim_{x \rightarrow \infty} [x\sqrt{x^2+1} - x]$

Soln:

Let, $\lim_{x \rightarrow \infty} [x\sqrt{x^2+1} - x] = \lim_{x \rightarrow \infty} x\sqrt{x^2+1} - x \left[\frac{\sqrt{x^2+1} + x}{\sqrt{x^2+1} + x} \right]$
 $= \lim_{x \rightarrow \infty} \frac{x[x^2+1 - x^2]}{\sqrt{x^2+1} + x}$
 $= \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2(1+\frac{1}{x^2})} + x}$
 $= \lim_{x \rightarrow \infty} \frac{x}{x\sqrt{1+\frac{1}{x^2}} + x}$
 $= \lim_{x \rightarrow \infty} \frac{x}{x[\sqrt{1+\frac{1}{x^2}} + 1]}$
 $= \frac{1}{\sqrt{1+\frac{1}{\infty}} + 1} = \frac{1}{2}$

9. Evaluate $\lim_{n \rightarrow \infty} \frac{3n^2 - n - 2}{5n^2 + 4n + 1}$

Soln:

$$\text{Let } f(n) = \frac{3n^2 - n - 2}{5n^2 + 4n + 1}$$

$$\begin{aligned} \therefore \lim_{n \rightarrow \infty} f(n) &= \lim_{n \rightarrow \infty} \frac{3n^2 - n - 2}{5n^2 + 4n + 1} \\ &= \lim_{n \rightarrow \infty} \frac{n^2 \left(3 - \frac{1}{n} - \frac{2}{n^2} \right)}{n^2 \left(5 + \frac{4}{n} + \frac{1}{n^2} \right)} \\ &= \frac{3 - \frac{1}{\infty} - \frac{2}{\infty}}{5 + \frac{4}{\infty} + \frac{1}{\infty}} = \frac{3}{5} \end{aligned}$$

HORIZONTAL ASYMPTOTE:

The line $y=L$ is called a horizontal asymptote of the curve $y=f(x)$ if either $\lim_{x \rightarrow \infty} f(x) = L$ (or) $\lim_{x \rightarrow -\infty} f(x) = L$.

PROBLEMS

1. Find the horizontal asymptote of the curve $\frac{x^2-1}{x^2+1}$

Soln:

$$\text{Given: } f(x) = \frac{x^2-1}{x^2+1}$$

$$\begin{aligned} \lim_{x \rightarrow -\infty} f(x) &= \lim_{x \rightarrow -\infty} \frac{x^2-1}{x^2+1} \\ &= \lim_{x \rightarrow -\infty} \frac{x^2 \left(1 - \frac{1}{x^2} \right)}{x^2 \left(1 + \frac{1}{x^2} \right)} \\ &= \frac{1-0}{1+0} = 1. \end{aligned}$$

$$\text{and } \lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} \frac{x^2-1}{x^2+1} = \lim_{x \rightarrow \infty} \frac{x^2 \left(1 - \frac{1}{x^2} \right)}{x^2 \left(1 + \frac{1}{x^2} \right)}$$

Hence the line $y=1$ is a horizontal asymptote of the given curve.

$$= \frac{1-0}{1+0} = 1.$$

Q. Find the horizontal and vertical asymptotes of the curve $\frac{\sqrt{2x^2+1}}{3x-5}$

Soln:

$$\text{Let } f(x) = \frac{\sqrt{2x^2+1}}{3x-5}$$

$$\begin{aligned} \lim_{x \rightarrow \infty} f(x) &= \lim_{x \rightarrow \infty} \frac{\sqrt{2x^2+1}}{3x-5} \\ &= \lim_{x \rightarrow \infty} \frac{\sqrt{x^2 \left(2 + \frac{1}{x^2}\right)}}{x \left(3 - \frac{5}{x}\right)} \\ &= \lim_{x \rightarrow \infty} \frac{x \sqrt{2 + \frac{1}{x^2}}}{x \left[3 - \frac{5}{x}\right]} \\ &= \frac{\sqrt{2+0}}{3} \\ &= \frac{\sqrt{2}}{3} \end{aligned}$$

$$\begin{aligned} \lim_{x \rightarrow -\infty} f(x) &= \lim_{x \rightarrow -\infty} \frac{\sqrt{2x^2+1}}{3x-5} \quad \text{when } x \rightarrow \infty \\ &= \lim_{x \rightarrow -\infty} \frac{-x \sqrt{2 + \frac{1}{x^2}}}{x \left(3 - \frac{5}{x}\right)} \quad \text{we have } \sqrt{x^2} = -x, x < 0 \\ &= \frac{-\sqrt{2}}{3} \end{aligned}$$

$\therefore$ Both the line $y = \frac{-\sqrt{2}}{3}$ and $\frac{\sqrt{2}}{3}$ are horizontal asymptotes. The vertical asymptotes occurs when the given function becomes either $-\infty$ or ∞ .

For $x = \frac{5}{3}$ the function becomes ∞ .

$$\therefore \lim_{x \rightarrow \left(\frac{5}{3}\right)^+} f(x) = \lim_{x \rightarrow \frac{5}{3}^+} \frac{\sqrt{2x^2+1}}{3x-5} = \infty$$

$$\text{and } \lim_{x \rightarrow \frac{5}{3}^-} f(x) = \lim_{x \rightarrow \frac{5}{3}^-} \frac{\sqrt{2x^2+1}}{3x-5} = -\infty.$$

$$-\sqrt{x^2 + x^2} \leq \sqrt{x^2 + x^2} \sin \frac{\pi}{n} \leq \sqrt{x^2 + x^2}$$

$$\lim_{x \rightarrow 0} -\sqrt{x^2 + x^2} \leq \lim_{x \rightarrow 0} \sqrt{x^2 + x^2} \sin \frac{\pi}{n} \leq \lim_{x \rightarrow 0} \sqrt{x^2 + x^2}$$

$$\text{since } \lim_{x \rightarrow 0} -\sqrt{x^2 + x^2} = 0 \text{ \& } \lim_{x \rightarrow 0} \sqrt{x^2 + x^2} = 0.$$

$$\therefore \text{By squeeze theorem, } \lim_{x \rightarrow 0} \sqrt{x^2 + x^2} \sin \frac{\pi}{n} = 0.$$

3. Show that $\lim_{x \rightarrow 0} x^2 \cos\left(\frac{1}{x^2}\right)$.

PROOF:

$$\text{Let, } -1 \leq \cos \frac{1}{x^2} \leq 1$$

$$-x^2 \leq x^2 \cos \frac{1}{x^2} \leq x^2$$

$$\lim_{x \rightarrow 0} -(x^2) \leq \lim_{x \rightarrow 0} x^2 \cos \frac{1}{x^2} \leq \lim_{x \rightarrow 0} x^2$$

$$\text{since } \lim_{x \rightarrow 0} (-x^2) = 0 \text{ \& } \lim_{x \rightarrow 0} (x^2) = 0$$

$$\text{By squeeze theorem, } \lim_{x \rightarrow 0} x^2 \cos\left(\frac{1}{x^2}\right) = 0.$$

CONTINUITY

A function 'f' is continuous at the point 'a' if $\lim_{x \rightarrow a} f(x) = f(a)$.

* A function 'f' is continuous from right at a point 'a' if $\lim_{x \rightarrow a^+} f(x) = f(a)$.

* A function 'f' is continuous from left at a point 'a' if $\lim_{x \rightarrow a^-} f(x) = f(a)$.

RESULT: f(x) is continuous if $\lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow a^-} f(x) = f(a)$.

PROBLEMS

1. Show that the function $f(x) = 1 - \sqrt{1 - x^2}$ is continuous on the interval $[-1, 1]$.

Soln:

$$\text{Let, } f(x) = 1 - \sqrt{1-x^2}$$

$$\begin{aligned}\lim_{x \rightarrow -1^+} f(x) &= \lim_{x \rightarrow -1^+} 1 - \sqrt{1-x^2} \\ &= 1 - \sqrt{1-(-1)^2} \\ &= 1\end{aligned}$$

$$\begin{aligned}\lim_{x \rightarrow 1^-} f(x) &= \lim_{x \rightarrow 1^-} 1 - \sqrt{1-x^2} \\ &= 1 - \sqrt{1-1} \\ &= 1\end{aligned}$$

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} 1 - \sqrt{1-x^2}$$

$$= 1 - \sqrt{1-a^2}$$

$$= f(a)$$

$$\therefore -1 < a < 1$$

f is continuous.

$$\therefore \lim_{x \rightarrow -1^+} f(x) = \lim_{x \rightarrow 1^-} f(x) = 1.$$

$\therefore f(x)$ is continuous on the interval $[-1, 1]$.

Q. For what value of the constant 'c' is the function 'f' continuous on $(-\infty, \infty)$, $f(x) = \begin{cases} cx^2 + 2x; & x < 2 \\ x^3 - cx; & x \geq 2 \end{cases}$

Soln:

The given function $f(x)$ is

continuous on $(-\infty, 2)$ and $(2, \infty)$. Now,

$$\begin{aligned}\lim_{x \rightarrow 2^-} f(x) &= \lim_{x \rightarrow 2^-} (cx^2 + 2x) \\ &= c(4) + 2(2) \\ &= 4c + 4\end{aligned}$$

$$\begin{array}{ccc} cx^2 + 2x & & x^3 - cx \\ \downarrow & & \downarrow \\ -\infty & 2 & \infty \end{array}$$

$$\begin{aligned}\text{Also, } \lim_{x \rightarrow 2^+} f(x) &= \lim_{x \rightarrow 2^+} (x^3 - cx) \\ &= 8 - 2c\end{aligned}$$

We know that a function 'f' is continuous at a point 'a' if

$$\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x)$$

$$\therefore 4c + 4 = 8 - 2c$$

$$4c + 2c = 4$$

$$c = 4/6$$

$$\boxed{c = 2/3}$$


APRIL/MAY 2019
JAN 2018.

3. Let $f(x) = \begin{cases} 2x-2, & x < -1 \\ ax+b, & -1 \leq x \leq 1 \\ 5x+7, & x \geq 1 \end{cases}$ is continuous for all

real 'x', Find the values of 'a' & 'b'.

Soln:

Given: $f(x)$ is continuous.



$$\begin{aligned} \therefore \lim_{x \rightarrow -1^-} f(x) &= \lim_{x \rightarrow -1^-} 2x-2 \\ &= -2-2 = -4 \end{aligned}$$

$$\begin{aligned} \& \lim_{x \rightarrow -1^+} f(x) &= \lim_{x \rightarrow -1^+} ax+b \\ &= a(-1)+b = -a+b \end{aligned}$$

$$\begin{aligned} \because f \text{ is continuous, } \lim_{x \rightarrow -1^-} f(x) &= \lim_{x \rightarrow -1^+} f(x) \\ -4 &= -a+b \\ \Rightarrow a-b &= 4 \quad \text{--- (1)} \end{aligned}$$

$$\begin{aligned} \text{Also, } \lim_{x \rightarrow 1^-} f(x) &= \lim_{x \rightarrow 1^-} ax+b \\ &= a(1)+b = a+b \end{aligned}$$

$$\begin{aligned} \& \lim_{x \rightarrow 1^+} f(x) &= \lim_{x \rightarrow 1^+} 5x+7 \\ &= 5+7 = 12 \end{aligned}$$

$$\begin{aligned} \because f \text{ is continuous, } \lim_{x \rightarrow 1^-} f(x) &= \lim_{x \rightarrow 1^+} f(x) \\ a+b &= 12 \quad \text{--- (2)} \end{aligned}$$

From (1) & (2)


$$\begin{array}{r} a+b = 12 \\ a-b = 4 \quad (+) \\ \hline 2a = 16 \\ \boxed{a = 8} \end{array}$$

$$\begin{aligned} \therefore (1) \Rightarrow 8-b &= 4 \\ \boxed{b} &= 4 \end{aligned}$$

$$\begin{aligned} \therefore a &= 8 \\ \& b &= 4. \end{aligned}$$

4. Find the domain where the function 'f' is continuous. Also find the number at which the function 'f' is discontinuous where $f(x) = \begin{cases} 1+x^2, & x \leq 0 \\ 2-x, & 0 < x \leq 2 \\ (x-2)^2, & x > 2 \end{cases}$

Soln:



At $x=0$,

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (1+x^2) = 1 \quad \text{--- (1)}$$

$$\& \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (2-x) = 2 \quad \text{--- (2)}$$

(1) $\neq$ (2) $\therefore f(x)$ is discontinuous at $x=0$.

At $x=2$,

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (2-x) = 0 \quad \text{--- (3)}$$


$$\& \lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (x-2)^2 = 0 \quad \text{--- (4)}$$

(3) = (4) $\therefore f(x)$ is continuous at $x=2$.

$\therefore$ Domain: $(-\infty, 0) \cup (0, \infty)$.

5. Find the values of 'a' and 'b' that make 'f' continuous on $(-\infty, \infty)$, $f(x) = \begin{cases} \frac{x^2-8}{x-2} & \text{if } x < 2 \\ ax^2-bx+3 & \text{if } 2 \leq x \leq 3 \\ 2x-a+b & \text{if } x \geq 3 \end{cases}$

Soln:



NOV/DEC 2018

At $x=2$,

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} \frac{x^2-8}{x-2} = \frac{2^2-8}{2-2} = \frac{-4}{0} = \infty$$

Apply L'Hospital's Rule,

$$= \lim_{x \rightarrow 2^-} \frac{2x}{1} = 2(2) = 4$$

Also, $\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} ax^2 - bx + 3$
 $= 4a - 2b + 3$

Given: f is continuous.

$$\therefore \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x)$$

$$\Rightarrow 4a - 2b + 3 = 4$$

$$4a - 2b = 1 \quad \text{--- (1)}$$

At $x=3$,

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} ax^2 - bx + 3$$

$$= 9a - 3b + 3 \quad \text{--- (2)}$$

Also, $\lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3^+} 2x - a + b$
 $= 6 - a + b \quad \text{--- (3)}$

Given: f is continuous,

$$\therefore \lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^+} f(x)$$

$$9a - 3b + 3 = 6 - a + b \quad \text{By (2) \& (3)}$$

$$10a - 4b = 3 \quad \text{--- (4)}$$

From (1) \& (4) $10a - 4b = 3$

$$(-) \quad 8a - 4b = 2 \quad [(1) \times 2]$$

$$\hline 2a = 1$$

$$\boxed{a = 1/2}$$

$$\therefore (1) \Rightarrow 4 \cdot \frac{1}{2} - 2b = 1$$

$$2 - 1 = 2b$$

$$\boxed{b = 1/2}$$

DERIVATIVE :

The derivative of a function 'f' at a number 'a', denoted by $f'(a)$, is

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} \text{ if this limit exists.}$$

PROBLEMS

1. If $f(x) = \sqrt{x}$, find the derivative of $f(x)$.

Soln:

By the definition of derivative,

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h} \times \frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \\ &= \lim_{h \rightarrow 0} \frac{x+h-x}{h(\sqrt{x+h} + \sqrt{x})} \\ &= \lim_{h \rightarrow 0} \frac{1}{\sqrt{x+h} + \sqrt{x}} = \frac{1}{\sqrt{x} + \sqrt{x}} = \frac{1}{2\sqrt{x}} \end{aligned}$$

2. If $f(x) = \sin x$, find the derivative of $f(x)$.

Soln:

$$\begin{aligned} \text{Let, } f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h} \\ &= \lim_{h \rightarrow 0} \frac{2 \cos\left(\frac{x+h}{2}\right) \sin \frac{h}{2}}{h} \quad \begin{array}{l} \because \sin A - \sin B \\ = 2 \cos\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right) \end{array} \\ &= \lim_{h \rightarrow 0} \frac{\sin \frac{h}{2}}{\frac{h}{2}} \cos\left(x + \frac{h}{2}\right) \\ &= \lim_{h \rightarrow 0} \frac{\sin h/2}{h/2} \times \lim_{h \rightarrow 0} \cos\left(x + \frac{h}{2}\right) \\ &= 1 \times \cos x \quad \begin{array}{l} \because \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \end{array} \\ &= \cos x \end{aligned}$$

3. Find the derivative of the function $f(x) = \frac{1}{\sqrt{x}}$.

Soln:

$$\begin{aligned}
 \text{Let, } f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{1}{\sqrt{x+h}} - \frac{1}{\sqrt{x}}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{\sqrt{x} - \sqrt{x+h}}{\sqrt{x} \sqrt{x+h} \cdot h}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sqrt{x} - \sqrt{x+h}}{\sqrt{x} \sqrt{x+h} \cdot h} \times \frac{\sqrt{x} + \sqrt{x+h}}{\sqrt{x} + \sqrt{x+h}} \\
 &= \lim_{h \rightarrow 0} \frac{x - (x+h)}{\sqrt{x} \sqrt{x+h} h [\sqrt{x} + \sqrt{x+h}]} \\
 &= \lim_{h \rightarrow 0} \frac{-h}{\cancel{h} \sqrt{x} \sqrt{x+h} [\sqrt{x} + \sqrt{x+h}]} \\
 &= \lim_{h \rightarrow 0} \frac{-1}{\sqrt{x} \sqrt{x+h} [\sqrt{x} + \sqrt{x+h}]} \\
 &= \frac{-1}{\sqrt{x} \sqrt{x} (\sqrt{x} + \sqrt{x})} = \frac{-1}{2x\sqrt{x}}
 \end{aligned}$$

RULES OF DIFFERENTIATION:

1. $\frac{d}{dx}(c) = 0$ 2. $\frac{d}{dx}(c \cdot u) = c \frac{du}{dx}$, [u is a function of x & c is a constant]

3. Product rule, $\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$

4. Quotient rule, $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{vu' - uv'}{v^2}$

5. Chain rule,

(i) If 'y' is a function of 'u' and 'u' itself is a function of x, then $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

(ii) If 'y' is a function of 'u', 'u' is a function of 'v', 'v' is a function of 'w' and 'w' is a function of 'x', then $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dw} \cdot \frac{dw}{dx}$

PROBLEMS

1. If $f(x) = e^x (x + x\sqrt{x})$, find the derivative of $f(x)$.

Soln:

$$\text{Given: } f(x) = e^x (x + x\sqrt{x})$$

$$= e^x (x + x^{3/2})$$

$$f'(x) = \frac{d}{dx} [e^x (x + x^{3/2})]$$

$$= e^x \frac{d}{dx} (x + x^{3/2}) + (x + x^{3/2}) \frac{d}{dx} e^x$$

$$= e^x \left(1 + \frac{3}{2} x^{\frac{3}{2}-1}\right) + (x + x^{3/2}) e^x$$

$$= e^x \left[1 + \frac{3}{2} x^{1/2} + x + x^{3/2}\right]$$

$$= e^x \left[1 + x + \frac{3\sqrt{x}}{2} + x\sqrt{x}\right]$$

2. Find $\frac{dy}{dx}$ if $y = x^2 e^{2x} (x^2 + 1)^4$.

APRIL/MAY 2019

Soln:

$$\text{Given: } y = x^2 e^{2x} (x^2 + 1)^4$$

$$\frac{dy}{dx} = \frac{d}{dx} [x^2 e^{2x} (x^2 + 1)^4]$$

$$= x^2 e^{2x} \frac{d}{dx} (x^2 + 1)^4 + x^2 (x^2 + 1)^4 \frac{d}{dx} e^{2x} + e^{2x} (x^2 + 1)^4 \frac{d}{dx} (x^2)$$

$$= x^2 e^{2x} 4(x^2 + 1)^3 (2x) + x^2 (x^2 + 1)^4 (2e^{2x}) + e^{2x} (x^2 + 1)^4 (2x)$$

$$= 8x^3 e^{2x} (x^2 + 1)^3 + 2x^2 e^{2x} (x^2 + 1)^4 + 2x e^{2x} (x^2 + 1)^4$$

$$= 2x e^{2x} (x^2 + 1)^3 [4x^2 + x(x^2 + 1) + x^2 + 1]$$

$$= 2x e^{2x} (x^2 + 1)^3 [4x^2 + x^3 + x + x^2 + 1]$$

$$= 2x e^{2x} (x^2 + 1)^3 (x^3 + 5x^2 + x + 1)$$

3. If $f(x) = \frac{1 - xe^x}{x + e^x}$, find the derivative of $f(x)$.

Soln:

$$\text{Let, } f'(x) = \frac{d}{dx} \left[\frac{1 - xe^x}{x + e^x} \right]$$

$$= \frac{(x + e^x) \frac{d}{dx} (1 - xe^x) - (1 - xe^x) \frac{d}{dx} (x + e^x)}{(x + e^x)^2}$$

$$\begin{aligned}
 &= \frac{(x+e^x)(-e^x - xe^x) - (1-xe^x)(1+e^x)}{(x+e^x)^2} \\
 &= \frac{-xe^x - x^2e^x - e^{2x} - xe^{2x} - 1 - e^x + xe^x + xe^{2x}}{(x+e^x)^2} \\
 &= \frac{-x^2e^x - e^{2x} - 1 - e^x}{(x+e^x)^2} \\
 &= \frac{-(x^2e^x + e^{2x} + e^x + 1)}{(x+e^x)^2}
 \end{aligned}$$

4. If $y = (1-x^2)^{10}$, find the derivative of y .

Soln:

Given: $y = (1-x^2)^{10}$

Let $u = 1-x^2 \therefore y = u^{10}$

$$\frac{du}{dx} = -2x \quad \frac{dy}{du} = 10u^9$$

$$\begin{aligned}
 \therefore \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\
 &= 10u^9 (-2x) = -20x(1-x^2)^9
 \end{aligned}$$

5. If $y = \tan(\sin x)$, find the derivative of y .

Soln:

Let $u = \sin x \therefore y = \tan u$

$$\frac{dy}{du} = \sec^2 u = \sec^2(\sin x)$$

$$\begin{aligned}
 \therefore \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\
 &= \sec^2(\sin x) \cos x
 \end{aligned}$$

6. If $y = \log(x + \sqrt{x^2-1})$, find the derivative of y .

Soln:

Let $y = \log u$ where $u = x + \sqrt{x^2-1}$

$$\begin{aligned}
 \frac{dy}{du} &= \frac{1}{u} \quad \frac{du}{dx} = 1 + \frac{1}{\sqrt{x^2-1}} \quad (\neq x) \\
 &= 1 + \frac{x}{\sqrt{x^2-1}}
 \end{aligned}$$

$$\begin{aligned}
 \therefore \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\
 &= \frac{1}{u} \left(1 + \frac{x}{\sqrt{x^2-1}} \right) \\
 &= \frac{1}{x+\sqrt{x^2-1}} \left(1 + \frac{x}{\sqrt{x^2-1}} \right) \\
 &= \frac{1}{x+\sqrt{x^2-1}} \cdot \frac{x+\sqrt{x^2-1}}{\sqrt{x^2-1}} \\
 &= \frac{1}{\sqrt{x^2-1}}
 \end{aligned}$$

7. If $y = \sin(\sin(\sin x))$, find the derivative of 'y'.

Soln:

$$\text{Let } y = \sin u, \quad u = \sin v, \quad v = \sin x$$

$$\frac{dy}{du} = \cos u, \quad \frac{du}{dv} = \cos v, \quad \frac{dv}{dx} = \cos x$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx}$$

$$= \cos u \cos v \cos x$$

$$= \cos(\sin v) \cos(\sin x) \cos x$$

$$= \cos(\sin(\sin x)) \cos(\sin x) \cos x.$$

8. If $y = \cos \sqrt{\sin(\tan \pi x)}$, find the derivative of 'y'.

Soln:

$$\text{Let } y = \cos \sqrt{u}, \quad u = \sin v, \quad v = \tan \pi x$$

$$\frac{dy}{du} = -\sin \sqrt{u} \left(\frac{1}{2\sqrt{u}} \right), \quad \frac{du}{dv} = \cos v, \quad \frac{dv}{dx} = \sec^2 \pi x (\pi)$$

$$\therefore \frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx}$$

$$= \frac{-1}{2\sqrt{u}} \sin \sqrt{u} \times \cos v \times \pi \sec^2 \pi x$$

$$= \frac{-1}{2\sqrt{\sin v}} \sin \sqrt{\sin v} \times \cos(\tan \pi x) \pi \sec^2 \pi x$$

$$= \frac{-\pi}{2} \frac{\sin \sqrt{\sin(\tan \pi x)}}{\sqrt{\sin(\tan \pi x)}} \times \cos(\tan \pi x) \sec^2 \pi x.$$

IMPLICIT DIFFERENTIATION:PROBLEMS

1. If $\sqrt{x} + \sqrt{y} = 1$ then find $\frac{dy}{dx}$

Soln:

Given: $\sqrt{x} + \sqrt{y} = 1$

Differentiate with respect to 'x'

$$\frac{1}{2\sqrt{x}} + \frac{1}{2\sqrt{y}} \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = \frac{-2\sqrt{y}}{2\sqrt{x}} = -\frac{\sqrt{y}}{\sqrt{x}}$$

2. Find y'' if $x^4 + y^4 = 16$

Soln:

Given $x^4 + y^4 = 16$

JAN 2018

Differentiate w.r. to 'x'.

$$4x^3 + 4y^3 \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = -\frac{x^3}{y^3} = y'$$

$$\therefore y'' = \frac{d^2y}{dx^2} = - \frac{y^3 (3x^2) - (x^3)(3y^2) \frac{dy}{dx}}{y^6}$$

$$= - \frac{3x^2y^3 - 3x^3y^2 \left(-\frac{x^3}{y^3} \right)}{y^6}$$

$$= - \frac{3x^2y^3 + \frac{3x^6}{y}}{y^6}$$

$$= - \frac{3x^2y^4 + 3x^6}{y^7}$$

$$= - \frac{3x^2}{y^7} [y^4 + x^4] = \frac{-3x^2}{y^7} (16) = \frac{-48x^2}{y^7}$$

NOV/DEC 2018

3. Find y' if $\cos(ny) = 1 + \sin y$

Soln:

Given $\cos ny = 1 + \sin y$

Diff. w.r. to 'x'.

$$-\sin ny \left[n \frac{dy}{dx} + y \cdot n \right] = \cos y \frac{dy}{dx}$$

$$-\sin ny \quad \checkmark \quad n \frac{dy}{dx} - y \sin ny - \cos y \quad \checkmark \quad \frac{dy}{dx} = 0$$

$$-\frac{dy}{dx} [n \sin ny + \cos y] = y \sin ny$$

$$\therefore \frac{dy}{dx} = \frac{-y \sin ny}{n \sin ny + \cos y}$$

4. If $x\sqrt{1+y} + y\sqrt{1+x} = 0$, then prove that $\frac{dy}{dx} = \frac{-1}{(1+x)^2}$.

Soln:

$$\text{Let } x\sqrt{1+y} = -y\sqrt{1+x}$$

Squaring on both sides,

$$x^2(1+y) = y^2(1+x)$$

$$x^2 + x^2y = y^2 + y^2x$$

$$(x^2 - y^2) + x^2y - y^2x = 0$$

$$(x+y)(x-y) + xy(x-y) = 0$$

$$x-y [x+y+xy] = 0$$

$$\text{If } x-y=0, \text{ then } 1 - \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = 1.$$

$$\text{III}^{\text{rd}} \text{ If } x+y+xy = 0$$

$$y+xy = -x$$

$$y(1+x) = -x$$

$$y = \frac{-x}{1+x}$$

$$\frac{dy}{dx} = - \frac{(1+x) \cdot 1 - x \cdot 1}{(1+x)^2}$$

$$= - \frac{1+x-x}{(1+x)^2} = \frac{-1}{(1+x)^2}.$$

DERIVATIVE OF TRIGONOMETRIC FUNCTIONS:

FORMULA'S

$$1. \frac{d}{dx} (\sin x) = \cos x$$

$$2. \frac{d}{dx} (\cos x) = -\sin x$$

$$3. \frac{d}{dx} (\tan x) = \sec^2 x$$

$$4. \frac{d}{dx} (\cot x) = -\operatorname{cosec}^2 x$$

$$5. \frac{d}{dx} (\sec x) = \sec x \tan x$$

$$6. \frac{d}{dx} (\operatorname{cosec} x) = -\operatorname{cosec} x \cot x$$

$$1. \frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$$

$$2. \frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$$

$$3. \frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}$$

$$4. \frac{d}{dx} (\cot^{-1} x) = \frac{-1}{1+x^2}$$

$$5. \frac{d}{dx} (\sec^{-1} x) = \frac{1}{x\sqrt{x^2-1}}$$

$$6. \frac{d}{dx} (\operatorname{cosec}^{-1} x) = \frac{-1}{x\sqrt{x^2-1}}$$

1. If $f(x) = \cos^{-1} \left[\frac{b+a \cos x}{a+b \cos x} \right]$, find the derivative of $f(x)$.

NOV/DEC 2018

Soln:

$$\text{Let } u = \frac{b+a \cos x}{a+b \cos x}$$

$$\therefore \text{Let } y = f(x) = \cos^{-1}(u)$$

$$\frac{dy}{dx} = \frac{-1}{\sqrt{1-u^2}} \frac{du}{dx}$$

$$= \frac{-1}{\sqrt{1-u^2}} \left[\frac{(a+b \cos x)(-a \sin x) - (b+a \cos x)(-b \sin x)}{(a+b \cos x)^2} \right]$$

$$= \frac{-1}{\sqrt{1-\left(\frac{b+a \cos x}{a+b \cos x}\right)^2}} \left[\frac{-a^2 \sin x - ab \sin x \cos x + b^2 \sin x + ab \sin x \cos x}{(a+b \cos x)^2} \right]$$

$$= \frac{-(a+b \cos x)}{\sqrt{1-\frac{b^2+a^2 \cos^2 x+2ab \cos x - b^2 - a^2 \cos^2 x - 2ab \cos x}{(a+b \cos x)^2}}} \frac{-(a^2-b^2) \sin x}{(a+b \cos x)^2}$$

$$\begin{aligned}
 &= \frac{(a^2 - b^2) \sin n}{\sqrt{a^2(1 - \cos^2 n) - b^2(1 - \cos^2 n)} (a + b \cos n)} \\
 &= \frac{(a^2 - b^2) \sin n}{\sqrt{a^2 \sin^2 n - b^2 \sin^2 n} (a + b \cos n)} \\
 &= \frac{(a^2 - b^2) \sin n}{\sin n \sqrt{a^2 - b^2} (a + b \cos n)} \\
 &= \frac{\sqrt{a^2 - b^2} \sin n}{a + b \cos n}
 \end{aligned}$$

Q. Find $\frac{dy}{dx}$ if $y = \tan^{-1} \left[\frac{\sqrt{1+x^2} - 1}{x} \right]$

Soln:

Let $x = \tan \theta$ ——— (1)

$$\begin{aligned}
 \therefore y &= \tan^{-1} \left[\frac{\sqrt{1 + \tan^2 \theta} - 1}{\tan \theta} \right] \\
 &= \tan^{-1} \left(\frac{\sec \theta - 1}{\tan \theta} \right) \\
 &= \tan^{-1} \left(\frac{1/\cos \theta - 1}{\sin \theta / \cos \theta} \right) \\
 &= \tan^{-1} \left(\frac{1 - \cos \theta}{\sin \theta} \right) \\
 &= \tan^{-1} \left(\frac{2 \sin^2(\theta/2)}{2 \sin(\theta/2) \cos(\theta/2)} \right) \\
 &= \tan^{-1} (\tan(\theta/2))
 \end{aligned}$$

$y = \theta/2$

$y = \frac{\tan^{-1} x}{2}$ By (1)

$\therefore \frac{dy}{dx} = \frac{1}{2} \left(\frac{1}{1+x^2} \right)$

DERIVATIVE OF LOGARITHMIC FUNCTIONS:-PROBLEMS1. Differentiate $y = n^n$

Soln:

Given $y = n^n$

Taking 'log' on both sides,

$$\log y = \log n^n$$

$$\log y = n \log n$$

Differentiate with respect to 'n'.

$$\begin{aligned} \frac{1}{y} \frac{dy}{dn} &= n \frac{1}{n} + \log n \quad (1) \\ &= 1 + \log n \end{aligned}$$

$$\begin{aligned} \frac{dy}{dn} &= y (1 + \log n) \\ &= n^n (1 + \log n) \end{aligned}$$

2. Find y' if $n^y = y^n$

Soln:

Given $y^n = n^y$

Taking 'log' on both sides,

$$\log n^y = \log y^n$$

$$y \log n = n \log y$$

Differentiate with respect to 'n'.

$$y \frac{1}{n} + \log n \frac{dy}{dn} = n \frac{1}{y} \frac{dy}{dn} + \log y \quad (1)$$

$$\frac{dy}{dn} \left(\log n - \frac{n}{y} \right) = \log y - \frac{y}{n}$$

$$\frac{dy}{dn} = \frac{n \log y - y}{n} \times \frac{y}{y \log n - n}$$

$$\frac{dy}{dn} = \frac{y (n \log y - y)}{n (y \log n - n)}$$

3. Differentiate $y = n^{\cos n}$

Soln:

Given: $y = n^{\cos n}$

Taking 'log' on both sides

$$\log y = \log (n^{\cos n})$$

$$\log y = \cos n \log n$$

Differentiate w.r. to 'n'

$$\frac{1}{y} \frac{dy}{dn} = \cos n \left(\frac{1}{n} \right) + \log n (-\sin n)$$

$$\begin{aligned} \frac{dy}{dn} &= y \left(\frac{\cos n - n \sin n \log n}{n} \right) \\ &= n^{\cos n} \left(\frac{\cos n - n \sin n \log n}{n} \right) \end{aligned}$$

TANGENT & NORMAL:

* The equation of tangent at a given point (x_1, y_1) is given by $y - y_1 = m(x - x_1)$.

* The equation of normal at a given point (x_1, y_1) is given by $y - y_1 = -\frac{1}{m}(x - x_1)$.

PROBLEMS

1. Find the points on the curve $y = x^4 - 6x^2 + 4$ where the tangent line is horizontal?

Soln:

Given: $y = x^4 - 6x^2 + 4$

Differentiate w.r. to 'n',

$$\frac{dy}{dn} = 4x^3 - 12x$$

The tangent line is horizontal if $y' = 0$

$$\therefore y' = 4x^3 - 12x = 0$$

$$\Rightarrow 4x(x^2 - 3) = 0$$

$$\begin{aligned} 4x &= 0, & x^2 - 3 &= 0 \\ \boxed{x=0} &, & \boxed{x = \pm\sqrt{3}} \end{aligned}$$

$\therefore$ So the given curve has horizontal tangents when $x=0, \sqrt{3}$ & $-\sqrt{3}$.

$$\begin{aligned} \text{when } x=0 &\Rightarrow y = 0 - 6(0) + 4 \Rightarrow \boxed{y=4} \\ &\Rightarrow (0, 4) \end{aligned}$$

$$\begin{aligned} \text{when } x=\sqrt{3} &\Rightarrow (\sqrt{3})^4 - 6(\sqrt{3})^2 + 4 = 4 \\ &\Rightarrow \boxed{y=-5} \Rightarrow (\sqrt{3}, -5) \end{aligned}$$

When $x = -\sqrt{3} \Rightarrow y = (-\sqrt{3})^4 - 6(-\sqrt{3})^2 + 4$

$$\boxed{y = -5} \Rightarrow (-\sqrt{3}, -5).$$

$\therefore$ The corresponding points are $(0, 4)$, $(\sqrt{3}, -5)$ & $(-\sqrt{3}, -5)$.

2. Find the equation of the tangent line to the curve $x^3 + y^3 = 6xy$ at the point $(3, 3)$ and what point the tangent line horizontal in the first quadrant.

JAN 2018

Soln:

Given: $x^3 + y^3 = 6xy$, diff. w.r.to 'x',

$$3x^2 + 3y^2 \frac{dy}{dx} = 6 \left[x \frac{dy}{dx} + y \right]$$

$$x^2 + y^2 \frac{dy}{dx} = 2x \frac{dy}{dx} + 2y$$

$$\frac{dy}{dx} = \frac{2y - x^2}{y^2 - 2x}, \text{ at } (3, 3) \frac{dy}{dx} = \frac{6-9}{9-6} = -1.$$

$$\therefore \boxed{m = -1} \quad \text{--- (1)}$$

The equation of the tangent at the point $(3, 3)$ is

$$y - 3 = -1(x - 3) \Rightarrow y - 3 = -x + 3$$

$$\boxed{x + y = 6}$$

The tangent line is horizontal if $y' = 0$

$$\text{(1)} \Rightarrow \frac{2y - x^2}{y^2 - 2x} = 0 \Rightarrow 2y - x^2 = 0$$

$$\boxed{y = x^2/2} \quad \text{--- (2)}$$

Let $x^3 + y^3 = 6xy$

$$x^3 + \left(\frac{x^2}{2}\right)^3 = 6x \frac{x^2}{2} \text{ by (2)}$$

$$x^3 + \frac{x^6}{8} = \frac{6x^3}{2} \Rightarrow x^3 + \frac{x^6}{8} = 3x^3$$

$$\Rightarrow 2x^3 = \frac{x^6}{8}$$

$$x^6 = 16x^3$$

$$\boxed{x^3 = 16}$$

$$\therefore x = 16^{1/3}$$

$$\Rightarrow \boxed{x = 2^{4/3}}$$

$$\therefore \text{(2)} \Rightarrow y = \frac{(2^{4/3})^2}{2} = \frac{2^{8/3}}{2} = 2^{5/3}$$

$\therefore$ The tangent is horizontal at $(2^{4/3}, 2^{5/3})$.

Applications: Maxima and minima of functions of one variable.

Let 'c' be a point in a domain 'D' of the function 'f'.

Then $f(c)$ is the * absolute maximum value of 'f' on 'D' if

$$f(c) \geq f(x), \text{ for all 'x' in 'D'.$$

* absolute minimum value of 'f' on 'D' if

$$f(c) \leq f(x) \text{ for all 'x' in 'D'.$$

Absolute Maximum and Minimum of $f(x)$:

Step (i): Find the critical numbers of $f(x)$.

$$\text{i.e. } \boxed{f'(x) = 0}$$

Step (ii): Substitute the critical points at $f(x)$.

Step (iii): Then the maximum value of $f(x)$ is absolute maximum and the minimum value of $f(x)$ is absolute minimum.

1. Find the absolute maximum and absolute minimum value of $f(x) = x - 2\sin x$ on $[0, 2\pi]$.

Soln:

$$\text{Given: } f(x) = x - 2\sin x$$

$$\therefore f'(x) = 1 - 2\cos x$$

$$\text{Critical numbers: } f'(x) = 0$$

$$\Rightarrow 1 - 2\cos x = 0$$

$$\cos x = 1/2$$

$$x = \frac{\pi}{2}, \frac{5\pi}{2}$$

$$\therefore f\left(\frac{\pi}{2}\right) = \frac{\pi}{2} - 2\sin\left(\frac{\pi}{2}\right) = -0.68485 \text{ [radian mode]}$$

$$f\left(\frac{5\pi}{2}\right) = \frac{5\pi}{2} - 2\sin\left(\frac{5\pi}{2}\right) = 6.968039$$

At end

$$\text{points, } f(0) = 0 - 2\sin 0 = 0$$

$$f(2\pi) = 2\pi - 2\sin 2\pi = 2\pi = 6.28.$$

$\therefore$ The absolute minimum value is $f\left(\frac{\pi}{2}\right) = -0.68485$

The absolute maximum value is $f\left(\frac{5\pi}{2}\right) = 6.9680.$

Q. Find the absolute maximum and absolute minimum value of $f(x) = 3x^4 - 4x^3 - 12x^2 + 11$ on $[-2, 3]$.

Soln:

$$\text{Given: } f(x) = 3x^4 - 4x^3 - 12x^2 + 11$$

$$f'(x) = 12x^3 - 12x^2 - 24x$$

The critical numbers of $f(x)$ are occurred at

$$\begin{aligned} f'(x) = 0 &\Rightarrow 12x^3 - 12x^2 - 24x = 0 \\ 12x(x^2 - x - 2) &= 0 \end{aligned}$$

$$\begin{array}{cc} -2 & -1 \\ -2 \times 1 & -2 \times 1 \end{array}$$

$$x = 0, x = 2, x = -1$$

The values of $f(x)$ at these critical numbers are

$$f(0) = 0 + 1 = 1$$

$$f(2) = 3(16) - 4(8) - 12(4) + 11 = -4$$

$$f(-1) = 3(1) - 4(-1) - 12(1) + 11 = -31$$

At end points,

$$f(-2) = 33$$

$$f(3) = 28$$

$\therefore$ The absolute minimum value is $f(-1) = -31$

The absolute maximum value is $f(-2) = 33$.

FIRST AND SECOND DERIVATIVE TEST:

INCREASING / DECREASING / CONCAVITY / INFLECTION POINTS :-

INCREASING & DECREASING FUNCTIONS:-

* If $f'(x) > 0$ in an interval (a, b) , then 'f' is increasing $\uparrow$.

* If $f'(x) < 0$ in an interval (a, b) , then 'f' is decreasing $\downarrow$.

CRITICAL NUMBER:

A critical number of a function 'f' is a number 'c' in the domain of 'f' such that $f'(c) = 0$ (or) $f'(c)$ does not exist.

FIRST DERIVATIVE TEST:

Suppose that 'c' is a critical number of a continuous function 'f' then

* If $f'(x)$ changes from positive to negative at 'c', then $f(x)$ has local maximum at 'c'.

* If $f'(x)$ changes from negative to positive at 'c', then $f(x)$ has local minimum at 'c'.

* If $f'(x)$ does not change sign at c , then $f(x)$ has no local maximum or minimum at c .

Note: The first derivative test is a consequence of the increasing and decreasing test.

CONCAVITY: CONCAVE UPWARDS / CONCAVE DOWNWARDS

* If $f''(x) > 0$ in any interval, then $f(x)$ is concave upwards [convex downwards].

* If $f''(x) < 0$ in any interval, then $f(x)$ is concave downwards [convex upwards].

INFLECTION POINTS:

A point 'P' on a curve $y = f(x)$ is called an inflection point if $f(x)$ is continuous and the curve changes from concave upwards to concave downwards or from concave downwards to concave upwards at 'P'.

SECOND DERIVATIVE TEST:

Suppose $f''(x)$ is continuous near c ,

* If $f'(c) = 0$ and $f''(c) > 0$, then $f(x)$ has a local minimum at c .

* If $f'(c) = 0$ and $f''(c) < 0$, then $f(x)$ has a local maximum at c .

PROBLEMS

1. The profit function of aersonic company is given by $P(x) = -0.02x^2 + 300x - 200,000$, dollars, where the function 'P' is increasing and where it is decreasing.

Soln:

The derivative P' of the function P is


$$P'(x) = -0.04x + 300$$

$$P'(x) = -0.04(x - 7500)$$

To find critical points: $P'(x) = 0$

$$-0.04(x - 7500) = 0$$


$$\boxed{x = 7500}$$



* $P'(x) > 0$ for 'x' in the interval $(0, 7500)$.

* $P'(x) < 0$ for 'x' in the interval $(7500, \infty)$.

This means that the profit function 'P' is increasing on $(0, 7500)$ and decreasing on $(7500, \infty)$.



Q. For the function $f(x) = 2 + 2x^2 - x^4$, find the intervals of increasing (or) decreasing, local maximum (or) minimum values and the intervals of concavity also the inflection points.

Soln:

Given: $f(x) = 2 + 2x^2 - x^4$

NOV/DEC 2018

$$f'(x) = 4x - 4x^3$$


To find the critical points:-

$$f'(x) = 0 \Rightarrow 4x - 4x^3 = 0$$

$$4x(1 - x^2) = 0$$

$$x = 0, 1 = x^2$$

$$\boxed{x = 0}, \boxed{x = \pm 1}$$



$\therefore$ The critical points are -1, 0, 1.

Interval	Sign of $f'(x)$	Behaviour of $f(x)$
$(-\infty, -1)$	$f'(-2) = 4(-2) - 4(-2)^3$ $= 24$ (ive)	Increasing.
$(-1, 0)$	$f'(-1/2) = 4(-1/2) - 4(-1/2)^3$ $= -3/2$ (ive)	Decreasing
$(0, 1)$	$f'(1/2) = 4(1/2) - 4(1/2)^3$ $= 3/2$, (ive)	Increasing
$(1, \infty)$	$f'(2) = 4(2) - 4(2)^3$ $= -24$ (ive)	Decreasing

Local maximum (or) minimum using First derivative Test:

From the above table,

* $f'(x)$ changes from +ive to -ive at $x = -1$.

$\therefore f(x)$ has a local maximum at $x = -1$.

$\therefore$ Local maximum value at $x = -1$, $f(-1) = 2 + 2(-1)^2 - (-1)^4$
 $= 3$.

* $f'(x)$ changes from +ive to +ive at $x = 0$.

$\therefore f(x)$ has a local minimum at $x = 0$.

$\therefore$ Local minimum value at $x = 0$, $f(0) = 2 + 2(0) - 0$
 $= 2$.

* $f'(x)$ changes from +ive to -ive at $x = 1$.

$\therefore f(x)$ has a local maximum at $x = 1$.

$\therefore$ Local maximum value at $x = 1$, $f(1) = 2 + 2(1) - 1$
 $= 3$.

Concavity:

$$f''(x) = 4 - 12x^2$$

$$f''(x) = 0 \Rightarrow 4 - 12x^2 = 0$$

$$4 = 12x^2$$

$$x^2 = \frac{1}{3} \Rightarrow x = \pm \frac{1}{\sqrt{3}}$$

$$= \pm 0.58$$



Interval	Sign of $f''(x)$	Behaviour of $f(x)$
$(-\infty, -0.58)$	$f(-1) = 4 - 12(-1)^2 = -8$, (-ive)	Concave downwards
$(-0.58, 0.58)$	$f(0) = 4 - 12(0) = 4$, (+ive)	Concave upwards
$(0.58, \infty)$	$f(1) = 4 - 12(1)^2 = -8$, (-ive)	Concave downwards

Local maximum (or) minimum using second derivative Test:

* $f'(-1) = 0$ and $f''(-1) = 4 - 12(-1)^2 = -8 < 0$ (From the above table)
 $\therefore f(x)$ is Local maximum at $x = -1$.

* $f'(0) = 0$ and $f''(0) = 4 - 12(0) = 4 > 0$
 $\therefore f(x)$ is Local minimum at $x = 0$.

$$* f'(1) = 0, \quad f''(1) = 4 - 12(1)^2 = -8 < 0$$

$\therefore f(x)$ is Local maximum at $x=1$.

Inflection points:

* Curve changes from concave downward to concave upward at $x = -0.58$.

$$\therefore f(-0.58) = 2 + 2(-0.58) - (-0.58)^4 \\ = 2.56$$

$\therefore$ Inflection pts are $(-0.58, 2.56)$.

* Also, curve changes from concave upward to concave downward at $x = 0.58$.

$$\therefore f(0.58) = 2 + (0.58) - (0.58)^4 \\ = 2.56$$

$\therefore$ Inflection pts are $(0.58, 2.56)$.

2. Find the Local maximum (or) minimum values of $f(x) = \sqrt{x} - 4\sqrt[4]{x}$ using both first and second derivative tests.

Soln:

$$\text{Given: } f(x) = \sqrt{x} - 4\sqrt[4]{x} \\ = x^{1/2} - x^{1/4}$$

$$f'(x) = \frac{1}{2}x^{-1/2} - \frac{1}{4}x^{-3/4} \\ = \frac{x^{-3/4}}{4} (2x^{1/4} - 1) \\ = \frac{2x^{1/4} - 1}{4x^{3/4}}$$

To find critical points:


$$f'(x) = 0 \Rightarrow \frac{2x^{1/4} - 1}{4x^{3/4}} = 0$$

$$\Rightarrow 2x^{1/4} - 1 = 0$$

$$\Rightarrow 2x^{1/4} = 1$$

$$x^{1/4} = 1/2$$

$$\Rightarrow x = (1/2)^4 = 1/16$$



Second Derivative Test:

$$\textcircled{1} \Rightarrow c''(n) = \frac{240}{n^3}$$

$$\Rightarrow c''(\sqrt{60}) > 0.$$

Hence 'c' is a maximum at $n = \sqrt{60}$

$$\therefore y = \frac{30}{n} = \frac{30}{\sqrt{60}} \text{ m.}$$

6. If $f(x) = 2x^3 + 3x^2 - 36x$, find the intervals on which it is increasing (or) decreasing, the local maximum (or) minimum values of $f(x)$.

Soln:

Given $f(x) = 2x^3 + 3x^2 - 36x$

April/May 2019

$$f'(x) = 6x^2 + 6x - 36$$

$$\text{Let } f'(x) = 0 \Rightarrow x^2 + x - 6 = 0$$

$$\begin{array}{cc} -6 & 1 \\ -2 \times 3 & -2 + 3 \end{array}$$

$$\boxed{x = -3, -2}$$

Increasing / decreasing:

Interval	sign of $f'(x)$	Behaviour of
$(-\infty, -3)$	$f'(-4) = 6(-4)^2 + 6(-4) - 36 = 36$ (Positive)	Increasing
$(-3, -2)$	$f'(-1) = 6(-1)^2 + 6(-1) - 36 = -36$ (Negative)	Decreasing
$(-2, \infty)$	$f'(3) = 6(3)^2 + 6(3) - 36 = 36 > 0$	Increasing

First Derivative Test: [Local maximum/minimum]

* $f'(x)$ changes positive to negative at $x = -3$

$\therefore f(x)$ has a local maximum at $x = -3$.

$$\therefore \boxed{f(-3) = 81}$$

* $f'(x)$ changes negative to positive at $x = -2$

$\therefore f(x)$ has a local minimum at $x = -2$.

$$\& \boxed{f(-2) = -44}$$

Second Derivative Test: $f''(x) = 12x + 6$

* $f'(-3) = 0$ and $f''(-3) = -30 < 0$

$\therefore f(x)$ is local maximum at $x = -3$.

* $f'(-2) = 0$ & $f''(-2) = 30 > 0$ $\therefore f(x)$ is local minimum at $x = -2$

$$\begin{aligned}\frac{\partial z}{\partial u} - \frac{\partial z}{\partial v} &= e^u \frac{\partial z}{\partial x} - e^{-u} \frac{\partial z}{\partial y} + e^{-v} \frac{\partial z}{\partial x} + e^v \frac{\partial z}{\partial y} \\ &= (e^u + e^{-v}) \frac{\partial z}{\partial x} - (e^{-u} - e^v) \frac{\partial z}{\partial y} = x \frac{\partial z}{\partial x} - y \frac{\partial z}{\partial y}\end{aligned}$$

⑦ Transform the equation $zx_x + 2zxy + zyy = 0$ by changing the independent variables using $u = x - y$ and $v = x + y$.

Soln:-

(AU June 202)

$u = x - y$	$v = x + y$
$\frac{\partial u}{\partial x} = 1$	$\frac{\partial v}{\partial x} = 1$
$\frac{\partial u}{\partial y} = -1$	$\frac{\partial v}{\partial y} = 1$
$\begin{aligned}\frac{\partial z}{\partial x} &= \frac{\partial z}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial z}{\partial v} \frac{\partial v}{\partial x} \\ &= \frac{\partial z}{\partial u} (1) + \frac{\partial z}{\partial v} (1) \\ &= \frac{\partial z}{\partial u} + \frac{\partial z}{\partial v}\end{aligned}$	$\begin{aligned}\frac{\partial z}{\partial y} &= \frac{\partial z}{\partial u} \frac{\partial u}{\partial y} + \frac{\partial z}{\partial v} \frac{\partial v}{\partial y} \\ &= \frac{\partial z}{\partial u} (-1) + \frac{\partial z}{\partial v} (1) \\ &= -\frac{\partial z}{\partial u} + \frac{\partial z}{\partial v}\end{aligned}$
$\frac{\partial}{\partial x} = \frac{\partial}{\partial u} + \frac{\partial}{\partial v}$	$\frac{\partial}{\partial y} = -\frac{\partial}{\partial u} + \frac{\partial}{\partial v}$
$\begin{aligned}\frac{\partial^2 z}{\partial x^2} &= \left(\frac{\partial}{\partial x}\right) \left(\frac{\partial z}{\partial x}\right) \\ &= \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v}\right) \left(\frac{\partial z}{\partial u} + \frac{\partial z}{\partial v}\right) \\ &= \frac{\partial^2 z}{\partial u^2} + \frac{\partial^2 z}{\partial u \partial v} + \frac{\partial^2 z}{\partial v \partial u} + \frac{\partial^2 z}{\partial v^2}\end{aligned}$	$\begin{aligned}\frac{\partial^2 z}{\partial y^2} &= \left(\frac{\partial}{\partial y}\right) \left(\frac{\partial z}{\partial y}\right) \\ &= \left(-\frac{\partial}{\partial u} + \frac{\partial}{\partial v}\right) \left(-\frac{\partial z}{\partial u} + \frac{\partial z}{\partial v}\right) \\ &= \frac{\partial^2 z}{\partial u^2} - \frac{\partial^2 z}{\partial u \partial v} - \frac{\partial^2 z}{\partial v \partial u} + \frac{\partial^2 z}{\partial v^2}\end{aligned}$

$$\begin{aligned}\frac{\partial^2 z}{\partial x \partial y} &= \left(\frac{\partial}{\partial x}\right) \left(\frac{\partial z}{\partial y}\right) = \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v}\right) \left(-\frac{\partial z}{\partial u} + \frac{\partial z}{\partial v}\right) \\ &= -\frac{\partial^2 z}{\partial u^2} + \frac{\partial^2 z}{\partial u \partial v} - \frac{\partial^2 z}{\partial v \partial u} + \frac{\partial^2 z}{\partial v^2}\end{aligned}$$


$$\begin{aligned}zx_x + 2zxy + zyy &= 0 \\ \Rightarrow \frac{\partial^2 z}{\partial u^2} + \frac{\partial^2 z}{\partial u \partial v} + \frac{\partial^2 z}{\partial v \partial u} + \frac{\partial^2 z}{\partial v^2} - 2\frac{\partial^2 z}{\partial u^2} + 2\frac{\partial^2 z}{\partial u \partial v} - 2\frac{\partial^2 z}{\partial v \partial u} + 2\frac{\partial^2 z}{\partial v^2} \\ &+ \frac{\partial^2 z}{\partial u^2} - \frac{\partial^2 z}{\partial u \partial v} - \frac{\partial^2 z}{\partial v \partial u} + \frac{\partial^2 z}{\partial v^2} = 0\end{aligned}$$

$$\begin{aligned}4\frac{\partial^2 z}{\partial v^2} + 2\frac{\partial^2 z}{\partial u \partial v} - 2\frac{\partial^2 z}{\partial v \partial u} &= 0 \Rightarrow 4Z_{vv} + 2Z_{uv} - Z_{vu} = 0 \\ \Rightarrow 2Z_{vv} + Z_{uv} - Z_{vu} &= 0\end{aligned}$$




Introduction

In Mathematics, integral assign numbers to functions in a way that can describe displacement, area, volume and other concepts that arise by combining infinitesimal data. Integral calculus plays a vital role in Mathematics, Engineering, Science and Economics.

First let us concentrate to solve the area problem. Given a function f which is continuous and non-negative on an interval $[a, b]$, find the areas between the graph of f and the interval $[a, b]$ on the x -axis.



From this diagram, S is bounded by the graph of a continuous function f [where $f(x) \geq 0$], the vertical lines $x=a$ and $x=b$ and the x axis.



For a rectangle, the area is defined as the product of the length and the width. The area of triangle is half of the base times the height. The area of a polygon is found by dividing it into triangles and adding the areas of the triangles. It is not easy to find the area of a region with curved sides. In defining a tangent we first approximated the slope of the tangent line by slopes of secant lines and then we took the limit of these approximations.

We follow a similar idea for areas. We first approximate the region S by rectangles and then we take the limit of the areas of these rectangles as we increase the number of rectangles.


The area problem.

① Use rectangles to estimate the area under the parabola $y = x^2$ from 0 to 1.

Solution:

Given that, the area S is between 0 and 1

We divide S into 4 strips S_1, S_2, S_3 and S_4 by drawing the vertical lines $x = 1/4, x = 1/2, x = 3/4,$



We can approximate each strip by a rectangle that has the same base as the strip and whose height is the same as the right edge of the strip.

The height of these rectangles are the values of the function $f(x) = x^2$ at the right end points of the of the sub-intervals. $[0, 1/4], [1/4, 1/2], [1/2, 3/4] \Delta [3/4, 1]$


Each rectangle has width $1/4$ and the height are $(1/4)^2, (1/2)^2, (3/4)^2$ and 1. If we let R_4 be the sum of the areas of these approximating rectangles, we get

$$R_4 = \frac{1}{4} \left(\frac{1}{4}\right)^2 + \frac{1}{4} \left(\frac{1}{2}\right)^2 + \frac{1}{4} \left(\frac{3}{4}\right)^2 + \frac{1}{4} (1)^2 = \frac{15}{32} = 0.46875$$

$\therefore$ From the above diagram we see that the area A of S is less than R_4 .

$$\therefore A < 0.46875$$

Instead of using the above rectangles, we can use the smaller rectangles from the following graph.



From the graph the heights are the values of f at the left end points of the sub intervals. The sum of the areas of these approximating rectangle is

$$L_4 = \frac{1}{4}(0)^2 + \frac{1}{4}\left(\frac{1}{4}\right)^2 + \frac{1}{4}\left(\frac{1}{2}\right)^2 + \frac{1}{4}\left(\frac{3}{4}\right)^2$$

$$L_4 = \frac{7}{32} = 0.21875$$

We see that the area of S is larger than L_4 , so we've lower and upper estimates for A .

$$\therefore 0.21875 < A < 0.46875$$

We should repeat this procedure with a larger number of strips. Now the given region is subdivided into 8 strips of equal width.

$$\begin{aligned} L_8 &= \frac{1}{8}(0)^2 + \frac{1}{8}\left(\frac{1}{8}\right)^2 + \frac{1}{8}\left(\frac{2}{8}\right)^2 + \frac{1}{8}\left(\frac{3}{8}\right)^2 + \frac{1}{8}\left(\frac{4}{8}\right)^2 \\ &\quad + \frac{1}{8}\left(\frac{5}{8}\right)^2 + \frac{1}{8}\left(\frac{6}{8}\right)^2 + \frac{1}{8}\left(\frac{7}{8}\right)^2 \\ &= \frac{1}{8} \left[\frac{1}{64} + \frac{1}{16} + \frac{9}{64} + \frac{1}{4} + \frac{25}{64} + \frac{9}{16} + \frac{49}{64} \right] \end{aligned}$$

$$\therefore L_8 = 0.2734375$$

$$\begin{aligned} R_8 &= \frac{1}{8}\left(\frac{1}{8}\right)^2 + \frac{1}{8}\left(\frac{2}{8}\right)^2 + \frac{1}{8}\left(\frac{3}{8}\right)^2 + \frac{1}{8}\left(\frac{4}{8}\right)^2 + \frac{1}{8}\left(\frac{5}{8}\right)^2 + \frac{1}{8}\left(\frac{6}{8}\right)^2 \\ &\quad + \frac{1}{8}\left(\frac{7}{8}\right)^2 + \frac{1}{8}(1)^2 \end{aligned}$$

$$R_8 = 0.3984375$$

$$\therefore 0.2734375 < A < 0.3984375$$

We can obtain better estimates by increasing the no. of strips

$\therefore$ A good estimate is obtained by averaging these numbers

n	L_n	R_n
10	0.285000	0.385000
20	0.3087500	0.3587500
30	0.3168519	0.3501852
50	0.3234000	0.3434000
100	0.328125	0.338125

② For the region S in $y=x^2$ from 0 to 1, show that the sum of the areas of the upper approximating rectangles, approaches $\frac{1}{3}$. i.e. $\lim_{n \rightarrow \infty} R_n = \frac{1}{3}$.

Solution: Let R_n be the sum of the areas of the n rectangles.

Each rectangle has width $\frac{1}{n}$ and the heights are the values of the function $f(x)=x^2$ at the points $\frac{1}{n}, \frac{2}{n}, \frac{3}{n}, \dots, \frac{n}{n}$. i.e. the heights are $(\frac{1}{n})^2, (\frac{2}{n})^2, (\frac{3}{n})^2, \dots, (\frac{n}{n})^2$.

$$\begin{aligned} \text{Then } R_n &= \frac{1}{n} \left(\frac{1}{n} \right)^2 + \frac{1}{n} \left(\frac{2}{n} \right)^2 + \dots + \frac{1}{n} \left(\frac{n}{n} \right)^2 \\ &= \frac{1}{n} \cdot \frac{1}{n^2} (1^2 + 2^2 + 3^2 + \dots + n^2) \end{aligned}$$

$$R_n = \frac{1}{n^3} \frac{n(n+1)(2n+1)}{6} = \frac{(n+1)(2n+1)}{6n^2}$$

$$\lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} \frac{(n+1)(2n+1)}{6n^2} = \lim_{n \rightarrow \infty} \frac{1}{6} n \frac{(1+\frac{1}{n})}{n} \cdot n \frac{(2+\frac{1}{n})}{n}$$

$$\therefore \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} \frac{1}{6} \left(1 + \frac{1}{n} \right) \left(2 + \frac{1}{n} \right) = \frac{1}{6} \cdot 1 \cdot 2 = \frac{1}{3}.$$

i.e. the sum of the areas of the upper approximating rectangles approaches $\frac{1}{3}$.

Definition:

The area of a region S that lies under the graph of the continuous function f is the limit of the sum of the areas of approximating rectangles.

$$A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} [f(x_1)\Delta x + f(x_2)\Delta x + f(x_3)\Delta x + \dots + f(x_n)\Delta x]$$

We can get the same value for left end points.

$$A = \lim_{n \rightarrow \infty} L = \lim_{n \rightarrow \infty} [f(x_0)\Delta x + f(x_1)\Delta x + f(x_2)\Delta x + \dots + f(x_{n-1})\Delta x]$$

$$\therefore A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i-1})\Delta x.$$

$[0, 1]$.

Solution:

Dividing $[0, 1]$ into n strips of equal length

$$\Delta x = \frac{1-0}{n} = \frac{1}{n}.$$

$$x_0 = 0, \quad x_1 = 0 + \frac{1}{n} = \frac{1}{n} \quad x_2 = 0 + 2 \cdot \frac{1}{n} = \frac{2}{n}, \dots, x_n = 1.$$

If R_n is the right end point approximation using n approximating rectangles, then

$$A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x.$$

$$A = \lim_{n \rightarrow \infty} \left[\left(\frac{1}{n}\right)^3 \frac{1}{n} + \left(\frac{2}{n}\right)^3 \frac{1}{n} + \left(\frac{3}{n}\right)^3 \frac{1}{n} + \dots + \left(\frac{n}{n}\right)^3 \frac{1}{n} \right]$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n^4} (1^3 + 2^3 + \dots + n^3)$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n^4} \left(\frac{n(n+1)}{2} \right)^2$$

$$= \frac{1}{4} \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n} \right)^2$$

$$\therefore A = \frac{1}{4}.$$

Similarly, if L_n is the ~~right~~ left end point approximation using n approximating rectangles, then.

$$A = \lim_{n \rightarrow \infty} L_n = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i-1}) \Delta x.$$

$$= \lim_{n \rightarrow \infty} \left[\left(\frac{0}{n}\right)^3 \left(\frac{1}{n}\right) + \left(\frac{1}{n}\right)^3 \left(\frac{1}{n}\right) + \left(\frac{2}{n}\right)^3 \left(\frac{1}{n}\right) + \dots + \left(\frac{n-1}{n}\right)^3 \left(\frac{1}{n}\right) \right]$$

$$= \lim_{n \rightarrow \infty} \left[\frac{1}{n^4} (1^3 + 2^3 + 3^3 + \dots + (n-1)^3) \right]$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n^4} \left(\frac{(n-1)n}{2} \right)^2$$

$$= \frac{1}{4} \lim_{n \rightarrow \infty} \frac{n^4 \left(1 - \frac{1}{n} \right)^2}{n^4}$$

$$\therefore A = \frac{1}{4}.$$

The Definite Integral

The limit of the form

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x = \lim_{n \rightarrow \infty} [f(x_1^*) \Delta x + f(x_2^*) \Delta x + \cdots + f(x_n^*) \Delta x]$$

If f is a function defined for $a \leq x \leq b$, we divide $[a, b]$ into n sub-intervals of equal width $\Delta x = \frac{b-a}{n}$. Let $x_0 = a, x_1, x_2, \dots, x_n = b$ be the end points of these sub-intervals and let $x_1^*, x_2^*, x_3^*, \dots, x_n^*$ be any sample points in these sub-intervals. So x_i^* lies in the i^{th} sub-interval $[x_{i-1}, x_i]$. Then the definite integral of f from a to b is given by

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x.$$

provided that this limit exists and gives the same value for all possible points. If it exists, then f is integrable on $[a, b]$.

Theorem 1

If f is continuous on $[a, b]$ or if f has only a finite number of discontinuities, then f is integrable on $[a, b]$. i.e. The definite integral $\int_a^b f(x) dx$ exists.

Theorem 2

If f is integrable on $[a, b]$ then

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x$$

where $\Delta x = \frac{b-a}{n}$ and $x_i = a + i\Delta x$

Properties of Definite integral

Consider the integral $\int_a^b f(x) dx$.

Let $a < b$. Then $\Delta x = \frac{b-a}{n}$

$$\therefore \int_a^b f(x) dx = - \int_b^a f(x) dx.$$

If $a=b$, then $\Delta x=0$, & $\int_a^b f(x) dx = 0$

Let us assume that f and g are continuous functions

Then (i) $\int_a^b c dx = c(b-a)$, where c is any constant

$$(ii) \int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx.$$

$$(iii) \int_a^b c f(x) dx = c \int_a^b f(x) dx, \text{ where } c \text{ is any constant.}$$

$$(iv) \int_a^b [f(x) - g(x)] dx = \int_a^b f(x) dx - \int_a^b g(x) dx.$$

$$(v) \int_a^c f(x) dx + \int_c^b f(x) dx = \int_a^b f(x) dx.$$

(vi) If $f(x) \geq 0$ for $a \leq x \leq b$, then $\int_a^b f(x) dx \geq 0$.

(vii) If $f(x) \geq g(x)$ for $a \leq x \leq b$, then $\int_a^b f(x) dx \geq \int_a^b g(x) dx$.

(viii) If $m \leq f(x) \leq M$ for $a \leq x \leq b$, then

$$m(b-a) \leq \int_a^b f(x) dx \leq M(b-a)$$

④ Let A be the area of the region that lies under the graph of $f(x) = e^{-x}$ between $x=0$ and $x=2$. (a) Using right end points, Find an expression for A as a limit. (b) Estimate the area by taking the sample points to be mid-points and using four sub-intervals and then ten sub-intervals.

Solution:

(a) Since $a=0$ and $b=2$, then $\Delta x = \frac{2-0}{n} = \frac{2}{n}$.

$$\therefore x_1 = \frac{2}{n}, x_2 = \frac{4}{n}, x_3 = \frac{6}{n}, \dots, x_i = \frac{2i}{n} \text{ \& } x_n = \frac{2n}{n}$$

$$\begin{aligned}\therefore R_n &= f(x_1)\Delta x + f(x_2)\Delta x + \dots + f(x_n)\Delta x \\ &= e^{-2/n} \left(\frac{2}{n}\right) + e^{-4/n} \left(\frac{2}{n}\right) + \dots + e^{-2n/n} \left(\frac{2}{n}\right)\end{aligned}$$

$$A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} \frac{2}{n} \left[e^{-2/n} + e^{-4/n} + \dots + e^{-2n/n} \right]$$

$$A = \lim_{n \rightarrow \infty} \frac{2}{n} \sum_{i=1}^n e^{-2i/n}$$

(b) When $n=4$, ~~$\Delta x = 0.5$~~ $\Delta x = 0.5$. The sub-intervals are $[0, 0.5]$, $[0.5, 1.0]$, $[1.0, 1.5]$ and $[1.5, 2]$.

The mid-points are $x_1^* = 0.25$, $x_2^* = 0.75$, $x_3^* = 1.25$ and $x_4^* = 1.75$

$$M_4 = \sum_{i=1}^4 f(x_i^*) \Delta x = f(0.25)\Delta x + f(0.75)\Delta x + f(1.25)\Delta x + f(1.75)\Delta x.$$

$$= e^{-0.25}(0.5) + e^{-0.75}(0.5) + e^{-1.25}(0.5) + e^{-1.75}(0.5)$$

$$= \frac{1}{2} (e^{-0.25} + e^{-0.75} + e^{-1.25} + e^{-1.75})$$

$$M_4 = 0.8557$$

When $n=10$ the sub-intervals are $[0, 0.2]$, $[0.2, 0.4]$, $[0.4, 0.6]$, $[0.6, 0.8]$, $[0.8, 1.0]$, $[1.0, 1.2]$, $[1.2, 1.4]$, $[1.4, 1.6]$, $[1.6, 1.8]$, $[1.8, 2.0]$

The mid points are 0.1, 0.3, 0.5, 0.7, 0.9, 1.1, 1.3, 1.5, 1.7 and 1.9.

$$A = M_{10} = f(0.1)\Delta x + f(0.3)\Delta x + f(0.5)\Delta x + f(0.7)\Delta x + \dots + f(1.9)\Delta x$$

$$= 0.2 [e^{-0.1} + e^{-0.3} + e^{-0.5} + e^{-0.7} + e^{-0.9} + e^{-1.1} + e^{-1.3} + e^{-1.5} + e^{-1.7} + e^{-1.9}]$$

$$A = 0.8632$$

- ⑤ Evaluate the Riemann sum for $f(x) = x^3 - 6x$, taking the sample points to be right end points and $a=0, b=3$ and $n=6$. Also evaluate $\int_0^3 (x^3 - 6x) dx$.

Solution:-

① When $n=6$, $\Delta x = \frac{b-a}{n} = \frac{1}{2}$

$\therefore$ the right end points are $x_1=0.5, x_2=1.0, x_3=1.5, x_4=2.0, x_5=2.5$ and $x_6=3.0$.

The Riemann sum is

$$\begin{aligned} R_6 &= \sum_{i=1}^6 f(x_i) \Delta x = f(x_1) \Delta x + f(x_2) \Delta x + f(x_3) \Delta x + f(x_4) \Delta x + f(x_5) \Delta x + f(x_6) \Delta x \\ &= f(0.5) \Delta x + f(1.0) \Delta x + f(1.5) \Delta x + f(2.0) \Delta x + f(2.5) \Delta x + f(3.0) \Delta x \end{aligned}$$

$$= \frac{1}{2} [-2.875 - 5 - 5.625 - 4 + 0.625 + 9]$$

$$R_6 = -3.9375$$

- ⑥ With n sub-intervals, we've $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.

$$\Rightarrow x_0=0, x_1=\frac{3}{n}, x_2=\frac{6}{n}, x_3=\frac{9}{n} \text{ and in general } x_i = \frac{3i}{n}$$

$$\int_0^3 (x^3 - 6x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{3i}{n}\right) \frac{3}{n}$$

$$= \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\left(\frac{3i}{n}\right)^3 - 6\left(\frac{3i}{n}\right) \right]$$

$$= \lim_{n \rightarrow \infty} \left[\frac{81}{n^4} \sum_{i=1}^n i^3 - \frac{54}{n^2} \sum_{i=1}^n i \right]$$

$$= \lim_{n \rightarrow \infty} \left[\frac{81}{n^4} \left(\frac{n(n+1)}{2} \right)^2 - \frac{54}{n^2} \left(\frac{n(n+1)}{2} \right) \right]$$

$$= \lim_{n \rightarrow \infty} \left[\frac{81}{4} \left(1 + \frac{1}{n} \right)^2 - 27 \left(1 + \frac{1}{n} \right) \right]$$

$$= \frac{81}{4} - 27 \quad \left\{ \because \sum_{i=1}^n i^3 = \left(\frac{n(n+1)}{2} \right)^2 \right.$$

$$\int_0^3 (x^3 - 6x) dx = -6.75$$

$$\left. \sum_{i=1}^n i = \frac{n(n+1)}{2} \right\}$$


⑥ Evaluate the following integrals by interpreting each in terms of areas. ① $\int_0^1 \sqrt{1-x^2} dx$ ② $\int_0^3 (x-1) dx$

Solution:

① Let $f(x) = \sqrt{1-x^2}$. This integral is the area under the curve $y = \sqrt{1-x^2}$ from 0 to 1.

Since $y^2 = 1-x^2 \Rightarrow x^2 + y^2 = 1$. (i.e. a quadratic circle with radius 1.)


$$\therefore \int_0^1 \sqrt{1-x^2} dx = \frac{1}{4} \pi r^2 = \frac{1}{4} \pi (1) = \pi/4.$$



② The graph of $y = x-1$ is the line with slope 1.

We compute the integral as the difference of the areas of the two triangles.

$$\begin{aligned} \int_0^3 (x-1) dx &= A_1 - A_2 \\ &= \frac{1}{2} (2 \times 2) - \frac{1}{2} (1 \times 1) \\ \int_0^3 (x-1) dx &= 1.5 \end{aligned}$$



⑦ Use mid point rule with $n=5$, to approximate $\int_1^2 \frac{1}{x} dx$.

Solution:

Let $a=1, b=2, n=5$

Then $\Delta x = \frac{b-a}{n} = \frac{1}{5}$.

$\therefore$ the end points of the sub-intervals are 1, 1.2, 1.4, 1.6, 1.8 and 2.

Also the mid points are 1.1, 1.3, 1.5, 1.7 & 1.9

$\therefore$ The mid point rule gives

$$\begin{aligned} \int_1^2 \frac{1}{x} dx &= \Delta x [f(1.1) + f(1.3) + f(1.5) + f(1.7) + f(1.9)] \\ &= \frac{1}{5} \left[\frac{1}{1.1} + \frac{1}{1.3} + \frac{1}{1.5} + \frac{1}{1.7} + \frac{1}{1.9} \right] \\ &= 0.691908 \end{aligned}$$

Since $f(x) = \frac{1}{x} > 0$ for $1 \leq x \leq 2$ the integral represents an area, and the approximation given by the mid-point rule is the sum of the area of the rectangles.

⑧ Prove that (a) $\int_a^b x dx = \frac{b^2 - a^2}{2}$ & (b) $\int_a^b x^2 dx = \frac{b^3 - a^3}{3}$

Solution:

(a) With n sub-intervals, we have $\Delta x = \frac{b-a}{n}$

$\Rightarrow x_i = a + \frac{(b-a)}{n} i$. To evaluate the integral, we use

Riemann sum,

$$\begin{aligned} \int_a^b x dx &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left(\frac{b-a}{n} \right) \left(a + \frac{(b-a)}{n} i \right) \\ &= \lim_{n \rightarrow \infty} \left(\frac{b-a}{n} \right) \sum_{i=1}^n \left[a + \frac{(b-a)}{n} i \right] \\ &= \lim_{n \rightarrow \infty} \frac{b-a}{n} \left[\sum_{i=1}^n a + \sum_{i=1}^n \left(\frac{b-a}{n} \right) i \right] \\ &= \lim_{n \rightarrow \infty} \left[\left(\frac{b-a}{n} \right) a \sum_{i=1}^n 1 + \left(\frac{b-a}{n} \right)^2 \sum_{i=1}^n i \right] \\ &= \lim_{n \rightarrow \infty} \left[a \left(\frac{b-a}{n} \right) \cdot n + \left(\frac{b-a}{n} \right)^2 \frac{n(n+1)}{2} \right] \\ &= \lim_{n \rightarrow \infty} \left[a(b-a) + \frac{(b-a)^2}{2} \left(1 + \frac{1}{n} \right) \right] \end{aligned}$$

$$\begin{aligned} &= ab - a^2 + \frac{1}{2} (a^2 - 2ab + b^2) \cdot 1 \\ &= \frac{1}{2} [2ab - 2a^2 + a^2 - 2ab + b^2] \end{aligned}$$

$$\int_a^b x dx = \frac{b^2 - a^2}{2}$$

(b) With n sub-intervals, we've $\Delta x = \frac{b-a}{n} \Rightarrow x_i = a + \frac{(b-a)}{n} i$

$$\begin{aligned} \therefore \int_a^b x^2 dx &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left(\frac{b-a}{n} \right) \left[a + \frac{(b-a)}{n} i \right]^2 \\ &= \lim_{n \rightarrow \infty} \frac{b-a}{n} \sum_{i=1}^n \left[a + \frac{(b-a)}{n} i \right]^2 \\ &= \lim_{n \rightarrow \infty} \left(\frac{b-a}{n} \right) \sum_{i=1}^n \left[a^2 + 2a \left(\frac{b-a}{n} \right) i + \left(\frac{b-a}{n} \right)^2 i^2 \right] \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[a^2 \frac{(b-a)}{n} + 2a \left(\frac{b-a}{n} \right)^2 i + \left(\frac{b-a}{n} \right)^3 i^2 \right] \end{aligned}$$

$$\int_a^b x^2 dx = \lim_{n \rightarrow \infty} \left[\frac{a^2(b-a)}{n} \sum_{i=1}^n 1 + 2a \left(\frac{b-a}{n} \right)^2 \sum_{i=1}^n i + \left(\frac{b-a}{n} \right)^3 \sum_{i=1}^n i^2 \right]$$

$$= a^2(b-a) + 2a(b-a)^2 \lim_{n \rightarrow \infty} \frac{1}{n^2} \frac{n(n+1)}{2} + (b-a)^3 \lim_{n \rightarrow \infty} \frac{1}{n^3} \frac{n(n+1)(2n+1)}{6}$$

$$= a^2(b-a) + a(b-a)^2 + \frac{(b-a)^3}{3}$$

$$= a^2b - a^3 + ab^2 - 2a^2b - a^3 + \frac{b^3}{3} - ab^2 + a^2b - \frac{a^3}{3}$$

$$\int_a^b x^2 dx = \frac{b^3 - a^3}{3}$$

⑨ Find the Riemann sum for $f(x) = \sin x$, $0 \leq x \leq \frac{3\pi}{2}$ with six terms, taking the sample points to be right end-points correct to six decimal places. Repeat the problem with mid-points as the sample points.

Solution:

given that $f(x) = \sin x$, $0 \leq x \leq \frac{3\pi}{2}$

$\Delta x = \frac{b-a}{2} = \frac{\pi}{4}$. Since we are using right end-points, $x_i^* = x_i$

$$\text{Now } R_6 = \sum_{i=1}^6 f(x_i) \Delta x = \Delta x [f(x_1) + f(x_2) + \dots + f(x_6)]$$

$$= \frac{\pi}{4} \left[f\left(\frac{\pi}{4}\right) + f\left(\frac{2\pi}{4}\right) + f\left(\frac{3\pi}{4}\right) + f(\pi) + f\left(\frac{5\pi}{4}\right) + f\left(\frac{6\pi}{4}\right) \right]$$

$$= \frac{\pi}{4} \left[\sin \frac{\pi}{4} + \sin \frac{\pi}{2} + \sin \frac{3\pi}{4} + \sin \pi + \sin \frac{5\pi}{4} + \sin \left(\frac{3\pi}{2}\right) \right]$$

$$= \frac{\pi}{4} \left[\frac{1}{\sqrt{2}} + 1 + \frac{1}{\sqrt{2}} + 0 - \frac{1}{\sqrt{2}} - 1 \right]$$

$$= \frac{\pi}{4} \frac{1}{\sqrt{2}}$$

$$R_6 = 0.555360$$

Since we are using the mid points $x_i^* = \bar{x}_i = \frac{1}{2}(x_{i-1} + x_i)$

① Find the derivative of the function $g(x) = \int_0^x \sqrt{1+t^2} dt$

Sol.

Since $f(t) = \sqrt{1+t^2}$ is continuous, Part 1 of the fundamental theorem of calculus gives

$$g'(x) = \sqrt{1+x^2}$$

② Evaluate the integral $\int_1^3 e^x dx$.

Sol.

Since $f(x) = e^x$ is continuous every where
Wkt anti-derivative of $f(x)$ is $F(x) = e^x$.

So part 2 of fundamental theorem gives

$$\int_1^3 e^x dx = F(3) - F(1) = e^3 - e^1.$$

③ What is wrong with the following calculation.

$$\int_{-1}^3 \left(\frac{1}{x^2}\right) dx = \left[\frac{x^{-1}}{-1} \right]_{-1}^3 = -\frac{4}{3}.$$

Sol.

Since $f(x) = \frac{1}{x^2} \geq 0$, $\int_a^b f(x) dx \geq 0$ when $f(x) \geq 0$.

Also $\frac{1}{x^2}$ is discontinuous on $[-1, 3]$, $f(x) = \frac{1}{x^2}$ has an infinite discontinuity at $x=0$.

$\therefore \int_{-1}^3 \left(\frac{1}{x^2}\right) dx$ does not exist.

④ Find the derivative of $y = \int_0^{\tan x} \sqrt{t+\sqrt{t}} dt$.

Sol

Let $u = \tan x$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

Since $\frac{du}{dx} = \sec^2 x$

$$\begin{aligned} \text{Then } \frac{d}{dx} \left\{ \int_0^{\tan x} \sqrt{t+\sqrt{t}} dt \right\} &= \frac{d}{dx} \left\{ \int_0^u \sqrt{t+\sqrt{t}} dt \right\} \\ &= \frac{d}{du} \left\{ \int_0^u \sqrt{t+\sqrt{t}} dt \right\} \frac{du}{dx} = \sqrt{u+\sqrt{u}} \sec^2 x \\ &= \sqrt{\tan x + \sqrt{\tan x}} \sec^2 x. \end{aligned}$$

Indefinite Integral

In calculus, an indefinite integral of a function $f(x)$ is a differentiable function F whose derivative is equal to the original function, $f(x)$. i.e. $F' = f(x)$.

Remark:

From the definite and indefinite integrals, we note that a definite integral $\int_a^b f(x) dx$ is a number, whereas an indefinite integral $\int f(x) dx$ is a function. The relation between these two integrals is given by part 2 of the fundamental theorem.

Formulae.

- ① $\int c f(x) dx = c \int f(x) dx$
 c is constant.
- ② $\int k dx = kx + c$
- ③ $\int x^n dx = \frac{x^{n+1}}{n+1} + c, n \neq -1$
- ④ $\int \frac{1}{x} dx = \log|x| + c$
- ⑤ $\int e^x dx = e^x + c$
- ⑥ $\int a^x dx = a^x \frac{1}{\log a} + c$
- ⑦ $\int \sin x dx = -\cos x + c$
- ⑧ $\int \cos x dx = \sin x + c$
- ⑨ $\int \sec^2 x dx = \tan x + c$
- ⑩ $\int \csc^2 x dx = -\cot x + c$
- ⑪ $\int \sec x \tan x dx = \sec x + c.$

- ⑫ $\int \operatorname{cosec} x \cot x dx = -\operatorname{cosec} x + c$
- ⑬ $\int \frac{dx}{x^2+1} = \tan^{-1} x \text{ (or) } -\cot^{-1}(x) + c$
- ⑭ $\int \frac{dx}{\sqrt{1-x^2}} = \sin^{-1}(x) + c \text{ or } -\cos^{-1}(x) + c$
- ⑮ $\int \sinh x dx = \cosh x + c$
- ⑯ $\int \cosh x dx = \sinh x + c$
- ⑰ $\int \operatorname{sech}^2 x dx = \tanh x + c$
- ⑱ $\int \operatorname{cosech}^2 x dx = -\coth x + c$
- ⑲ $\int \operatorname{sech} x \tanh x dx = -\operatorname{sech} x + c$
- ⑳ $\int \operatorname{cosech} x \coth x dx = -\operatorname{cosech} x + c$
- ㉑ $\int \frac{dx}{x(x^2-1)} = \sec^{-1}(x) + c \text{ or } \operatorname{cosec}^{-1}(x) + c$
- ㉒ $\int \frac{dx}{\sqrt{1+x^2}} = \sinh^{-1}(x) + c$
- ㉓ $\int \frac{dx}{\sqrt{x^2-1}} = \cosh^{-1} x + c$
- ㉔ $\int \frac{dx}{x^2-1} = \tanh^{-1} x + c \text{ (or) } \coth^{-1}(x) + c.$

① Evaluate $\int (10x^4 - 2\sec^2 x) dx$.
Sol Let $I = \int (10x^4 - 2\sec^2 x) dx$.

$$= 10 \frac{x^5}{5} - 2 \tan x + C$$

$$\therefore I = 2x^5 - 2 \tan x + C$$

② $\int \frac{\cos \theta}{\sin^2 \theta} d\theta = ?$

Sol: Let $I = \int \frac{\cos \theta}{\sin^2 \theta} d\theta$.

$$I = \int \frac{\cos \theta}{\sin \theta} \cdot \frac{1}{\sin \theta} d\theta$$

$$I = \int \cot \theta \cdot \operatorname{cosec} \theta d\theta$$

$$\therefore I = -\operatorname{cosec} \theta + C$$

③ Evaluate $\int [\sqrt{x^3} + \sqrt[3]{x^2}] dx$

Sol. Let $I = \int [\sqrt{x^3} + \sqrt[3]{x^2}] dx$.

Then $I = \int (x^{3/2} + x^{2/3}) dx$.

$$I = \frac{x^{5/2}}{5/2} + \frac{x^{5/3}}{5/3} + C$$

$$= \frac{1}{5} [2x^{5/2} + 3x^{5/3}] + C$$

④ $\int \frac{\sin 2x}{\sin x} dx = ?$

Sol
$$\int \frac{\sin 2x}{\sin x} dx = \int \frac{2 \sin x \cos x}{\sin x} dx$$

$$= \int 2 \cos x dx$$

$$= 2 \sin x + C$$

⑤ $\int [(x^2+1) + \frac{1}{x^2+1}] dx = ?$

Sol
$$\int [x^2+1 + \frac{1}{x^2+1}] dx$$

$$= \frac{x^3}{3} + x + \tan^{-1} x + C$$

⑥ Evaluate $\int_1^2 (\frac{1}{x^2} - \frac{4}{x^3}) dx$.

Sol

$$I = \int_1^2 (\frac{1}{x^2} - \frac{4}{x^3}) dx = \int_1^2 (x^{-2} - 4x^{-3}) dx$$

$$I = \left[\frac{x^{-2+1}}{-2+1} - 4 \frac{x^{-3+1}}{-3+1} \right]_1^2$$

$$\therefore I = \left[\frac{x^{-1}}{-1} + \frac{4}{3} x^{-2} \right]_1^2 = \frac{-1}{2} + \frac{1}{2} - 2$$

$$\therefore I = -1$$

⑦ Evaluate $\int_0^1 (5x - 5^x) dx$.

Sol
$$I = \int_0^1 (5x - 5^x) dx$$

$$I = 5 \left[\frac{x^2}{2} \right]_0^1 - \left[\frac{5^x}{\log 5} \right]_0^1$$

$$= \frac{5}{2} [1-0] - \frac{1}{\log 5} [5^1 - 5^0]$$

$$I = \frac{5}{2} - \frac{4}{\log 5}$$

⑧ Evaluate $\int_{\pi/4}^{\pi/3} \operatorname{cosec}^2 \theta d\theta$.

Sol

$$\int_{\pi/4}^{\pi/3} \operatorname{cosec}^2 \theta d\theta = \left[-\cot \theta \right]_{\pi/4}^{\pi/3}$$

$$= -[\cot \pi/3 - \cot \pi/4]$$

$$= \cot \pi/4 - \cot \pi/3$$

—

$$= \int \frac{\sin^2 x + \cos^2 x}{\sin^2 x \cos^2 x} dx.$$

$$= \int \frac{1}{\cos^2 x} dx + \int \frac{1}{\sin^2 x} dx.$$

$$= \int \sec^2 x dx + \int \csc^2 x dx.$$

$$= \tan x - \cot x + C.$$

$$(2) \int \sqrt{1 + \sin 2x} dx.$$

$$\text{Sol } \int \sqrt{1 + \sin 2x} dx.$$

$$= \int \sqrt{\cos^2 x + \sin^2 x + 2 \sin x \cos x} dx$$

$$= \int \sqrt{(\cos x + \sin x)^2} dx$$

$$= \int (\cos x + \sin x) dx.$$

$$= \sin x - \cos x + C.$$

$$(3) \int \frac{1}{1 - \cos x} dx.$$

$$\text{Sol } \int \frac{1}{1 - \cos x} dx$$

$$= \int \frac{1}{1 - \cos x} \times \frac{1 + \cos x}{1 + \cos x} dx$$

$$= \int \frac{1 + \cos x}{1 - \cos^2 x} dx.$$

$$= \int \frac{1 + \cos x}{(1 - \cos x)(1 + \cos x)} dx.$$

$$= \int \frac{1 + \cos x}{\sin^2 x} dx.$$

$$= -\cot x - \csc x + C.$$

Properties of Definite Integrals

We assume that f and g are continuous functions

$$1. \int_a^b f(x) dx = \int_a^b f(t) dt$$

$$2. \int_a^b f(x) dx = - \int_b^a f(x) dx.$$

$$3. \int_a^c f(x) dx = \int_a^b f(x) dx + \int_b^c f(x) dx$$

$a < b < c.$

$$4. \int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$5. \int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a-x) dx$$

$$6. \int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a-x) dx$$

iff $f(2a-x) = f(x)$

$$(ii) \int_0^{2a} f(x) dx = 0 \text{ iff } f(2a-x) = -f(x)$$

$$7(i) \text{ If } f(x) \text{ is an even func. then } \int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx.$$

$$(ii) \text{ If } f(x) \text{ is an odd func. then } \int_{-a}^a f(x) dx = 0.$$

① Evaluate $\int_{-1}^1 (2-|x|) dx$.

Sol Let $I = \int_{-1}^1 (2-|x|) dx$

Then $I = \int_{-1}^1 2 dx - \int_{-1}^1 |x| dx$.

$= 4 - 2 \int_0^1 x dx$ { $|x|$ is even }

$= 4 - 2 \left[\frac{x^2}{2} \right]_0^1 \Rightarrow I = 3$

② $\int_0^{\pi/2} \frac{1}{1+\tan x} dx$.

Let $I = \int_0^{\pi/2} \frac{1}{1+\tan x} dx$.

$I = \int_0^{\pi/2} \frac{\cos x}{\sin x + \cos x} dx$ — (1)

{ $\therefore \int_a^b f(x) dx = \int_a^b f(a+x) dx$ }

$\therefore I = \int_0^{\pi/2} \frac{\cos(\pi/2-x)}{\sin(\pi/2-x) + \cos(\pi/2-x)} dx$

$\Rightarrow I = \int_0^{\pi/2} \frac{\sin x}{\cos x + \sin x} dx$ — (2)

(1) + (2) $\Rightarrow \pi/2$

$2I = \int_0^{\pi/2} \frac{\sin x + \cos x}{\sin x + \cos x} dx$.

$2I = \int_0^{\pi/2} dx \Rightarrow 2I = [x]_0^{\pi/2}$

$\therefore I = \pi/4$.

③ $\int_0^{\pi} \frac{e^{\cos x}}{e^{\cos x} + e^{-\cos x}} dx$.

Sol Let $I = \int_0^{\pi} \frac{e^{\cos x}}{e^{\cos x} + e^{-\cos x}} dx$ — (1)

Then we've $\int_0^a f(x) dx = \int_0^a f(a-x) dx$

$\therefore I = \int_0^{\pi} \frac{e^{\cos(\pi-x)}}{e^{\cos(\pi-x)} + e^{-\cos(\pi-x)}} dx$

$I = \int_0^{\pi} \frac{e^{-\cos x}}{e^{-\cos x} + e^{\cos x}} dx$ — (2)

(1) + (2) $\Rightarrow 2I = \int_0^{\pi} \frac{e^{\cos x} + e^{-\cos x}}{e^{-\cos x} + e^{\cos x}} dx$

$\Rightarrow 2I = \int_0^{\pi} dx \Rightarrow 2I = [x]_0^{\pi} = \pi$

$\Rightarrow I = \pi/2$

④ $\int_{-2}^2 |x+1| dx$.

Sol Since $|x+1| = \begin{cases} -(x+1), & -2 \leq x < -1 \\ (x+1), & -1 \leq x \leq 2 \end{cases}$

$\therefore I = \int_{-2}^2 |x+1| dx = \int_{-2}^{-1} -(x+1) dx + \int_{-1}^2 (x+1) dx$

$\therefore I = \left[-\frac{x^2}{2} - x \right]_{-2}^{-1} + \left[\frac{x^2}{2} + x \right]_{-1}^2$

$= -\left[\frac{1}{2} - 0 \right] + \left[4 - \left(-\frac{1}{2} \right) \right]$

$I = 5$

⑤ Evaluate $\int_{-\pi/2}^{\pi/2} \sin^{199} x \, dx$.

Sol Let $I = \int_{-\pi/2}^{\pi/2} \sin^{199} x \, dx$

Let $f(x) = \sin^{199} x$

Then $f(-x) = \sin^{199}(-x)$
 $= -\sin^{199} x$
 $= -f(x)$

$\therefore f(x)$ is an ~~even~~ odd function.

$\therefore \int_{-\pi/2}^{\pi/2} \sin^{199} x \, dx = 0$.

$\therefore \int_{-a}^a f(x) \, dx = 0$, where $f(x)$ is odd.

Integration by Substitution rule.

Type I (A).

$\int [f(x)]^n f'(x) \, dx$ (or)

$\int \phi[f(x)] f'(x) \, dx$ — ①

Substitute $u = f(x)$,
 $du = f'(x) \, dx$

$\Rightarrow \int u^n \, du$ or $\int \phi(u) \, du$
 and then proceed.

① Solve $\int \frac{1}{x^2} \sqrt{2 - \frac{1}{x}} \, dx$.

Sol Let $I = \int \frac{1}{x^2} \sqrt{2 - \frac{1}{x}} \, dx$

$\therefore \int f'(x) \phi[f(x)] \, dx$

$\therefore u = 2 - \frac{1}{x} \quad du = \frac{1}{x^2} \, dx$

$\therefore I = \int \sqrt{u} \, du = \frac{2}{3} u^{3/2} + C$

Hence $I = \frac{2}{3} \left(2 - \frac{1}{x}\right)^{3/2} + C$.

② Evaluate $\int \frac{x^2}{\sqrt{x+5}} \, dx$.

Sol let $u = \sqrt{x+5}$

$du = \frac{1}{2\sqrt{x+5}} \, dx \Rightarrow 2du = \frac{dx}{\sqrt{x+5}}$

Now $u^2 = x+5 \Rightarrow x^2 = (u^2-5)^2$

$\therefore x^2 = u^4 - 10u^2 + 25$

$\therefore I = \int (u^4 - 10u^2 + 25) 2du$

$= 2 \left(\frac{u^5}{5} - \frac{10u^3}{3} + 25u \right) + C$

$= \frac{2}{5} (x+5)^{5/2} - \frac{20}{3} (x+5)^{3/2} + 50(x+5)^{1/2} + C$

③ Evaluate $\int \frac{1}{(3x-4)^{3/2}} \, dx$.

Sol Let $I = \int \frac{1}{(3x-4)^{3/2}} \, dx$.

Let $u = 3x-4 \quad du = 3 \, dx$
 $\Rightarrow \frac{du}{3} = dx$

$\therefore I = \int \frac{du/3}{u^{3/2}} = -\frac{2}{3} \frac{1}{\sqrt{u}} + C$

$\Rightarrow I = -\frac{2}{3} \frac{1}{\sqrt{3x-4}} + C$

⑥ Evaluate $\int e^{ax} \cos bx \, dx$.

Sol

Let $I = \int e^{ax} \cos bx \, dx$

$u = \cos bx \quad dv = e^{ax} \, dx$

$du = -\sin bx(b) \quad v = \frac{e^{ax}}{a}$

$\therefore I = \frac{e^{ax}}{a} \cos bx + \int \frac{e^{ax}}{a} \sin bx(b) \, dx$

$= \frac{e^{ax}}{a} \cos bx + \frac{b}{a} \int e^{ax} \sin bx \, dx$

Take $\int e^{ax} \sin bx \, dx$ ⑦

$u = \sin bx \quad dv = e^{ax} \, dx$

$du = b \cos bx \, dx \quad v = \frac{e^{ax}}{a}$

$\therefore \int e^{ax} \sin bx \, dx$

$= \frac{e^{ax}}{a} \sin bx - \int \frac{e^{ax}}{a} b \cos bx \, dx$

$= \frac{e^{ax}}{a} \sin bx - \frac{b}{a} I$

$\therefore \text{⑦} \Rightarrow I = \frac{e^{ax}}{a} \cos bx + \frac{b}{a} \left(\frac{e^{ax}}{a} \sin bx - \frac{b}{a} I \right)$

$\Rightarrow + \frac{b}{a} \frac{e^{ax}}{a} \sin bx - \frac{b^2}{a^2} I$

$\Rightarrow I + \frac{b^2}{a^2} I = \frac{e^{ax}}{a} \cos bx + b \frac{e^{ax}}{a^2} \sin bx$

$\Rightarrow I \left(\frac{a^2 + b^2}{a^2} \right) = \frac{e^{ax}}{a^2} [a \cos bx + b \sin bx]$

$\Rightarrow I = \frac{e^{ax}}{a^2 + b^2} [a \cos bx + b \sin bx] + c$

⑦ Evaluate $\int e^{ax} \sin bx \, dx$.

⑧ Evaluate $\int e^{-ax} \cos bx \, dx$.

⑨ Evaluate $\int e^{-ax} \sin bx \, dx$.

⑩ Find the reduction formula for $\int \sin^m x \, dx$ and also find $\int_0^{\pi/2} \sin^m x \, dx$, $\int_0^{\pi/2} \sin^5 x \, dx$

Sol Let $I_n = \int \sin^n x \, dx$

Then $I_n = \int \sin^{n-1} x \sin x \, dx$

$u = \sin^{n-1} x \quad dv = \sin x \, dx$

$du = (n-1) \sin^{n-2} x \cos x \, dx$

$v = -\cos x$

$\therefore I_n = -\cos x \sin^{n-1} x$

$+ (n-1) \int \cos x (\sin^{n-2} x \cos x \, dx)$

$= -\cos x \sin^{n-1} x$

$+ (n-1) \int \sin^{n-2} x \cos^2 x \, dx$

$= -\cos x \sin^{n-1} x$

$+ (n-1) \int \sin^{n-2} x (1 - \sin^2 x) \, dx$

$= -\cos x \sin^{n-1} x$

$+ (n-1) \int \sin^{n-2} x \, dx$

$- (n-1) \int \sin^n x \, dx$

$$\textcircled{2} \int \sec x \, dx = ?$$

Sol Let $I = \int \sec x \, dx$

~~Let~~ $I = \int \sec x \left(\frac{\sec x + \tan x}{\sec x + \tan x} \right) dx$

Let $u = \sec x + \tan x$

$du = (\sec x \tan x + \sec^2 x) dx$

$\therefore I = \int \frac{1}{u} du = \log u + C.$

$\Rightarrow I = \log(\sec x + \tan x) + C.$

$\textcircled{2}$ Evaluate $\int e^{\tan^{-1}x} \left[\frac{1+x+x^2}{1+x^2} \right] dx$

Let $I = \int e^{\tan^{-1}x} \left[\frac{1+x+x^2}{1+x^2} \right] dx$

Let $u = \tan^{-1}x \quad du = \frac{dx}{1+x^2}$

$\Rightarrow \tan u = x$

$1+x+x^2 = \tan u + \sec^2 u$

$\therefore I = \int e^u (\tan u + \sec^2 u) du$

put $t = e^u \tan u$

$dt = [e^u \sec^2 u + \tan u e^u] du$

$\therefore I = \int dt = t + C$

$\Rightarrow I = x e^{\tan^{-1}x} + C.$

TECHNIQUES OF INTEGRATIONS

Integration by parts method

$\int u \, dv = uv - \int v \, du.$

$\textcircled{1}$ Solve $\int_0^1 \frac{y}{e^{2y}} dy$

Let $I = \int_0^1 y e^{-2y} dy$

Let $u = y \quad dv = e^{-2y} dy$

$du = dy \quad v = e^{-2y} / -2$

$\therefore I = y \left(\frac{e^{-2y}}{-2} \right) \Big|_0^1 + \int_0^1 \frac{e^{-2y}}{2} dy$

$= -\frac{1}{2} e^{-2} + \frac{1}{2} \left[\frac{e^{-2y}}{-2} \right]_0^1$

$= -\frac{1}{2} e^{-2} - \frac{1}{4} [e^{-2} - 1]$

$\Rightarrow I = \frac{1}{4} - \frac{3}{4} e^{-2}$

$\textcircled{2}$ Evaluate $\int \left(\frac{\log x}{x} \right)^2 dx.$

Sol Let $I = \int (\log x)^2 / x^2 dx.$

~~Let~~ $u = (\log x)^2 \quad dv = \frac{1}{x^2} dx.$

$du = 2 \log x (1/x) dx \quad v = -\frac{1}{x}.$

$\therefore I = -\frac{1}{x} (\log x)^2 + 2 \int (\log x) \left(\frac{1}{x^2} \right) dx$

Consider $\int (\log x) \left(\frac{1}{x^2} \right) dx$ ①

Let $u = \log x \quad dv = dx / x^2$

$du = dx / x \quad v = -1/x.$

$\therefore \int (\log x) \left(\frac{1}{x^2} \right) dx$

$= -\frac{1}{x} \log x + \int \frac{1}{x^2} dx$

$= -\frac{1}{x} \log x + \frac{1}{x} + C.$

$\therefore I = -\frac{1}{x} (\log x)^2 - 2 \frac{1}{x} \log x - 2 \left(\frac{1}{x} \right)$

$+ 2C$

$$\Rightarrow n I_n = \sin x \cos^{n-1} x + (n-1) I_{n-2}$$

$$\therefore I_n = \frac{1}{n} \sin x \cos^{n-1} x + \frac{n-1}{n} I_{n-2}$$

$$\text{Now } I_n = \int_0^{\pi/2} \cos^n x \, dx.$$

$$= \left. \frac{1}{n} \sin x \cos^{n-1} x \right|_0^{\pi/2} + \frac{n-1}{n} I_{n-2}$$

$$I_n = \frac{n-1}{n} I_{n-2}; I_{n-2} = \frac{n-3}{n-2} I_{n-4}$$

In general,

$$I_n = \begin{cases} \frac{n-1}{n} \frac{n-3}{n-2} \dots \frac{1}{2} I_0 & (n \text{ is even}) \\ \frac{n-1}{n} \frac{n-3}{n-2} \dots \frac{2}{3} I_1 & (n \text{ is odd}) \end{cases}$$

$$I_0 = \int_0^{\pi/2} \cos^0 x \, dx = \pi/2$$

$$I_1 = \int_0^{\pi/2} \cos x \, dx = \sin x \Big|_0^{\pi/2}$$

$$\therefore I_1 = 1$$

$$\textcircled{C} \Rightarrow I_n = \begin{cases} \frac{n-1}{n} \frac{n-3}{n-2} \dots \frac{1}{2} \cdot \frac{\pi}{2}, & n \text{ is even} \\ \frac{n-1}{n} \frac{n-3}{n-2} \dots \frac{2}{3} \cdot 1, & n \text{ is odd} \end{cases}$$

$$\text{also } \int_0^{\pi/2} \cos^9 x \, dx = \frac{8}{9} \cdot \frac{6}{7} \cdot \frac{4}{5} \cdot \frac{2}{3} \cdot 1$$

$$\therefore \int_0^{\pi/2} \cos^9 x \, dx = \frac{128}{315}$$

(12) Evaluate $\int \sec^n x \, dx$

$$\text{Let } I_n = \int \sec^n x \, dx. \quad \text{--- (1)}$$

$$\text{Then } I_n = \int \sec^{n-2} x \sec^2 x \, dx.$$

$$\int \sec^{n-2} x \, d(\tan x)$$

$$\int \sec^{n-2} x \, d(\tan x)$$

$$\int \sec^{n-2} x \, d(\tan x)$$

$$\therefore I_n = \int \sec^{n-2} x \, d(\tan x) \, dx.$$

$$u = \sec^{n-2} x$$

$$dv = d(\tan x)$$

$$du = (n-2) \sec^{n-3} x (\sec x \tan x \, dx)$$

$$\& \quad v = \tan x.$$

$$\therefore I_n = \tan x \sec^{n-2} x$$

$$- \int (n-2) \sec^{n-2} x \tan^2 x \, dx.$$

$$I_n = \tan x \sec^{n-2} x$$

$$- (n-2) \int \sec^{n-2} x (1 + \sec^2 x) \, dx$$

$$I_n = \tan x \sec^{n-2} x.$$

$$- (n-2) \left[\int \sec^2 x \, dx + \int \sec^n x \, dx \right]$$

$$I_n = \tan x \sec^{n-2} x$$

$$- (n-2) I_n + (n-2) I_{n-2}.$$

$$\Rightarrow I_n + (n-2) I_n$$

$$= \tan x \sec^{n-2} x + (n-2) I_{n-2}$$

$$\Rightarrow \frac{(n-1)}{n} I_n = \tan x \sec^{n-2} x + (n-2) I_{n-2}$$

$$\Rightarrow I_n = \frac{n-1}{n} \tan x \sec^{n-2} x + \frac{n-2}{n-1} I_{n-2}, \quad n \geq 2.$$

$$I_0 = \int dx = x + c \quad (\text{from (1)})$$

$$I_1 = \int \sec x \, dx = \log(\sec x + \tan x) + c$$

(13) Evaluate $\int (\log x)^n dx$.

also find $\int (\log x)^2 dx$.

Sol Let $I_n = \int (\log x)^n dx$.

$$u = (\log x)^n \quad dv = dx.$$

$$du = n(\log x)^{n-1} \frac{1}{x} dx \quad v = x.$$

$$\therefore I_n = x(\log x)^n - \int x n(\log x)^{n-1} \frac{1}{x} dx$$

$$= x(\log x)^n - n \int (\log x)^{n-1} dx.$$

$$\therefore I_n = x(\log x)^n - n I_{n-1}$$

$$I_0 = \int dx = x + c$$

To find $\int (\log x)^2 dx$,
put $n=2$ in I_n .

$$\therefore I_2 = \int (\log x)^2 dx$$

$$= x(\log x)^2 - 2I_1$$

$$= x(\log x)^2 - 2I_1 \quad \text{--- (1)}$$

$$\text{But } I_1 = x \log x - I_0$$

$$\therefore I_1 = x \log x - x + c$$

$$\therefore \text{--- (1)} \Rightarrow I_2 = x(\log x)^2 - 2(x \log x - x) + c$$

$$I_2 = x(\log x)^2 - 2x \log x + 2x + c.$$

(14) Find the reduction formula for $\int x^n \sin mx dx$

Sol Let $I_n = \int x^n \sin mx dx$.

$$= \int x^n d\left(-\frac{\cos mx}{m}\right)$$

$$= -x^n \frac{\cos mx}{m} + \int \frac{\cos mx}{m} d(mx)$$

$$= -x^n \frac{\cos mx}{m} + \frac{1}{m} \int \cos mx (n x^{n-1}) dx.$$

$$= -x^n \frac{\cos mx}{m} + \frac{n}{m} \int x^{n-1} d\left(\frac{\sin mx}{m}\right)$$

$$= -x^n \frac{\cos mx}{m} + \frac{n}{m} \left[x^{n-1} \frac{\sin mx}{m} - \int \frac{\sin mx}{m} (n-1) x^{n-2} dx \right]$$

$$= -x^n \frac{\cos mx}{m} + \frac{n}{m^2} x^{n-1} \sin mx - \frac{n(n-1)}{m^2} \int x^{n-2} \sin mx dx.$$

$$\therefore I_n = -x^n \frac{\cos mx}{m} + \frac{n}{m^2} x^{n-1} \sin mx - \frac{(n-1)n}{m^2} I_{n-2}.$$

(15) Evaluate $\int x^n e^x dx$

$$\text{Let } u = x^n \quad dv = e^x dx$$

$$du = nx^{n-1} dx \quad v = e^x$$

$$\therefore \int x^n e^x dx = x^n e^x - \int e^x nx^{n-1} dx$$

$$\therefore I_n = x^n e^x - n I_{n-1}$$

Ans.

$$\therefore I_n = -\cos x \sin^{n-1} x + (n-1) I_{n-2} - (n-1) I_n.$$

$$\Rightarrow I_n = \frac{-1}{n} \cos x \sin^{n-1} x + \frac{n-1}{n} I_{n-2}.$$

$$\text{Now } I_n = \int_0^{\pi/2} \sin^n x \, dx.$$

$$= \left. \frac{-1}{n} \cos x \sin^{n-1} x \right|_0^{\pi/2} + \frac{n-1}{n} I_{n-2}$$

$$I_n = \frac{n-1}{n} I_{n-2}.$$

$$I_{n-2} = \frac{n-3}{n-2} I_{n-4}.$$

$$I_{n-4} = \frac{n-5}{n-4} I_{n-6}.$$

In general,

$$\int_0^{\pi/2} \sin^n x \, dx = \begin{cases} \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdots \frac{1}{2} I_0 & (n \text{ is even}) \\ \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdots \frac{2}{3} I_1 & (n \text{ is odd}) \end{cases}$$

$$\text{Since } I_n = \int_0^{\pi/2} \sin^n x \, dx.$$

$$I_0 = \int_0^{\pi/2} dx = \pi/2$$

$$I_1 = \int_0^{\pi/2} \sin x \, dx = -\cos x \Big|_0^{\pi/2}$$

$$\therefore I_1 = 1$$

$$\therefore \textcircled{1} \Rightarrow I_n = \int_0^{\pi/2} \sin^n x \, dx.$$

$$= \begin{cases} \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdots \frac{1}{2} \cdot \frac{\pi}{2}, & n \text{ is even} \\ \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdots \frac{2}{3} \cdot 1, & n \text{ is odd} \end{cases}$$

$$\text{Also } \int_0^{\pi/2} \sin^5 x \, dx.$$

$$= \frac{4}{5} \cdot \frac{2}{3} \cdot 1 = \frac{8}{15}$$

① Reduction formula for $\int \cos^n x \, dx$ and hence find $\int_0^{\pi/2} \cos^n x \, dx$, $\int_0^{\pi/2} \cos^9 x \, dx$

Sol Let $I_n = \int \cos^n x \, dx$

Then $I_n = \int \cos^{n-1} x \cos x \, dx.$

$$u = \cos^{n-1} x \quad dv = \cos x \, dx$$

$$du = (n-1) \cos^{n-2} x (-\sin x) \, dx$$

$$v = \sin x.$$

$$\therefore I_n = \sin x \cos^{n-1} x + (n-1) \int \sin^2 x \cos^{n-2} x \, dx.$$

$$= \sin x \cos^{n-1} x + (n-1) \int (1 - \cos^2 x) \cos^{n-2} x \, dx.$$

$$= \sin x \cos^{n-1} x + (n-1) \int (\cos^{n-2} x \, dx - \cos^n x \, dx) \\ = \sin x \cos^{n-1} x + (n-1) I_{n-2} - (n-1) I_n$$

Problems based on trigonometric integrals

① Evaluate $\int \sin^6 x \cos^3 x dx$

Sol Let $I = \int \sin^6 x \cos^3 x dx$.

$$\text{Then } I = \int \sin^6 x \cos^2 x \cos x dx.$$

$$= \int \sin^6 x (1 - \sin^2 x) \cos x dx$$

put $u = \sin x \quad du = \cos x dx$

$$I = \int u^6 (1 - u^2) du = \int (u^6 - u^8) du$$

$$\therefore I = \frac{u^7}{7} - \frac{u^9}{9} + c.$$

$$\therefore I = \frac{\sin^7 x}{7} - \frac{\sin^9 x}{9} + c$$

② Solve $I = \int_0^{\pi/3} \tan^5 x \sec^4 x dx$

Sol $I = \int_0^{\pi/3} \tan^4 x \sec^3 x (\sec x \tan x) dx$

$$= \int_0^{\pi/3} (\sec^2 x - 1)^2 \sec^3 x (\sec x \tan x) dx.$$

let $u = \sec x \quad du = \sec x \tan x dx$

x	0	$\pi/3$
u	1	2

$$\therefore I = \int_1^2 (u^2 - 1)^2 u^3 du$$

$$= \int_1^2 (u^4 - 2u^2 + 1) u^3 du$$

$$= \int_1^2 (u^7 - 2u^5 + u) du.$$

$$= \left[\frac{u^8}{8} - \frac{2u^6}{6} + \frac{u^2}{2} \right]_1^2 = \frac{117}{8}.$$

Expression	Substitution	Identity
$\sqrt{a^2 - x^2}$	$x = a \sin \theta$ $-\pi/2 \leq \theta \leq \pi/2$	$1 - \sin^2 \theta = \cos^2 \theta.$
$\sqrt{a^2 + x^2}$	$x = a \tan \theta$ $-\pi/2 < \theta < \pi/2$	$1 + \tan^2 \theta = \sec^2 \theta.$
$\sqrt{x^2 - a^2}$	$x = a \sec \theta$ $0 \leq \theta < \pi/2, \pi \leq \theta < \frac{3\pi}{2}$	$\sec^2 \theta - 1 = \tan^2 \theta.$

① Evaluate $\int \frac{\sqrt{9-x^2}}{x^2} dx$.

Sol

$$I = \int \frac{\sqrt{9-x^2}}{x^2} dx.$$

put $x = 3 \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2$

$$dx = 3 \cos \theta d\theta$$

$$\therefore I = \int \frac{\sqrt{9-9\sin^2 \theta}}{9\sin^2 \theta} (3\cos \theta) d\theta.$$

$$= \int \frac{3\sqrt{1-\sin^2 \theta}}{9\sin^2 \theta} (3\cos \theta) d\theta.$$

$$= \int \frac{\cos \theta}{\sin^2 \theta} d\theta = \int (\cot \theta - \csc \theta) d\theta$$

$$= -\cot \theta - \theta + c.$$

$$\therefore I = -\frac{\sqrt{9-x^2}}{x} - \sin^{-1}\left(\frac{x}{3}\right) + c.$$

② Evaluate $I = \int \frac{1}{\sqrt{a^2 - x^2}}$

$$x = a \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2$$

$$dx = a \cos \theta d\theta.$$

$$I = \int \frac{1}{\frac{\tan^2 \theta}{\sqrt{a^2 - a^2 \sin^2 \theta}}} a \cos \theta d\theta$$

$$= \int \frac{1}{a \cos \theta} a \cos \theta d\theta$$

$$\therefore I = \int d\theta = \theta + C = \sin^{-1}\left(\frac{x}{a}\right) + C$$

③ Let $I = \int \frac{1}{x^2 \sqrt{x^2 - 1}} dx, x > 1$

put $x = \sec \theta$

$$dx = \sec \theta \tan \theta d\theta$$

$$\therefore I = \int \frac{1}{\sec^2 \theta \tan \theta} \sec \theta \tan \theta d\theta$$

$$= \int \cos \theta d\theta$$

$$= \sin \theta + C$$

$$= \frac{\sqrt{x^2 - 1}}{x} + C$$

Partial Fraction method.

Solve $\int \frac{dx}{(x+1)(x+2)}$

Sol $\frac{1}{(x+1)(x+2)} = \frac{A}{x+1} + \frac{B}{x+2}$

$$\Rightarrow A(x+2) + B(x+1) = 1$$

put $x = -2, -B = 1 \Rightarrow \boxed{B = -1}$

put $x = -1, \boxed{A = 1}$

$$\therefore \int \frac{dx}{(x+1)(x+2)} = \int \frac{dx}{x+1} - \int \frac{dx}{x+2}$$

$$= \log \left(\frac{x+1}{x+2} \right) + C$$

② Solve $\int \frac{x^2 + 1}{(x-3)(x-2)^2} dx$

Sol

$$\frac{x^2 + 1}{(x-3)(x-2)^2} = \frac{A}{x-3} + \frac{B}{x-2} + \frac{C}{(x-2)^2}$$

$$A(x-2)^2 + B(x-2)(x-3) + C(x-3) = x^2 + 1$$

put $x = 3, \boxed{A = 10}$

put $x = 2, \boxed{C = -5}$

equating coefft of x^2 on both sides. $1 = A + B$

$$\therefore \boxed{B = -9}$$

$$\therefore \Rightarrow \frac{x^2 + 1}{(x-3)(x-2)^2} = \frac{10}{x-3} - \frac{9}{x-2} - \frac{5}{(x-2)^2}$$

$$\therefore \int \frac{x^2 + 1}{(x-3)(x-2)^2} dx = 10 \int \frac{dx}{x-3} - 9 \int \frac{dx}{x-2} - 5 \int \frac{dx}{(x-2)^2}$$

$$I = 10 \log(x-3) - 9 \log(x-2) + 5 \left(\frac{1}{x-2} \right) + C$$

③ Solve $\int \frac{10 dx}{(x-1)(x^2+9)}$

Sol

$$\frac{10}{(x-1)(x^2+9)} = \frac{A}{x-1} + \frac{Bx+C}{x^2+9}$$

$$\Rightarrow A(x^2+9) + (Bx+C)(x-1) = 10$$

put $x = 1, \boxed{A = 1}$

put $x = 0, 9A - C = 10$

$$\Rightarrow \boxed{C = -1}$$

Equating the coefficient of x^2 , we get, $A+B=0$
 $\Rightarrow \boxed{B=-1}$

$$\therefore \frac{1}{(x-1)(x^2+9)} = \frac{1}{x-1} + \frac{(-x-1)}{x^2+9}$$

$$\text{Now } \int \frac{1}{(x-1)(x^2+9)} dx = \int \frac{dx}{x-1} - \int \frac{x dx}{x^2+9} - \int \frac{dx}{x^2+9}$$

$$\therefore I = \log(x-1) - \frac{1}{2} \log(x^2+9) - \frac{1}{3} \tan^{-1}\left(\frac{x}{3}\right) + C$$

④ Integrate the following.

$$\int \frac{x^4 - 2x^2 + 4x + 1}{x^3 - x^2 - x + 1} dx$$

Sol Since $\deg(Nr) > \deg(Dr)$, it is improper

$$\text{So, } f(x) = q(x) + \frac{r(x)}{p(x)}$$

$$\begin{array}{r} \text{Q:} \\ x^3 - x^2 - x + 1 \overline{) x^4 - 0x^3 - 2x^2 + 4x + 1} \\ \underline{x^4 - x^3 - x^2 + x} \\ x^3 - x^2 + 3x + 1 \\ \underline{x^3 - x^2 - x + 1} \\ 4x \end{array}$$

$$\therefore \frac{x^4 - 2x^2 + 4x + 1}{x^3 - x^2 - x + 1}$$

$$= (x+1) + \frac{4x}{x^3 - x^2 - x + 1} \quad \text{--- (1)}$$

$$\text{Now } \frac{4x}{x^3 - x^2 - x + 1} = \frac{4x}{(x-1)^2(x+1)}$$

$$= \frac{4x}{(x-1)^2(x+1)} = \frac{A}{(x-1)^2} + \frac{B}{(x-1)} + \frac{C}{x+1}$$

$$\Rightarrow 4x = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

$$\text{put } x=1, \quad \boxed{B=2}$$

$$\text{put } x=-1, \quad \boxed{C=-1}$$

equating the coefft. of x^2 ,

$$A+C=0 \Rightarrow \boxed{A=1}$$

$$\therefore \text{ (1)} \Rightarrow \frac{x^4 - 2x^2 + 4x + 1}{x^3 - x^2 - x + 1}$$

$$= (x+1) + \frac{1}{x-1} + \frac{2}{(x-1)^2} - \frac{1}{x+1}$$

$$\therefore I = \int (x+1) dx + \int \frac{dx}{x-1} + 2 \int \frac{dx}{(x-1)^2} - \int \frac{dx}{x+1}$$

$$I = \frac{x^2}{2} + x + \log(x-1) + 2 \left(\frac{-1}{(x-1)} \right) - \log(x+1) + C$$

$$\therefore I = \frac{x^2}{2} + x - \frac{2}{x-1} + \log\left(\frac{x-1}{x+1}\right) + C$$

The given integral is of the form $\int \frac{dx}{ax^2+bx+c}$

① Evaluate $\int \frac{dx}{3x^2-4x-5}$

Sol

Let $I = \int \frac{dx}{3x^2-4x-5}$

Then $I = \frac{1}{3} \int \frac{dx}{x^2 - \frac{4}{3}x - \frac{5}{3}}$

Now

$$\begin{aligned} x^2 - \frac{4}{3}x - \frac{5}{3} &= x^2 - \frac{4}{3}x + \left(\frac{2}{3}\right)^2 - \left(\frac{2}{3}\right)^2 - \frac{5}{3} \\ &= \left(x - \frac{2}{3}\right)^2 - \frac{19}{9} \\ &= \left(x - \frac{2}{3}\right)^2 - \left(\frac{\sqrt{19}}{3}\right)^2 \end{aligned}$$

$$\therefore I = \frac{1}{3} \int \frac{dx}{\left(x - \frac{2}{3}\right)^2 - \left(\frac{\sqrt{19}}{3}\right)^2}$$

$$\left. \begin{aligned} \therefore \int \frac{dx}{x^2 - a^2} \\ = \frac{1}{2a} \log \left(\frac{x-a}{x+a} \right) + c \end{aligned} \right\}$$

$$\therefore I = \frac{1}{3} \frac{1}{2\left(\frac{\sqrt{19}}{3}\right)} \log \left[\frac{\left(x - \frac{2}{3}\right) - \frac{\sqrt{19}}{3}}{\left(x - \frac{2}{3}\right) + \frac{\sqrt{19}}{3}} \right] + c$$

$$\Rightarrow I = \frac{1}{2\sqrt{19}} \log \left[\frac{3x-2-\sqrt{19}}{3x-2+\sqrt{19}} \right] + c$$

The given integral is of the form $\int \frac{px+q}{ax^2+bx+c}$

$$Nx = A \frac{d}{dx}(Dx) + B.$$

$$\text{i.e.} \int \frac{px+q}{ax^2+bx+c} = \int \frac{A \frac{d}{dx}(ax^2+bx+c)}{ax^2+bx+c}$$

$$+ \int \frac{B}{ax^2+bx+c} dx.$$

① $\int \frac{2x+3}{x^2+2x+5} dx = ?$

Sol Let $2x+3 = A \frac{d}{dx}(x^2+2x+5) + B.$

$$\Rightarrow 2x+3 = A(2x+2) + B$$

Equating the Coefficients of x we get

$$2A = 2 \Rightarrow \boxed{A=1} \text{ \& } \cancel{B=3}$$

$$2A+B=3 \Rightarrow \boxed{B=1}$$

$$\therefore 2x+3 = (2x+2)+1.$$

$$\therefore I = \int \frac{2x+3}{x^2+2x+5} dx.$$

$$= \int \frac{(2x+2)+1}{x^2+2x+5} dx$$

$$= \int \frac{(2x+2)dx}{x^2+2x+5} + \int \frac{dx}{x^2+2x+5}$$

$$= \log(x^2+2x+5) + \int \frac{dx}{x^2+2x+5} \quad \text{②}$$

like $\int \frac{dx}{x^2+2x+5}$

$$\int \frac{dx}{x^2+2x+5} = \int \frac{dx}{x^2+2x+1^2+4}$$

$$= \int \frac{dx}{(x+1)^2+4}$$

$$= \frac{1}{2} \tan^{-1}\left(\frac{x+1}{2}\right) + C.$$

$$\Rightarrow I = \log(x^2+2x+5) + \frac{1}{2} \tan^{-1}\left(\frac{x+1}{2}\right) + C$$

The given integral is of the form

$$\int \frac{px+q}{\sqrt{ax^2+bx+c}} dx$$

$$px+q = A \frac{d}{dx}(ax^2+bx+c) + B$$

Solve $\int \frac{2x-1}{\sqrt{x^2+5x+6}} dx$

or $2x-1 = A \frac{d}{dx}(x^2+5x+6) + B$

$$\Rightarrow 2x-1 = (2x+5)A + B$$

Equating the coefficients of x we get

$$2A = 2 \Rightarrow \boxed{A=1}$$

Also, $-1 = 5A + B$

$$\Rightarrow -1 = 5 + B \Rightarrow \boxed{B=-6}$$

$$\therefore I = \int \frac{2x-1}{\sqrt{x^2+5x+6}} dx$$

$$= \int \frac{(2x+5)}{\sqrt{x^2+5x+6}} dx - 6 \int \frac{dx}{\sqrt{x^2+5x+6}}$$

$$\therefore I = 2\sqrt{x^2+5x+6}$$

$$- 6 \int \frac{dx}{\left(x+\frac{5}{2}\right)^2 + 6 - \frac{25}{4}}$$

$$= 2\sqrt{x^2+5x+6}$$

$$- 6 \int \frac{dx}{\left(x+\frac{5}{2}\right)^2 - \left(\frac{1}{2}\right)^2}$$

$$= 2\sqrt{x^2+5x+6}$$

$$- 6 \log\left[\left(x+\frac{5}{2}\right) + \sqrt{x^2+5x+6}\right] + C$$

The given integral is of the form $\int (px+q)\sqrt{ax^2+bx+c} dx$

put $px+q = A \frac{d}{dx}(ax^2+bx+c) + B$

$$\int (px+q)\sqrt{ax^2+bx+c} dx.$$

$$= A \int \sqrt{ax^2+bx+c} d(ax^2+bx+c)$$

$$+ B \int \sqrt{ax^2+bx+c} dx$$

$$= A \sqrt{u} du + B \sqrt{ax^2+bx+c} dx$$

① Solve $\int (3x-2)\sqrt{x^2+x+1} dx$

or $3x-2 = A \frac{d}{dx}(x^2+x+1) + B$

$$3x-2 = A(2x+1) + B$$

Equating the coefficients of x , we get

$$2A = 3 \Rightarrow \boxed{A = \frac{3}{2}}$$

also, $A+B = -2 \Rightarrow \boxed{B = -\frac{7}{2}}$

$$\begin{aligned}
 \therefore I &= \int (3x-2) \sqrt{x^2+x+1} \, dx. \\
 &= \frac{3}{2} \int \sqrt{x^2+x+1} \, d(x^2+x+1) \\
 &\quad - \frac{7}{2} \int \sqrt{x^2+x+1} \, dx. \\
 &= \frac{3}{2} \frac{(x^2+x+1)^{3/2}}{3/2} \\
 &\quad - \frac{7}{2} \int \sqrt{\left(x+\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2} \, dx. \\
 &= \frac{3}{2} (x^2+x+1)^{3/2} \\
 &\quad - \frac{7}{2} \left[\left(\frac{2x+1}{4}\right) \sqrt{x^2+x+1} \right. \\
 &\quad \left. + \frac{3}{8} \log \left[x + \frac{1}{2} + \sqrt{x^2+x+1} \right] \right] + C
 \end{aligned}$$

Improper Integrals

In a regular (proper) definite integral $\int_a^b f(x) dx$ it is assumed that the limits of integration are finite and that the integrand $f(x)$ is continuous for every value of x in the interval $a \leq x \leq b$.

If atleast one of these conditions is violated, then the integral is known as an improper integral. (Singular or indefinite integrals)

Defn. of an improper integral (Type I)

① If $\int_a^t f(x) dx$ exists for every number $t > a$, then

$$\int_a^\infty f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx.$$

② If $\int_t^b f(x) dx$ exists for every number $t \leq b$, then

$$\int_{-\infty}^b f(x) dx = \lim_{t \rightarrow -\infty} \int_t^b f(x) dx$$

provided this limit exists. (as a finite no.)

③ If both $\int_a^\infty f(x) dx$ and $\int_{-\infty}^a f(x) dx$ are convergent, if the corresponding limit exists and divergent if the limit does not exist.

④ If $\int_a^\infty f(x) dx$ and $\int_{-\infty}^a f(x) dx$ are convergent, then we define

$$\int_{-\infty}^\infty f(x) dx = \int_{-\infty}^a f(x) dx + \int_a^\infty f(x) dx$$

'a' is any real number.

Defn. of Improper Integral

Type: 2

① If f is continuous on $[a, b)$ and is discontinuous at b , then $\int_a^b f(x) dx$

$$= \lim_{t \rightarrow b^-} \int_a^t f(x) dx \text{ If this limit exists.}$$

② If f is continuous on $(a, b]$ and is discontinuous at a , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx$$

If this limit exist

The improper integral $\int_a^b f(x) dx$ is convergent

If the corresponding limit exists and divergent if the limit does not exist.

③ If f has a discontinuity at c , where $a < c < b$ and both $\int_a^c f(x) dx$ and $\int_c^b f(x) dx$ are convergent, then we define

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$

① Determine whether integral $\int_1^\infty \frac{1}{x} dx$ is convergent or divergent

Sol given $\int_1^\infty \frac{1}{x} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x} dx$

$$= \lim_{t \rightarrow \infty} [\log x]_1^t$$

$$= \lim_{t \rightarrow \infty} [\log t - \log 1]$$

$$= \lim_{t \rightarrow \infty} \log t = \infty \text{ (divergent)}$$

$$\therefore \int_1^\infty \frac{1}{x} dx \text{ is divergent}$$

② Determine whether the integral $\int_1^\infty \frac{\log x}{x^2} dx$ is convergent or divergent

Sol Take $\int \frac{\log x}{x^2} dx$

$$u = \log x \quad dv = \frac{1}{x^2} dx$$

$$du = \frac{dx}{x} \quad \Rightarrow v = -\frac{1}{x}$$

$$\therefore \int \frac{\log x}{x^2} dx$$

$$= -\frac{1}{x} \log x + \int \frac{1}{x^2} dx$$

$$= -\frac{1}{x} \log x - \frac{1}{x}$$

$$= -\frac{1}{x} [\log x + 1]$$

$$\therefore \int_1^{\infty} \frac{\log x}{x^2} dx$$

$$= \lim_{t \rightarrow \infty} \int_1^t \frac{\log x}{x^2} dx$$

$$= \lim_{t \rightarrow \infty} \left[\frac{-1}{x} (\log x + 1) \right]_1^t$$

$$= \lim_{t \rightarrow \infty} \left[\frac{-1}{t} (\log t + 1) + \log 1 + 1 \right]$$

$$= \lim_{t \rightarrow \infty} \left(\frac{-\log t}{t} \right) = \left(\frac{\infty}{\infty} \text{ form} \right)$$

Apply L'Hospital rule,

$$\therefore I = \lim_{t \rightarrow \infty} \left(\frac{-\log t}{t} \right)$$

$$I = \lim_{t \rightarrow \infty} \left(\frac{-1/t}{1} \right) = 0$$

$$\therefore I = \int_1^{\infty} \frac{\log x}{x^2} dx = 1 \text{ (finite)}$$

$$\therefore \int_1^{\infty} \frac{\log x}{x^2} \text{ is Convergent.}$$

③ Evaluate $\int_{-\infty}^0 x e^x dx$

Sol $I = \lim_{t \rightarrow -\infty} \int_t^0 x e^x dx.$

$$= \lim_{t \rightarrow -\infty} x d(e^x)$$

$$= \lim_{t \rightarrow -\infty} \left[x e^x \right]_t^0 - \int_t^0 e^x dx$$

$$= \lim_{t \rightarrow -\infty} \left[-t e^t - (1 - e^t) \right]$$

$$= \lim_{t \rightarrow -\infty} \left[-t e^t - 1 + e^t \right] = -1 \text{ (finite)}$$

$$\therefore \int_{-\infty}^0 x e^x dx \text{ is Convergent.}$$

④ For what values of p in the integral $\int_1^{\infty} \frac{1}{x^p} dx$ Convergent?

Sol

If $p \neq 1$, $\lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^p} dx$

$$= \lim_{t \rightarrow \infty} \int_1^t x^{-p} dx.$$

$$= \lim_{t \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^t$$

$$= \lim_{t \rightarrow \infty} \left[\frac{t^{-p+1}}{-p+1} - \frac{1}{-p+1} \right]$$

$$= \lim_{t \rightarrow \infty} \frac{1}{p-1} \left[1 - \frac{1}{t^{p-1}} \right]$$

$$= \frac{1}{p-1}, \quad p > 1, \text{ Converges}$$

$$\infty, \quad p \leq 1, \text{ diverges}$$

⑤ $\int_3^{\infty} \frac{1}{(x-2)^{3/2}} dx = ?$

Sol take $\int \frac{1}{(x-2)^{3/2}} dx$ — ①

put $u = x - 2$; $du = dx$


$$\text{①} \Rightarrow \int \frac{dx}{(x-2)^{3/2}} = \int \frac{du}{u^{3/2}}$$

$$= \int u^{-3/2} du$$

Hydrostatic force and force.

Sea divers realize that water pressure increases as they dive deeper into the sea. This significant feature happens because the weight of the water above them increases.

Let us assume that a thin horizontal plate with area A square meters is submerged in a fluid of density ' ρ ' kilograms per cubic meter at a depth ' d ' meters below the surface of the fluid.



The fluid directly above the plate has volume $V = Ad$.

Then its mass is $m = \rho V = \rho Ad$.

The force exerted by the fluid on the plate

$$F = mg = \rho g Ad.$$

where ' g ' is the acceleration due to gravity.

$\therefore$ The pressure on the plate is

$$P = \frac{F}{A} = \rho g d.$$

The SI unit for measuring pressure is Newton per square meter is called Pascal.

We know that the density of water is given by $\rho = 1000 \text{ kg/m}^3$

The pressure at the bottom of a swimming pool 2 meter 2-meter deep is given by

$$P = \rho g d = 1000 \text{ kg/m}^3 \times 9.8 \text{ m/s}^2 \times 2 \text{ meter.}$$


$$\therefore P = 19,600 \text{ Pa} = 19.6 \text{ kPa}$$

The principle of fluid pressure is that, at any point in a liquid the pressure is the same in all directions. Therefore, the divers has the same pressure on nose and both ears.

Hence the pressure in any direction at a depth "d" in a fluid with mass density " ρ " is given by $P = \rho g d = \delta d$.

This relation helps to determine the hydrostatic force against a vertical plate or wall or dam in a fluid.

- ① A dam has the shape of the trapezoid as shown in figure.



The height is 20m and the width is 50m at the top and 30m at the bottom. Find the force on the dam due hydrostatic pressure if the water level is 4m from the top of the dam.


at the surface of the water and directed down ward.

Then the depth of the water is 16 m.

$\therefore$ the interval $[0, 16]$ can be divided into sub-intervals of equal length with end points x_i , $x_i \in [x_{i-1}, x_i]$

Let us consider the i^{th} horizontal strip of the dam is approximated by a rectangle with height Δx and width w_i

$$\therefore \frac{a}{16-x_i} = \frac{10}{20} \Rightarrow a = \frac{16-x_i}{2} = 8 - \frac{x_i}{2}$$



$$w_i = 2(15+a) = 2(15+8-\frac{x_i}{2})$$

$$\therefore w_i = 46 - x_i$$

If A_i is the area of the i^{th} strip, then

$$A_i = w_i \Delta x = (46 - x_i) \Delta x$$

If Δx is very small, then the pressure P_i on the ~~i^{th}~~ i^{th} strip is almost constant.

$$\therefore P = \rho g d = \rho g x_i$$

$$P_i = w_i \rho g x_i$$

Now the hydrostatic force F_i is

$$F_i = P_i A_i = 1000 g x_i (46 - x_i) \Delta x.$$

As $n \rightarrow \infty$,

$$F = \lim_{n \rightarrow \infty} \sum_{i=1}^n 1000 g x_i (46 - x_i) \Delta x.$$

$$= \int_0^{16} 1000 g x (46 - x) dx$$

$$= 1000 g \int_0^{16} (46x - x^2) dx$$


$$= 1000 (9.8) \left[\frac{46x^2}{2} - \frac{x^3}{3} \right]_0^{16}$$

$$= 9800 \left[23 (16)^2 - \frac{1}{3} (16)^3 \right]$$

$$= 4.43 \times 10^7 \text{ N}.$$

Find the hydrostatic force on one end of a cylindrical drum with radius 3 feet if the drum is submerged in water 10 feet deep.

The circle has a simple equation $x^2 + y^2 = 9$. ①



We divide the Circular region into horizontal strips of equal width.

From (1), the length of the i th strip is given by $2\sqrt{9-(y_i)^2}$

∴ Its area (strip) is $A_i = 2\sqrt{9-(y_i)^2} \Delta y$.

The pressure on the strip is $\delta_i = 62.7(7-y_i)$

∴ the force on the strip is

$$\delta_i A_i = 62.7(7-y_i) 2\sqrt{9-(y_i)^2} \Delta y$$

∴ The total force is

$$F = \lim_{n \rightarrow \infty} \sum_{i=1}^n 62.7(7-y_i) 2\sqrt{9-(y_i)^2} \Delta y$$

$$= 125 \int_{-3}^3 (7-y) \sqrt{9-y^2} dy$$

$$= 125 \times 7 \times \int_{-3}^3 \sqrt{9-y^2} dy - 125 \int_{-3}^3 y \sqrt{9-y^2} dy$$

$$= 875 \int_{-3}^3 \sqrt{9-y^2} dy - 0$$

$$= 875 \int_{-3}^3 \sqrt{3^2-y^2} dy$$

$$= 875 \left[\frac{1}{2} y \sqrt{9-y^2} \right. \\ \left. + \frac{1}{2} 3^2 \sin^{-1}(y/3) \right]_{-3}^3$$


$$= 875 \left[\left(\frac{3}{2}(0) + \frac{9}{2} \sin^{-1}(1) \right) - \left(-\frac{3}{2}(0) - \frac{9}{2} \sin^{-1}(-1) \right) \right]$$

$$= 875 [9 \sin^{-1}(1)]$$

$$F = 12,375 \text{ lb}$$

$$\left\{ \begin{aligned} \int \sqrt{a^2-x^2} dx \\ = \frac{1}{2} x \sqrt{a^2-x^2} \\ + \frac{1}{2} a^2 \sin^{-1}(x/a) \end{aligned} \right\}$$

where m_1x_1 & m_2x_2 are moments of m_1 & m_2 and $m = m_1 + m_2$



Suppose we've 'n' particles with corresponding masses $m_1, m_2, m_3, \dots, m_n$ located at the points $x_1, x_2, \dots, x_n$ on the x-axis, then the centre of mass of the entire system is

$$\bar{x} = \frac{\sum_{i=1}^n m_i x_i}{\sum_{i=1}^n m_i} = \frac{\sum_{i=1}^n m_i x_i}{m}$$

where $m = \sum m_i$ is the total mass of the system and the sum of the individual moments

$M = \sum_{i=1}^n m_i x_i$ is called the moment of the system about the origin.

$$\therefore M = m\bar{x}.$$

Let us consider a system of n particles with masses $m_1, m_2, m_3, \dots, m_n$ located at $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ in the xy-plane.

$\therefore$ the moment of the system about y-axis is given by $M_y = \sum_{i=1}^n m_i x_i$, $M_x = \sum_{i=1}^n m_i y_i$

Then M_y measures the tendency of the system to rotate about the y-axis and M_x measures the tendency to rotate about the x-axis.

$\therefore (\bar{x}, \bar{y})$ of the centre of mass are given by

$$\bar{x} = \frac{M_y}{m} \quad \text{and} \quad \bar{y} = \frac{M_x}{m}$$

Where $m = \sum_{i=1}^n m_i$ is the total mass.

Since $m\bar{x} = M_y$ & $m\bar{y} = M_x$, the centre of mass $(\bar{x}, \bar{y})$ is the point where a single particle of mass m will have the same moments.

Centroid and Symmetry

Consider a flat plate which is called as a lamina with uniform density ρ that occupies a region R of the plane. We wish to locate the centre of mass of the plate which is called as the centroid of the region R .

If R is symmetric about a line l , then the centroid of R lies on l is called principle of symmetry. Hence the centroid of a rectangle is its centre.

Moments should be defined so that if the entire mass of a region is concentrated at the centre of mass, then its moments remains unchanged. The moments of the union of two non-overlapping regions should be the sum of the moments of the individual regions.

The moment of the region R above y -axis is given by

$$M_y = \lim_{n \rightarrow \infty} \sum_{i=1}^n \rho \bar{x}_i f(\bar{x}_i) \Delta x = \rho \int_a^b x f(x) dx$$

① Find the centre of mass of a semicircular plate of radius r .

Sol

Equation of the circle

$$\text{is } x^2 + y^2 = r^2 \Rightarrow y^2 = r^2 - x^2$$

$$\therefore y = f(x) = \sqrt{r^2 - x^2} \text{ and}$$

$$a = -r, b = r$$

Since the centre of mass must lie on the y -axis, $\bar{x} = 0$


$$\therefore \text{Area } A = \frac{1}{2} \pi r^2$$

$$\text{Now } \bar{y} = \frac{1}{A} \int_{-r}^r \frac{1}{2} [f(x)]^2 dx = \frac{1}{\frac{1}{2} \pi r^2} \int_{-r}^r \frac{1}{2} [r^2 - x^2] dx$$

$$\therefore \bar{y} = \frac{2}{\pi r^2} \int_0^r (r^2 - x^2) dx = \frac{2}{\pi r^2} \left[r^2 x - \frac{x^3}{3} \right]_0^r$$

$$\bar{y} = \frac{2}{\pi r^2} \left[r^3 - \frac{r^3}{3} \right] \Rightarrow \bar{y} = \frac{4r}{3\pi}$$

$$\therefore (\bar{x}, \bar{y}) = \left(0, \frac{4r}{3\pi} \right)$$




② Find the centroid of the region bounded by the curves $x=0$, $x=\pi/2$, $y=0$ and $y=\cos x$.

Sol

The area of the region

$$A = \int_0^{\pi/2} \cos x \, dx$$

$$= [\sin x]_0^{\pi/2} \Rightarrow \boxed{A=1}$$



We know that the centroid of the region is given by $(\bar{x}, \bar{y})$, where

$$\bar{x} = \frac{1}{A} \int_a^b x f(x) dx \quad \& \quad \bar{y} = \frac{1}{A} \int_a^b \frac{1}{2} [f(x)]^2 dx$$

The moment of the region R about the x -axis is

$$M_x = \lim_{n \rightarrow \infty} \sum_{i=1}^n \rho \cdot \frac{1}{2} [f(x_i)]^2 \Delta x = \rho \int_a^b \frac{1}{2} [f(x)]^2 dx$$

For system of particles, the centre of mass of the plate is defined as $m\bar{x} = M_y$ & $m\bar{y} = M_x$.

The mass of the plate is the product of its density and area.

$$m = \rho A = \rho \int_a^b f(x) dx.$$

Then

$$\bar{x} = \frac{M_y}{m} = \frac{\rho \int_a^b x f(x) dx}{\rho \int_a^b f(x) dx} = \frac{\int_a^b x f(x) dx}{\int_a^b f(x) dx}.$$

$$\bar{y} = \frac{M_x}{m} = \frac{\int_a^b \frac{1}{2} [f(x)]^2 dx}{\int_a^b f(x) dx}.$$

$\therefore$ The centre of mass of the plate or the centroid of R is located at the point $(\bar{x}, \bar{y})$ where $\bar{x} = \frac{1}{A} \int_a^b x f(x) dx$

and

$$\bar{y} = \frac{1}{A} \int_a^b \frac{1}{2} [f(x)]^2 dx.$$

We know that $\bar{x} = \frac{1}{A} \int_a^b x [f(x) - g(x)] dx.$

$$\bar{x} = \frac{1}{(1/6)} \int_0^1 x(x - x^2) dx$$

$$\bar{x} = 6 \int_0^1 (x^2 - x^3) dx = 6 \left[\frac{x^3}{3} - \frac{x^4}{4} \right]_0^1$$

$$\boxed{\bar{x} = \frac{1}{2}}$$

$$\bar{y} = \frac{1}{A} \int_a^b \frac{1}{2} [(f(x))^2 - (g(x))^2] dx.$$

$$= \frac{1}{46} \int_0^1 \frac{1}{2} [x^2 - x^4] dx$$

$$= \frac{1}{3} \left[\frac{x^3}{3} - \frac{x^5}{5} \right]_0^1$$

$$\boxed{\bar{y} = \frac{2}{5}}$$

$\therefore$ the centroid is $(\frac{1}{2}, \frac{2}{5})$

$$\text{Now } \bar{x} = \frac{1}{A} \int_a^b x f(x) dx = \int_0^{\pi/2} x \cos x dx.$$

$$u = x \quad u' = 1 \quad u'' = 0.$$

$$v = \cos x \quad v_1 = \sin x \quad v_2 = -\cos x.$$

$$\therefore \bar{x} = \left[x \sin x - \int \sin x dx \right]_0^{\pi/2} = \left(\frac{\pi}{2} + 0 \right) - (0 - 1)$$

$$\therefore \bar{x} = \frac{\pi}{2} - 1$$

$$\bar{y} = \frac{1}{A} \int_a^b \frac{1}{2} [f(x)]^2 dx = \frac{1}{2} \int_0^{\pi/2} \cos^2 x dx$$

$$\bar{y} = \frac{1}{4} \int_0^{\pi/2} (1 + \cos 2x) dx = \frac{1}{4} \left[x + \frac{\sin 2x}{2} \right]_0^{\pi/2}$$

$$\bar{y} = \frac{\pi}{8}$$

$\therefore$ the centroid is $(\bar{x}, \bar{y}) : \left(\frac{\pi}{2} - 1, \frac{\pi}{8} \right)$

③ Note

If the region R lies between $y=f(x)$ and $y=g(x)$ where $f(x) \geq g(x)$ then the centroid of R is $(\bar{x}, \bar{y})$, where $\bar{x} = \frac{1}{A} \int_a^b x[f(x) - g(x)] dx$ and $\bar{y} = \frac{1}{A} \int_a^b \frac{1}{2} [f(x)^2 - g(x)^2] dx$.

③ Find the centroid of the region bounded by the line $y=x$ and the parabola $y=x^2$.


Sol

$$\text{let } f(x) = x \text{ \& } g(x) = x^2$$

$$a=0 \text{ \& } b=1$$

$$\text{Now Area } A = \int_0^1 (x - x^2) dx$$

$$= \left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 \Rightarrow \boxed{A = \frac{1}{6}}$$



Multiple Integrals

Double integrals - change of order of integration -

Double integrals in Polar co-ordinates - Area enclosed

by plane curves - Triple integrals - volume of solids

change of variables in double and triple integrals -

Applications: Moments and centre of mass, moment of inertia.

Double integrals in cartesian co-ordinates:

Type: 1 (Limits are constants)

1) Evaluate $\int_0^a \int_0^b xy(x-y) dx dy$

Answer:

$$\begin{aligned}\int_0^a \int_0^b xy(x-y) dx dy &= \int_0^a \int_0^b (x^2y - xy^2) dx dy \\&= \int_0^a \left(\frac{x^3y}{3} - \frac{x^2y^2}{2} \right) \Big|_0^b dy \\&= \int_0^a \left(\frac{b^3y}{3} - \frac{b^2y^2}{2} \right) dy \\&= \int_0^a b^2 \left[\frac{by}{3} - \frac{y^2}{2} \right] dy \\&= b^2 \left\{ \frac{by^2}{6} - \frac{y^3}{6} \right\} \Big|_0^a \\&= b^2 \left\{ \frac{a^2b}{6} - \frac{a^3}{6} \right\} = \frac{a^2b^2}{6} (b-a)\end{aligned}$$

2) Evaluate $\int_0^1 \int_1^2 x(x+y) dy dx$

Answer: $\int_0^1 \int_1^2 x(x+y) dy dx = \int_0^1 \int_1^2 (x^2 + xy) dy dx$

$$= \int_0^1 \left(x^2y + \frac{xy^2}{2}\right)_1^2 dx$$

$$= \int_0^1 (2x^2 + 2x) - \left(x^2 + \frac{x}{2}\right) dx$$

$$= \int_0^1 \left(x^2 + \frac{3x}{2}\right) dx$$

$$= \left(\frac{x^3}{3} + \frac{3x^2}{4}\right)_0^1 = \left(\frac{1}{3} + \frac{3}{4}\right) - (0+0)$$

$$= \frac{13}{12}$$

3) Evaluate $\int_2^a \int_2^b \frac{dx dy}{xy}$

Ans:

$$\int_2^a \int_2^b \frac{dx dy}{xy} = \int_2^a \frac{dx}{x} \int_2^b \frac{dy}{y}$$

$$= (\log x)_2^a (\log y)_2^b$$

$$= (\log a - \log 2) (\log b - \log 2)$$

$$= \log \frac{a}{2} \log \frac{b}{2}$$

4) Evaluate $\int_0^3 \int_0^2 e^{x+y} dy dx$

Ans: $\int_0^3 \int_0^2 e^{x+y} dy dx = \int_0^3 e^x dx \int_0^2 e^y dy$

$$= (e^x)_0^3 (e^y)_0^2$$

$$= (e^3 - e^0) (e^2 - e^0) = (e^3 - 1)(e^2 - 1)$$

5) Evaluate $\int_1^2 \int_1^3 xy^2 dx dy$

Ans: $\int_1^2 \int_1^3 xy^2 dx dy = \int_1^2 x dx \int_1^3 y^2 dy$

$$= \left(\frac{x^2}{2}\right)_1^2 \left(\frac{y^3}{3}\right)_1^3$$

$$= \left(\frac{4}{2} - \frac{1}{2}\right) \left(\frac{27}{3} - \frac{1}{3}\right) = \frac{3}{2} \cdot \frac{26}{3} = 13.$$

Type-II (Limits are variable)

1) Evaluate $\int_0^1 \int_0^{1-x} y \, dy \, dx$

Ans:

$$\int_0^1 \int_0^{1-x} y \, dy \, dx = \int_0^1 \int_0^{1-x} y \, dx \, dy \quad (\text{correct form})$$

$$= \int_0^1 \left(\frac{y^2}{2} \right)_0^{1-x} dx$$

$$= \int_0^1 \frac{(1-x)^2}{2} dx = \frac{1}{2} \left\{ \frac{(1-x)^3}{-3} \right\}_0^1$$

$$= -\frac{1}{6} (0-1) = \frac{1}{6}$$

2) Evaluate $\int_0^1 \int_x^{\sqrt{x}} xy(x+y) \, dx \, dy$

$$\text{Ans: } \int_0^1 \int_x^{\sqrt{x}} xy(x+y) \, dx \, dy = \int_0^1 \int_x^{\sqrt{x}} (x^2y + xy^2) \, dy \, dx \quad (\text{correct form})$$

$$= \int_0^1 \int_x^{\sqrt{x}} (x^2y + xy^2) \, dy \, dx$$

$$= \int_0^1 \left(\frac{x^2y^2}{2} + \frac{xy^3}{3} \right)_x^{\sqrt{x}} dx$$

$$= \int_0^1 \left[\frac{x^2(\sqrt{x})^2}{2} - \frac{x(\sqrt{x})^3}{3} \right] - \left[\frac{x^2 \cdot x^2}{2} + \frac{x \cdot x^3}{3} \right] dx$$

$$= \int_0^1 \frac{x^3}{2} - \frac{x^{5/2}}{3} - \frac{x^4}{2} - \frac{x^4}{3} dx$$

$$= \left[\frac{x^4}{8} + \frac{x^{7/2}}{3(7/2)} - \frac{x^5}{10} - \frac{x^5}{15} \right]_0^1$$

$$= \frac{1}{8} + \frac{2}{81} - \frac{1}{10} - \frac{1}{15} = \frac{3}{56}$$

3) Evaluate $\int_0^1 \int_0^{x^2} e^{y/x} \, dy \, dx$

Ans: $\int_0^4 \int_0^{x^2} e^{y/x} dy dx$ (correct form)

$$= \int_0^4 \left\{ \frac{e^{y/x}}{1/x} \right\}_0^{x^2} dx$$

$$= \int_0^4 x \{ e^{x^2/x} - e^0 \} dx$$

$$= \int_0^4 x (e^x - 1) dx$$

$$= \int_0^4 (xe^x - x) dx$$

$$= \left\{ xe^x - e^x - \frac{x^2}{2} \right\}_0^4$$

$$= (4e^4 - e^4 - 8) - (-1) = 3e^4 - 7$$

4) Evaluate $\int_0^1 \int_y^{y^2+1} x^2 y dx dy$

Ans:

$$\int_0^1 \int_y^{y^2+1} x^2 y dx dy = \int_0^1 y \left(\frac{x^3}{3} \right)_y^{y^2+1} dy$$

$$= \frac{1}{3} \int_0^1 y \{ (y^2+1)^3 - y^3 \} dy$$

$$= \frac{1}{3} \int_0^1 y \{ y^6 + 1 + 3y^4 + 3y^2 - y^3 \} dy$$

$$= \frac{1}{3} \left\{ \frac{y^8}{8} + \frac{y^2}{2} + \frac{3y^6}{6} + \frac{3y^4}{4} - \frac{y^5}{5} \right\}_0^1$$

$$= \frac{1}{3} \left\{ \frac{1}{8} + \frac{1}{2} + \frac{1}{2} + \frac{3}{4} - \frac{1}{5} - 0 \right\}$$

$$= \frac{1}{3} \left\{ \frac{1}{8} + 1 + \frac{3}{4} - \frac{1}{5} \right\}$$

$$= \frac{1}{3} \left(\frac{67}{40} \right)$$

$$= \frac{67}{120}$$

Sketch the region of integration:

1) sketch the region of integration $\int_0^a \int_{\sqrt{ax-x^2}}^{\sqrt{a^2-x^2}} dx dy$

Answer:

$$\int_0^a \int_{\sqrt{ax-x^2}}^{\sqrt{a^2-x^2}} dx dy = \int_0^a \int_{\sqrt{ax-x^2}}^{\sqrt{a^2-x^2}} dy dx \quad (\text{correct form})$$

$$y: \sqrt{ax-x^2} \rightarrow y: \sqrt{a^2-x^2}$$

$$y = \sqrt{ax-x^2} \rightarrow y = \sqrt{a^2-x^2}$$

$$y^2 = ax-x^2 \rightarrow y^2 = a^2-x^2$$

$$x^2+y^2-ax=0 \rightarrow y^2+x^2=a^2$$


$$x^2+y^2-ax+\frac{a^2}{4}-\frac{a^2}{4}=0 \rightarrow x^2+y^2=a^2$$

$$(x-\frac{a}{2})^2+(y-0)^2=(\frac{a}{2})^2 \rightarrow x^2+y^2=a^2$$

Circle, $C = (a/2, 0)$ $\rightarrow$ Circle, $C = (0, 0)$
radi = $a/2$ $\rightarrow$ radi = a

Also $x: 0 \rightarrow a$
 $x=0 \rightarrow x=a$

Rough
 $2b = -a$
 $b = -a/2$
 $b^2 = \frac{a^2}{4}$



2) Sketch the region of integration for $\int_0^1 \int_0^x f(x,y) dy dx$

Ans:


Given $y: 0 \rightarrow x$

$y=0 \rightarrow y=x$

Also $x: 0 \rightarrow 1$

$x=0 \rightarrow x=1$

N/D'15
M/3/16
M/D 18



3) Sketch the region $\int_0^a \int_0^{\sqrt{a^2-x^2}} dx dy$

Ans: $\int_0^a \int_0^{\sqrt{a^2-x^2}} dx dy = \int_0^a \int_0^{\sqrt{a^2-x^2}} dy dx$ [Correct form]

Given $y: 0 \rightarrow \sqrt{a^2-x^2}$


$y=0 \rightarrow y=\sqrt{a^2-x^2}$

$y^2 = a^2 - x^2$

$x^2 + y^2 = a^2$ circle, $(0,0)$ & $rad = a$

Also $x: 0 \rightarrow a$

$x=0 \rightarrow x=a$



Change of order of integration:

When we change the order of integration the limits are also changed. The following points are very useful when the change of order of integration.

1) If the limits of the inner integral is a function of x . (or function of y) then the first integration should be wrt y (or wrt x).


2) Draw the region of integration by using the given limits.

3) If the integration is first wrt x keeping y as a constant then consider the horizontal strip and vice versa.


4) Then change into vertical strip, to find the limits of x & y .

1) change the order of integration in $\int_0^1 \int_0^x f(x,y) dy dx$

Ans: Given $y: 0 \rightarrow x$
 $y=0 \rightarrow y=x$
 Also $x: 0 \rightarrow 1$
 $x=0 \rightarrow x=1$



Consider the strip is $\parallel$ to Y axis.



To change the strip is $\parallel$ to X axis

$$x=y \rightarrow x=1$$

$$y=0 \rightarrow y=1$$


$$\therefore \int_0^1 \int_0^x f(x,y) dy dx = \int_0^1 \int_y^1 f(x,y) dx dy$$

2) change the order of integration in $\int_0^\infty \int_x^\infty \frac{e^{-y}}{y} dy dx$ & hence evaluate it.


Ans: $\int_0^\infty \int_x^\infty \frac{e^{-y}}{y} dy dx$ (correct form)

Given $y: x \rightarrow \infty$
 $y=x \rightarrow y=\infty$

Also $x: 0 \rightarrow \infty$
 $x=0 \rightarrow x=\infty$



Consider the strip is || to y axis



to change the strip is || to x axis

$$\begin{aligned}
 \text{Hence } \int_0^{\infty} \int_0^{\infty} \frac{e^{-y}}{y} dy dx &= \int_0^{\infty} \int_0^y \frac{e^{-y}}{y} dx dy \\
 &= \int_0^{\infty} \frac{e^{-y}}{y} (x)_0^y dy \\
 &= \int_0^{\infty} \frac{e^{-y}}{y} y dy = \int_0^{\infty} e^{-y} dy \\
 &= -1 [e^{-\infty} - e^0] \\
 &= -1(0 - 1) = 1
 \end{aligned}$$

3) change the order of integration & hence evaluate

$$\int_0^a \int_y^a \frac{x}{x^2 + y^2} dx dy.$$

Jan'2014
m/j 2016
A/m 2018

Ans

$$\text{Given } \int_0^a \int_y^a \frac{x}{x^2 + y^2} dx dy$$



The region is bounded by $x=y, x=a, y=0$ & $y=a$

$$x: y \rightarrow a$$

$$x=y \rightarrow x=a$$

$$y: 0 \rightarrow a$$

$$y \rightarrow y=a$$



$$\begin{aligned}
 \therefore \int_0^a \int_y^a \frac{x}{x^2+y^2} dx dy &= \int_0^a \int_0^x \frac{x}{x^2+y^2} dy dx \\
 &= \int_0^a x \left(\frac{1}{x} \tan^{-1} \frac{y}{x} \right)_0^x dx \\
 &= \int_0^a (\tan^{-1} \frac{y}{x})_0^x dx \\
 &= \int_0^a \tan^{-1}(1) - \tan^{-1}(0) dx \\
 &= \int_0^a \left(\frac{\pi}{4} - 0 \right) dx = \frac{\pi}{4} (x)_0^a = \frac{a\pi}{4}
 \end{aligned}$$

4) change the order of integration & hence evaluate it

$$\int_0^{4a} \int_{x^2/4a}^{2\sqrt{ax}} xy dy dx$$

Given $\int_0^{4a} \int_{x^2/4a}^{2\sqrt{ax}} xy dy dx$

$$y: x^2/4a \rightarrow 2\sqrt{ax}$$

$$y = x^2/4a \rightarrow y = 2\sqrt{ax}$$

$$x^2 = 4ay \rightarrow y^2 = 4ax$$



(Parabola's)

Also $x: 0 \rightarrow 4a$

$$x=0 \rightarrow x=4a$$

Scanned with CamScanner

(10)



$$x: \frac{y^2}{4a} \rightarrow 2\sqrt{ay}$$

$$y: 0 \rightarrow 4a$$

$$\begin{aligned} \therefore \int_0^{4a} \int_{\frac{y^2}{4a}}^{2\sqrt{ay}} xy \, dx \, dy &= \int_0^{4a} \int_{\frac{y^2}{4a}}^{2\sqrt{ay}} xy \, dx \, dy \\ &= \int_0^{4a} y \left(\frac{x^2}{2} \right)_{\frac{y^2}{4a}}^{2\sqrt{ay}} dy \\ &= \frac{1}{2} \int_0^{4a} y \left\{ (2\sqrt{ay})^2 - \left(\frac{y^2}{4a} \right)^2 \right\} dy \\ &= \frac{1}{2} \int_0^{4a} y \left\{ 4ay - \frac{y^4}{16a^2} \right\} dy \\ &= \frac{1}{2} \int_0^{4a} 4ay^2 - \frac{y^5}{16a^2} dy \\ &= \frac{1}{2} \left\{ 4a \frac{y^3}{3} - \frac{y^6}{6 \times 16a^2} \right\}_0^{4a} \\ &= \frac{1}{2} \left\{ (4a)^4 \right\} \left\{ \frac{1}{3} - \frac{(4a)^2}{96a^2} \right\} \\ &= \frac{256a^4}{2} \left(\frac{1}{3} - \frac{1}{6} \right) \\ &= 128a^4 \left(\frac{1}{6} \right) = \frac{64}{3} a^4 \end{aligned}$$

5 change the order of integration & hence evaluate

$$\int_0^a \int_{x^2/a}^{2a-x} xy \, dx \, dy$$

Ans: Given $\int_0^a \int_{x^2/a}^{2a-x} xy \, dy \, dx$ (Correct form)



$$y: x^2/a \rightarrow 2a-x$$

$$y = x^2/a \rightarrow y = 2a-x$$

$$x^2 = ay \rightarrow y = 2a-x$$

Also $x: 0 \rightarrow a$

$$x=0 \rightarrow x=a$$



Consider I_1

$$x: 0 \rightarrow \sqrt{ay}$$

$$y: 0 \rightarrow a$$

$$\therefore I_1 = \int_0^a \int_0^{\sqrt{ay}} xy \, dx \, dy$$

$$= \int_0^a y \left\{ \frac{x^2}{2} \right\}_0^{\sqrt{ay}} dy$$

$$= \int_0^a y \frac{(\sqrt{ay})^2}{2} dy = \frac{1}{2} \int_0^a y (ay) dy$$

$$\begin{aligned}
 &= \frac{a}{2} \int_0^a y^2 dy \\
 &= \frac{a}{2} \left(\frac{y^3}{3} \right)_0^a \\
 &= \frac{a}{2} \left(\frac{a^3}{3} \right) = \frac{a^4}{6}
 \end{aligned}$$

$$\begin{aligned}
 I_2 &= \int_a^{2a} \int_0^{2a-y} xy \, dx \, dy \\
 &= \int_a^{2a} \left[\frac{x^2}{2} \right]_0^{2a-y} dy \\
 &= \frac{1}{2} \int_a^{2a} y \{ 4a^2 - 4ay + y^2 \} dy \\
 &= \frac{1}{2} \left\{ 4 \frac{a^2 y^2}{2} - \frac{4ay^3}{3} + \frac{y^4}{4} \right\}_a^{2a} \\
 &= \frac{1}{2} \left\{ \left[2a^2(2a)^2 - \frac{4a}{3}(2a)^3 + \frac{(2a)^4}{4} \right] - \left[2a^2(a)^2 - \frac{4a}{3}(a)^3 + \frac{a^4}{4} \right] \right\} \\
 &= \frac{a^4}{2} \left\{ 8 - \frac{32}{3} + 4 - 2 + \frac{4}{3} - \frac{1}{4} \right\} \\
 &= \frac{a^4}{2} \left\{ 10 - \frac{28}{3} - \frac{1}{4} \right\} = \frac{5a^4}{24}
 \end{aligned}$$

$$\begin{aligned}
 \therefore I &= I_1 + I_2 = \frac{a^4}{6} + \frac{5a^4}{24} = a^4 \left\{ \frac{1}{6} + \frac{5}{24} \right\} \\
 &= \frac{9a^4}{24} = \frac{3a^4}{8}
 \end{aligned}$$

Double integration in polar co-ordinates

Type: I (limits are constants).

1) Evaluate $\int_0^{\pi/2} \int_0^2 r \, dr \, d\theta$

Ans $\int_0^{\pi/2} \int_0^2 r \, dr \, d\theta = \int_0^{\pi/2} \left(\frac{r^2}{2} \right)_0^2 d\theta$

$$= \frac{1}{2} \int_0^{\pi/2} (4-0) d\theta$$

$$= \frac{4}{2} \left\{ \theta \right\}_0^{\pi/2}$$

$$= 2 \left\{ \pi/2 \right\} = \pi$$

2) Evaluate $\int_0^{\pi/2} \int_0^{\infty} \frac{r}{(r^2+a^2)^2} dr d\theta$

$$I = \int_0^{\pi/2} \int_0^{\infty} r (r^2+a^2)^{-2} dr d\theta$$

$$= \int_0^{\pi/2} \int_0^{\infty} \frac{1}{2} 2r (r^2+a^2)^{-2} dr d\theta$$

$$= \frac{1}{2} \int_0^{\pi/2} \left[\frac{(r^2+a^2)^{-2+1}}{-2+1} \right]_0^{\infty} d\theta$$

$$= -\frac{1}{2} \int_0^{\pi/2} \left[\frac{1}{r^2+a^2} \right]_0^{\infty} d\theta$$

$$= -\frac{1}{2} \int_0^{\pi/2} \left(0 - \frac{1}{a^2} \right) d\theta$$

$$= \frac{1}{2a^2} (\theta)_0^{\pi/2} = \frac{1}{2a^2} \pi/2 = \frac{\pi}{4a^2}$$

Formula

$$\int [f(x)]^n df(x) dx = \frac{[f(x)]^{n+1}}{n+1}$$

Type: 2 (Limits are variables)

1) Evaluate $\int_0^{\pi} \int_0^{\sin \theta} r dr d\theta$

Ans

Given $\int_0^{\pi} \int_0^{\sin \theta} r dr d\theta = \int_0^{\pi} \left(\frac{r^2}{2} \right)_0^{\sin \theta} d\theta$

$$= \frac{1}{2} \int_0^{\pi} \sin^2 \theta d\theta$$

$$= \frac{1}{2} \int_0^{\pi} \left(\frac{1 - \cos 2\theta}{2} \right) d\theta$$

$$= \frac{1}{4} \left\{ \theta - \frac{\sin 2\theta}{2} \right\}_0^{\pi} = \frac{1}{4} \{ (\pi - 0) - (0 - 0) \} = \frac{\pi}{4}$$

2) Evaluate $\int_0^{\pi/2} \int_0^{\sin \theta} r \, dr \, d\theta$

Answer:

$$I = \int_0^{\pi/2} \left\{ \frac{r^2}{2} \right\}_0^{\sin \theta} d\theta$$

$$= \frac{1}{2} \int_0^{\pi/2} \sin^2 \theta \, d\theta$$

$$= \frac{1}{2} \int_0^{\pi/2} \left(1 - \frac{\cos 2\theta}{2} \right) d\theta$$

$$= \frac{1}{4} \left\{ \theta - \frac{\sin 2\theta}{2} \right\}_0^{\pi/2}$$


$$= \frac{1}{4} \{ (\pi/2 - 0) - (0 - 0) \}$$

$$= \pi/8$$

Area Enclosed by plane curves:

Formula $A = \iint_R dx \, dy$

1) Find the area bounded by the lines $x=0$, $y=1$ & $y=x$



Consider the horizontal Strip

$$x: 0 \rightarrow y$$

$$y: 0 \rightarrow 1$$

$$\therefore \text{Area} = \iint_R dx \, dy$$

$$= \int_0^1 \int_0^y dx \, dy$$

$$= \int_0^1 \left\{ x \right\}_0^y dy$$


$$= \int_0^1 y \, dy$$

$$= \left\{ \frac{y^2}{2} \right\}_0^1 = \frac{1}{2}$$

Scanned with CamScanner

2) Find the area of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

Ans:



N/D 2018
A/M 2020

Consider the horizontal strip
Strip starts from ellipse y-axis
 $\therefore$ lower lt of $x=0$
Strip ends at ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\Rightarrow \frac{x^2}{a^2} = 1 - \frac{y^2}{b^2} \quad \therefore x: 0 \rightarrow \frac{a}{b} \sqrt{b^2 - y^2}$$

$$\Rightarrow x = \frac{a}{b} \sqrt{b^2 - y^2}$$

Also $y: 0 \rightarrow b$

$\therefore$ Area = 4 x Area in the I quadrant

$$= 4 \iint_R dx dy$$

$$= 4 \int_0^b \int_0^{\frac{a}{b} \sqrt{b^2 - y^2}} dx dy = 4 \int_0^b (x)_0^{\frac{a}{b} \sqrt{b^2 - y^2}} dy$$

$$= 4 \int_0^b \frac{a}{b} \sqrt{b^2 - y^2} dy$$

$$= \frac{4a}{b} \left\{ \frac{y}{2} \sqrt{b^2 - y^2} + \frac{b^2}{2} \sin^{-1} \frac{y}{b} \right\}_0^b$$

$$= \frac{4a}{b} \left\{ \frac{b^2}{2} \sin^{-1}(1) - \frac{b^2}{2} \sin^{-1}(0) \right\}$$

$$= \frac{4a}{b} \cdot \frac{b^2}{2} \cdot \frac{\pi}{2} = \pi ab$$

3) show that the area b/w the parabolas

$$y^2 = 4ax \text{ \& } x^2 = 4ay \text{ is } \frac{16a^2}{3}$$


Ans

Given $y^2 = 4ax$

$$y = x^2/4a$$

x	-2a	-a	0	a	2a	3a	4a
y	-	-	0	2a	-	-	4a

x	-2a	-a	0	a	2a	3a	4a
y	-	-	0	-	-	-	4a



Consider the horizontal strip.

Strip starts from $y^2 = 4ax$
 $\Rightarrow x = y^2/4a$

strip ends at $x^2 = 4ay$
 $x = 2\sqrt{ay}$

$$\therefore x: \frac{y^2}{4a} \rightarrow 2\sqrt{ay}$$

$$y: 0 \rightarrow 4a$$

$$\therefore \text{Area} = \int_0^{4a} \int_{y^2/4a}^{2\sqrt{ay}} dx dy$$

$$= \int_0^{4a} \left\{ x \right\}_{y^2/4a}^{2\sqrt{ay}} dy$$

$$= \int_0^{4a} \left\{ 2\sqrt{ay} - \frac{y^2}{4a} \right\} dy$$

$$= 2\sqrt{a} \frac{y^{3/2}}{3/2} - \frac{y^3}{4a \times 3} \Big|_0^{4a}$$

$$= \frac{4\sqrt{a}}{3} (4a)^{3/2} - \frac{(4a)^3}{12a}$$

$$= \frac{4\sqrt{a} \cdot 4a(4a)^{1/2}}{3} - \frac{(4a)(4a)(4a)}{12a}$$

$$= \frac{32a^2}{3} - \frac{16a^2}{3} = \frac{16a^2}{3}$$

4) Using double integration, find the area enclosed by the curves $y = 2x^2$ & $y^2 = 4x$

Ans

Find the intersection points

$$y = 2x^2 \rightarrow ①$$

$$y^2 = 4x \rightarrow ②$$

From Q 14

$$4x = (2x^2)^2$$


$$4x^4 - 4x = 0$$

$$x \{ x^3 - 1 \} = 0$$

$$x = 0, x = 1$$

$$\text{When } x=0 \Rightarrow y=0$$

$$x=1 \Rightarrow y=2$$



Consider the horizontal strip.

Here

$$x: \sqrt{y/2} \rightarrow y$$

$$dy: 0 \rightarrow 2$$

$$\therefore \text{Area} = \iint dx dy$$

$$= \int_0^2 \int_{\sqrt{y/2}}^y dx dy = \int_0^2 (x) \Big|_{\sqrt{y/2}}^{\sqrt{y/2}} dy$$

$$= \int_0^2 \frac{\sqrt{y}}{\sqrt{2}} - \frac{y^2}{4} dy$$

$$= \left[\frac{y^{3/2}}{\sqrt{2} \cdot 3/2} - \frac{y^3}{3 \times 4} \right]_0^2$$

$$= \frac{2^{3/2}}{\sqrt{2} \cdot 3/2} - \frac{2^3}{3 \times 4}$$

$$= \frac{2\sqrt{2} \times 2}{3\sqrt{2}} - \frac{8}{3 \times 4}$$

$$= \frac{4}{3} - \frac{2}{3}$$

$$= \frac{2}{3}$$

Type: 2 (Polar co-ordinates).

$$\text{Area} = \int_{\theta_1}^{\theta_2} \int_{r=f_1(\theta)}^{r=f_2(\theta)} r dr d\theta$$

1) Find the area of the cardioid $r = a(1 + \cos \theta)$

Ans: $r = a(1 + \cos \theta)$

It is symmetrical about the initial line.

$$\theta: 0 \rightarrow \pi$$

$$r: 0 \rightarrow a(1 + \cos \theta)$$

$$\text{Area} = 2 \int_0^\pi \int_0^{a(1+\cos \theta)} r \, dr \, d\theta$$

$$= 2 \int_0^\pi \left(\frac{r^2}{2} \right)_0^{a(1+\cos \theta)} d\theta$$

$$= \frac{2}{2} \int_0^\pi [a(1 + \cos \theta)]^2 d\theta$$

$$= a^2 \int_0^\pi \{1 + \cos^2 \theta + 2\cos \theta\} d\theta$$


$$= a^2 \int_0^\pi \left\{ 1 + \frac{1 + \cos 2\theta}{2} + 2\cos \theta \right\} d\theta$$

$$= \frac{a^2}{2} \int_0^\pi \{3 + \cos 2\theta + 4\cos \theta\} d\theta$$

$$= \frac{a^2}{2} \left\{ 3\theta + \frac{\sin 2\theta}{2} + 4\sin \theta \right\}_0^\pi$$

$$= \frac{a^2}{2} \left\{ (3\pi + \frac{\sin 2\pi}{2} + 4\sin \pi) - (0 + \sin 0 + 4\sin 0) \right\}$$


$$= \frac{a^2}{2} (3\pi) = \frac{3}{2} \pi a^2$$



2) Find the area of the cardioid $r = a(1 - \cos \theta)$

$$\text{Area} = 2 \int_0^\pi \int_0^{a(1-\cos \theta)} r \, dr \, d\theta$$

$$= \frac{3}{2} \pi a^2$$



Scanned with CamScanner

3) Find the area of the lemniscate $r^2 = a^2 \cos 2\theta$ by double integration.

Ans Given $r^2 = a^2 \cos 2\theta \Rightarrow r = a\sqrt{\cos 2\theta}$

$$\theta: 0 \rightarrow \pi/4$$

$$r: 0 \rightarrow a\sqrt{\cos 2\theta}$$

$$\text{Area} = 4 \int_0^{\pi/4} \int_0^{a\sqrt{\cos 2\theta}} r \, dr \, d\theta$$


$$= 4 \int_0^{\pi/4} \left. \frac{r^2}{2} \right|_0^{a\sqrt{\cos 2\theta}} d\theta$$

$$= \frac{4}{2} \int_0^{\pi/4} (a\sqrt{\cos 2\theta})^2 d\theta$$

$$= 2 \int_0^{\pi/4} a^2 \cos 2\theta \, d\theta$$


$$= 2 \left\{ a^2 \frac{\sin 2\theta}{2} \right\}_0^{\pi/4}$$

$$= \frac{2a^2}{2} \{1 - 0\} = a^2$$



4) Find the area that lies inside the cardioid $r = a(1 + \cos \theta)$ & outside the circle $r = a$ by double integration.

Ans Given $r = a(1 + \cos \theta)$, $r = a$



$$\theta: 0 \rightarrow \pi/2$$

$$r: a \rightarrow a(1 + \cos \theta)$$

$$\text{Area} = 2 \int_0^{\pi/2} \int_a^{a(1 + \cos \theta)} r \, dr \, d\theta$$

$$= 2 \int_0^{\pi/2} \left(\frac{r^2}{2} \right)_a^{a(1+\cos\theta)} d\theta$$

$$= \int_0^{\pi/2} (a^2(1+\cos\theta)^2 - a^2) d\theta$$

$$= a^2 \int_0^{\pi/2} \{1 + \cos^2\theta + 2\cos\theta - 1\} d\theta$$

$$= a^2 \int_0^{\pi/2} \left\{ 1 + \frac{\cos 2\theta}{2} + 2\cos\theta \right\} d\theta$$

$$= \frac{a^2}{2} \int_0^{\pi/2} (1 + \cos 2\theta + 4\cos\theta) d\theta$$

$$= \frac{a^2}{2} \left\{ \theta + \frac{\sin 2\theta}{2} + 4\sin\theta \right\}_0^{\pi/2}$$

$$= \frac{a^2}{2} \left\{ \pi/2 + \frac{\sin \pi}{2} + 4\sin \pi/2 - (0+0+0) \right\}$$

$$= \frac{a^2}{2} \{ \pi/2 + 4 \}$$

$$= \frac{a^2}{4} \{ \pi + 8 \}$$

- 5) Find the area that lies inside the cardioid $r = 2(1 + \cos\theta)$ & outside the circle $r = 2$ by double integration.

Ans Same as problem 4, Replace 'a' by 2 we get area = $\pi + 8$

Triple Integrals

1) Evaluate $\int_0^a \int_0^b \int_0^c xyz \, dz \, dy \, dx$

$$= \left(\frac{z^2}{2} \right)_0^c \left(\frac{y^2}{2} \right)_0^b \left(\frac{x^2}{2} \right)_0^a$$

$$= \frac{a^2 b^2 c^2}{8}$$

2) Evaluate $\int_0^a \int_0^b \int_0^c e^{x+y+z} dz dy dx$

Ans
$$I = \int_0^a e^x dx \int_0^b e^y dy \int_0^c e^z dz$$

$$= (e^x)_0^a (e^y)_0^b (e^z)_0^c$$

$$= (e^a - 1) (e^b - 1) (e^c - 1)$$

3) Evaluate $\int_0^1 \int_0^2 \int_0^3 xyz dz dy dx$

Ans
$$I = \int_0^1 x dx \int_0^2 y dy \int_0^3 z dz$$

$$= \left(\frac{x^2}{2}\right)_0^1 \left(\frac{y^2}{2}\right)_0^2 \left(\frac{z^2}{2}\right)_0^3$$

$$= \frac{1}{2} \cdot \frac{2^2}{2} \cdot \frac{3^2}{2}$$

$$= \frac{36}{8} = \frac{9}{2}$$

4) Evaluate $\int_0^a \int_0^b \int_0^c (x^2 + y^2 + z^2) dx dy dz$

$$I = \int_0^a \int_0^b \left(\frac{x^3}{3} + y^2 x + x z^2\right)_0^c dy dz$$

$$= \int_0^a \int_0^b \left(\frac{c^3}{3} + cy^2 + cz^2\right) dy dz$$

$$= \int_0^a \left(\frac{c^3 y}{3} + \frac{cy^2}{2} + cz^2 y\right)_0^b dz$$

$$= \int_0^a \left\{ \frac{c^3 b}{3} + \frac{cb^2}{2} z + \frac{cbz^3}{3} \right\}_0^a dz$$

$$= \frac{c^3 ba}{3} + \frac{cb^2 a^2}{2} + \frac{cb a^3}{3}$$

$$= \frac{abc}{3} \{a^2 + b^2 + c^2\}$$

5) Evaluate $\int_0^{\log a} \int_0^x \int_0^{x+y} e^{x+y+m} dm dy dx$

$$I = \int_0^{\log a} \int_0^x e^x e^y (e^m)_0^{x+y} dy dx$$

$$= \int_0^{\log a} \int_0^x e^x e^y (e^{x+y} - 1) dy dx$$

$$= \int_0^{\log a} \int_0^x (e^{2x} e^{2y} - e^x e^y) dy dx$$

$$= \int_0^{\log a} \left\{ e^{2x} \frac{e^{2y}}{2} - e^x e^y \right\}_0^x dx$$

$$= \int_0^{\log a} \left[\frac{e^{4x}}{2} - e^{2x} \right] - \left[\frac{e^{2x}}{2} - e^x \right] dx$$

$$= \int_0^{\log a} \frac{e^{4x}}{2} - \frac{3e^{2x}}{2} + e^x dx$$

$$= \left[\frac{e^{4x}}{8} - \frac{3e^{2x}}{4} + e^x \right]_0^{\log a}$$

$$= \frac{e^{4 \log a}}{8} - \frac{3e^{2 \log a}}{4} + e^{\log a} - \left(\frac{1}{8} - \frac{3}{4} + 1 \right)$$

$$= \left(\frac{a^4}{8} - \frac{3a^2}{4} + a \right) - \left(\frac{1-6+8}{8} \right)$$

$$= \frac{a^4 - 6a^2 + 8a}{8} - \left(\frac{3}{8} \right)$$

$$= \frac{1}{8} (a^4 - 6a^2 + 8a - 3)$$

Volume of solids

$$\text{Volume} = \iiint_R dx dy dz$$

1) Find the volume of the sphere $x^2 + y^2 + z^2 = a^2$ without transformation.

Ans Given $x^2 + y^2 + z^2 = a^2$

Volume of the sphere = 8 x volume in the I octant

$$= 8 \iiint dx dy dz$$

Here $z: 0 \rightarrow \sqrt{a^2 - x^2 - y^2}$

$y: 0 \rightarrow \sqrt{a^2 - x^2}$

$x: 0 \rightarrow a$

$$= 8 \int_0^a \int_0^{\sqrt{a^2 - x^2}} \int_0^{\sqrt{a^2 - x^2 - y^2}} dz dy dx$$

$$= 8 \int_0^a \int_0^{\sqrt{a^2 - x^2}} (z)_0^{\sqrt{a^2 - x^2 - y^2}} dy dx$$

$$= 8 \int_0^a \int_0^{\sqrt{a^2 - x^2}} \sqrt{a^2 - x^2 - y^2} dy dx$$

$$= 8 \int_0^a \left[\frac{y}{2} \sqrt{a^2 - x^2 - y^2} + \left(\frac{a^2 - x^2}{2} \right) \sin^{-1} \left\{ \frac{y}{\sqrt{a^2 - x^2}} \right\} \right]_0^{\sqrt{a^2 - x^2}} dx$$

$$= 8 \int_0^a \left[0 + \frac{a^2 - x^2}{2} \sin^{-1}(1) + 0 \right] dx$$

$$= 8 \int_0^a \frac{a^2 - x^2}{2} \cdot \frac{\pi}{2} dx$$

$$= 2\pi \left\{ a^2 x - \frac{x^3}{3} \right\}_0^a$$

$$= 2\pi \left\{ a^3 - \frac{a^3}{3} \right\}$$

$$= \frac{4}{3} \pi a^3$$

2) Find the Volume of the tetrahedron bounded by the planes $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$ and the co-ordinates planes (A.U.)

Ans

The region bounded by $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$, $x=0$, $y=0$ & $z=0$.

Limits

$$x: 0 \rightarrow a$$

$$y: 0 \rightarrow b(1 - \frac{x}{a})$$

$$z: 0 \rightarrow c[1 - \frac{x}{a} - \frac{y}{b}]$$

$$\begin{aligned} \therefore \text{Volume} &= \iiint dz dy dx \\ &= \int_0^a \int_0^{b(1-\frac{x}{a})} \int_0^{c(1-\frac{x}{a}-\frac{y}{b})} dz dy dx \\ &= \int_0^a \int_0^{b(1-\frac{x}{a})} c(1-\frac{x}{a}-\frac{y}{b}) dy dx \\ &= c \int_0^a \left\{ y - \frac{xy}{a} - \frac{y^2}{2b} \right\}_0^{b(1-\frac{x}{a})} dx \\ &= c \int_0^a \left\{ b(1-\frac{x}{a}) - \frac{xb(1-\frac{x}{a})}{a} - \frac{b^2(1-\frac{x}{a})^2}{2b} \right\} dx \\ &= c \int_0^a b(1-\frac{x}{a}) \left\{ (1-\frac{x}{a}) - \frac{(1-\frac{x}{a})}{2} \right\} dx \\ &= bc \int_0^a (1-\frac{x}{a})^2 dx \\ &= \frac{bc}{2} \left\{ \frac{(1-\frac{x}{a})^3}{3(-\frac{1}{a})} \right\}_0^a \\ &= \frac{bc}{2} (-a)(0-1) = \frac{abc}{6} \end{aligned}$$

3) Find the Volume of the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ which lies in the I octant

Ans:

Volume of the ellipsoid in the I octant is bounded by the planes $x=0, y=0, z=0$ & the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$

Limits

$$x: 0 \rightarrow a$$

$$y: 0 \rightarrow b\sqrt{1-\frac{x^2}{a^2}}$$

$$z: 0 \rightarrow c\sqrt{1-\frac{x^2}{a^2}-\frac{y^2}{b^2}}$$

$\therefore$ Volume in I octant

$$= \int_0^a \int_0^{b\sqrt{1-\frac{x^2}{a^2}}} \int_0^{c\sqrt{1-\frac{x^2}{a^2}-\frac{y^2}{b^2}}} dz dy dx$$

$$= \int_0^a \int_0^{b\sqrt{1-\frac{x^2}{a^2}}} (z)_0^{c\sqrt{1-\frac{x^2}{a^2}-\frac{y^2}{b^2}}} dy dx$$

$$= c \int_0^a \int_0^{b\sqrt{1-\frac{x^2}{a^2}}} \frac{1}{b} \sqrt{b^2(1-\frac{x^2}{a^2})-y^2} dy dx$$

$$= \frac{c}{b} \int_0^a \left\{ \frac{y}{2} \sqrt{b^2(1-\frac{x^2}{a^2})-y^2} + \frac{b^2(1-\frac{x^2}{a^2})}{2} \sin^{-1} \left\{ \frac{y}{b\sqrt{1-\frac{x^2}{a^2}}} \right\} \right\}_0^{b\sqrt{1-\frac{x^2}{a^2}}} dx$$

$$= \frac{c}{2b} \int_0^a b^2 \left(1-\frac{x^2}{a^2}\right) \pi/2 dx$$

$$= \frac{\pi c}{4b} b^2 \left\{ x - \frac{x^3}{3a^2} \right\}_0^a = \frac{\pi b c}{4} \frac{2a^3}{3a^2}$$

$$= \frac{\pi abc}{6}$$

Change of variables

1) change into polar co-ordinates & hence evaluate

$$\int_0^a \int_0^{\sqrt{a^2-x^2}} \sqrt{x^2+y^2} \, dy \, dx$$

Ans $y: 0 \rightarrow \sqrt{a^2-x^2}$

$$y=0 \rightarrow y=\sqrt{a^2-x^2}$$

$$x^2+y^2=a^2$$

Circle with $(= (0,0))$
& radi = a

Also $x: 0 \rightarrow a$

put $x=r \cos \theta$, $y=r \sin \theta$
 $dx dy = r \, dr \, d\theta$

$$\therefore I = \int_0^{\pi/2} \int_0^a \sqrt{r^2} \, r \, dr \, d\theta$$



$$= \int_0^{\pi/2} \int_0^a r^2 \, dr \, d\theta$$

$$= \int_0^{\pi/2} \left(\frac{r^3}{3} \right)_0^a d\theta$$

$$= \frac{1}{3} \int_0^{\pi/2} a^3 \, d\theta$$

$$= \frac{a^3}{3} (\theta)_0^{\pi/2}$$

$$= \frac{a^3}{3} (\pi/2) = \frac{a^3 \pi}{6}$$



2) By changing into polar co-ordinates, evaluate

$$\int_0^2 \int_0^{\sqrt{2x-x^2}} \frac{x}{x^2+y^2} dx dy$$

Ans

$$I = \int_0^2 \int_0^{\sqrt{2x-x^2}} \frac{x}{x^2+y^2} dy dx \quad (\text{correct form})$$

$$y: 0 \rightarrow \sqrt{2x-x^2}$$

$$y=0 \rightarrow y=\sqrt{2x-x^2}$$

$$y^2 = 2x - x^2$$

$$x^2 + y^2 - 2x = 0$$

Circle with

$$C = (1, 0)$$

& radi = 1

$$\text{Also } y \text{ } x: 0 \rightarrow 2$$

$$x=0 \rightarrow x=2$$

$$\therefore I = \int_0^{\pi/2} \int_0^{2\cos\theta} \frac{x \cos\theta}{r^2} r dr d\theta$$


$$= \int_0^{\pi/2} \cos\theta (r)_0^{2\cos\theta} d\theta$$

$$= \int_0^{\pi/2} \cos\theta (2\cos\theta) d\theta$$

$$= 2 \int_0^{\pi/2} \frac{1+\cos 2\theta}{2} d\theta$$

$$= \frac{2}{2} \left\{ \theta + \frac{\sin 2\theta}{2} \right\}_0^{\pi/2}$$

$$= \pi/2$$



$$\text{Take } x = r \cos\theta$$


$$y = r \sin\theta$$

$$dx dy = r dr d\theta$$

Applications:

- 1) Find the mass and centre of mass of the lamina that occupies the region D and has the density function ρ . D is bounded by $y = x^2$ & $y = x + 2$, $\rho(x, y) = kx$

Ans:



we know that, the mass of homogeneous lamina is given by

$$m = \iint_R \rho(x, y) dA$$

$$m = \int_{-1}^2 \int_{x^2}^{x+2} kx dy dx$$

$$= \int_{-1}^2 kx (x - x^2 + 2) dx$$

$$= k \left(\frac{x^3}{3} - \frac{x^4}{4} + \frac{2x^2}{2} \right) \Big|_{-1}^2$$

$$= k \left(\frac{8}{3} - \frac{16}{4} + 4 \right) - \left(-\frac{1}{3} - \frac{1}{4} + 1 \right)$$

$$= k \left(\frac{32}{12} - \frac{15}{12} \right) = k \left(\frac{27}{12} \right) = \frac{9}{4} k$$

$$\bar{x} = \frac{1}{m} \iint_R x \rho(x, y) dA$$

$$= \frac{1}{\left(\frac{9}{4} k \right)} \int_{-1}^2 \int_{x^2}^{x+2} kx^2 dy dx$$

$$= 7/5$$

$$\bar{y} = \frac{1}{m} \iint_R y \rho(x, y) dA$$

$$= \frac{1}{\left(\frac{9}{4}k\right)} \int_{-1}^2 \int_{x^2}^{x+2} kxy dy dx$$

$$= \frac{2}{9} \int_{-1}^2 (x^3 + 4x + 4x^2 - x^5) dx$$


$$= \frac{2}{9} \left\{ 12 - \frac{6}{8} \right\}$$

$$= 5/2$$

$\therefore$ Centre of gravity of the lamina is $\left(\frac{7}{5}, \frac{5}{2}\right)$

- 2) Find the mass and centre of mass of the lamina that occupies the region D and has the given density function ρ . D is bounded by the parabolas $y = x^2$, $x = y^2$; $\rho(x, y) = \sqrt{x}$.

Ans



$$m = \int_0^1 \int_{x^2}^{\sqrt{x}} \sqrt{x} dy dx = \int_0^1 \sqrt{x} (y)_{x^2}^{\sqrt{x}} dx$$

$$= \int_0^1 \sqrt{x} (\sqrt{x} - x^2) dx$$

$$= \int_0^1 x - x^{5/2} dx$$

$$= \left(\frac{x^2}{2} - \frac{2}{7} x^{7/2} \right) \Big|_0^1 = 3/14$$

Scanned with CamScanner

